



# House dust mites in three contrasting climatic regions of Saudi Arabia

Riyadh Hussain<sup>1,2</sup> · S. Aeban<sup>1,2</sup> · Medjedline Hani<sup>1</sup> · Henk R. Braig<sup>3</sup> · M. Alejandra Perotti<sup>1</sup>

Received: 13 October 2025 / Accepted: 19 February 2026  
© The Author(s) 2026

## Abstract

Just a handful of mite species are suspected inhabiting the dust of houses within arid areas of the Arabian Peninsula. In this study, we identified and compared the house dust mite diversity in three different climatic areas within Saudi Arabia. Sampling was carried out over one year and two months, with samples collected in the desert, the coast, and in the highlands. It reviews all known mites of house dust of Saudi Arabia. A total of 998 mites were collected belonging to 23 species in 14 families. Pyroglyphidae (Astigmata) were the most abundant. The interaction among pyroglyphid species in the climatic regions was analysed. While variation in abundance of Pyroglyphidae among regions was weak, there was a strong desert region *Cheyletus eruditus*–*Dermatophagoides farinae* association, perhaps explaining that *D. farinae* populations from the desert have become *C. eruditus*' favourite prey. Another interesting association is that of Pyroglyphidae with Acaridae. *D. pteronyssinus* shows significant numerical dominance in the highlands. A newly discovered species of *Dermatophagoides* did not interact with any of the other 2 species and was restricted to the coastal region. *Dermatophagoides* n. sp. and *Cheletopsis* sp. can be used as markers of location in forensic analyses (provenance), based on their unique association with the coast and desert, respectively. A new species of *Dermatophagoides* has been identified. Five species are new to Saudi Arabia and the Arabian Peninsula including house dust, ten species are new to the house dust mite fauna of Saudi Arabia and the Arabian Peninsula but have been recorded as storage mites, totalling 15 new house dust mites for the Arabian Peninsula. Two species are new to house dust in Saudi Arabia, but already known from house dust in Kuwait. Of the fifteen new house dust mites for the Arabian Peninsula, five species are new to the dust mite fauna worldwide: Prostigmata: *Cheletopsis* sp. (Cheyletidae); Mesostigmata: *Proctolaelaps pygmaeus* (Melicharidae), *Macrocheles muscaedomesticae* (Macrochelidae); and Oribatida: *Oribatula tibialis* (Oribatulidae), and *Ceratozetes* sp. (Ceratozetidae).

**Keywords** Acari · House dust mites · Allergens · Biodiversity · Pyroglyphidae · Cheyletidae · *Dermatophagoides* · Oribatida · HDM · Arabian Peninsula

---

Henk R. Braig and M. Alejandra Perotti contributed equally.

---

Extended author information available on the last page of the article

## Introduction

Domestic mites (Acari) inhabit human dwellings, predominantly as house dust or pulvicolous mites, as stored products mites, and as mites of pet animals and house mice (Ogata et al. 1980; Faccini and Coutinho 1987; Hagstrum et al. 2013b; Abdel-Rahman et al. 2020; Kontschán et al. 2022; Kamezaxi et al. 2024; Kontschán and Hornok 2025; Yu et al. 2025). The mites associated with human habitation are being studied extensively and from almost every corner of the world with recent additions from, for example, Egypt (Bakr et al. 2022; Ibrahim et al. 2022), India (Saw et al. 2024), Iran (Zare et al. 2021), Ireland (Aljohani et al. 2024), Japan (Kamezaki et al. 2022), Panama (Lezcano et al. 2020), the Philippines (Corpuz-Raros 2022), Poland (Solarz et al. 2021), or Türkiye (Akdemir et al. 2023; Aykut 2024). House dust mites belonging mainly to the family Pyroglyphidae are feeding on mould (fungi) and, among other things, on human and pet animal skin flakes. The most ubiquitous species are *Dermatophagoides farinae* Hughes, 1961, *D. pteronyssinus* (Trouessart, 1897), *D. evansi* Fain, Hughes & Johnston, 1967, and *Euroglyphus maynei* (Cooreman, 1950). Storage mites include mites that feed on the stored products such as grain, and fungivore and other detritivore mites. The most prominent families are Acaridae, Echymopodidae, Glycyphagidae, Chortoglyphidae and Suidassidae represented by *Acarus siro* Linnaeus, 1758, *Thyreophagus entomophagus* (Laboulbène & Robin, 1862), *Tyrophagus putrescentiae* (Schrank 1781); *Blomia tropicalis* Van Bronswijk, De Cock and Oshima 1973; *Glycyphagus domesticus* (de Geer, 1778), *Lepidoglyphus destructor* (Schrank 1781); *Chortoglyphus arcuatus* (Troupeau, 1879); *Suidasia medanensis* Oudemans, 1924, and *S. pontifica* Oudemans, 1905 (Wharton 1976; Malainual et al. 1995; Petrova and Zheltikova 2000; Takaoka 2000; Pike and Wickens 2008; Yu et al. 2015; Podder et al. 2021; Solarz et al. 2021; Acuna-Cantillo et al. 2024; Aykut 2024). The concentration of storage mites might be underestimated compared to house dust mites (Reboux et al. 2019). There is some overlap between the two groupings, for example, *B. tropicalis* is often also considered a house dust mite. In animal studies, storage mites might produce greater inflammation of the lungs than house dust mites (Kim et al. 2024). Dust mites only by appearance, which are not associated with houses or storage such as date palm dust mite, *Oligonychus afrasiaticus* (McGregor, 1939), are not covered here.

The mite fauna of indoors can be quite rich. Oribatida (soil mites) can be brought inside houses (Chen et al. 2023), members of the Tetranychidae (Prostigmata) can also be found associated with ornamental plants. All these mites attract their natural predators to come indoors as well, particularly from the family Cheyletidae with *Cheyletus malaccensis* (Oudemans, 1903) and *C. eruditus* (Schrank 1781) being most widely distributed in houses (Zhou et al. 2022). Ticks only occur together with their household pets, the study of their development under domestic conditions has just begun (Buczek et al. 2023; Shinohara et al. 2023).

A new build is practically free of mites (Ouchi et al. 1976; Crowther et al. 2000). Humans introduce house dust mites mostly with their clothing (Perotti and Braig 2009; Clarke et al. 2015). When houses become uninhabited, the dust mites start to disappear too but are replaced by other species like *Tyrophagus putrescentiae*, species of Glycyphagidae, Tarsosomidae, Cheyletidae and Oribatida, as this has been seen in houses uninhabited for 6–7 years in Fukushima, Japan; where the total amount of mites stayed roughly the same (Shi-

nohara et al. 2023). The species more dominant in uninhabited houses are known from dust of houses built in the past more open to the environment.

Domestic mites may pose health risks (Cui 2014; Hubert et al. 2018). Most dust mites are allergenic, but only a fraction of the 280 or so storage mite species are major sources of allergens (Hagstrum et al. 2013a; Fernández-Caldas et al. 2014). Micro- and nanoplastics increase the allergenic potential of mite allergens (Wang et al. 2025; Wu and Yang 2025). The indoor mite fauna is also of increasing forensic interest (Perotti et al. 2009; Solarz 2009; Frost et al. 2010; Çakan et al. 2015; Perotti and Braig 2019; Solarz et al. 2022).

The fast adaptation to the peculiarities of human household environments is reflected in the genomes of house dust and storage mites (Xiong et al. 2022; Hubert et al. 2023; Klimov et al. 2024; Vidal-Quist et al. 2025). Humidity and temperature are crucial parameters in the distribution of domestic mites. House dust mites mainly absorb water from humid air. The active breeding zone has been defined for a relative humidity of 55 to 75% at 15 to 35 °C. (Arlian 1992; Pei et al. 2020; Vackova et al. 2021, 2023). Eggs of *D. pteronyssinus* withstand for 80% 40 °C dry heat, but for direct sunlight and 50 °C dry heat, the thermal death point occurs at 3 and 5 h, and for 60 and 70 °C, at 30 min (Mahakittikun et al. 2011). This impacts domestic mites in desertic regions like the Arabian Peninsula.


In this work we were interested in exploring the diversity and abundance of the domestic mite fauna in major Saudi cities in three ecoregions of the Arabian Peninsula, the desert and xeric shrublands with Riyadh and Al-Kharj, the coastal fog desert with Jeddah, and the highlands which themselves can be subdivided in the foothills savanna with Shafa and the montane woodlands with Taif (Fig. 1).

The importance of house dust mites for producing aeroallergens in respiratory diseases and disorders such as allergic rhinitis, asthma, asthmatic flare-ups, cat allergen sensitization and atopic dermatitis is often high or even the highest across the climate ranges of Saudi Arabia and the Arabian Peninsula (Table 1). This importance is measured against other indoor allergens like cockroaches and pet dander (cats, dogs), and outdoor allergens such as plants (prickly saltwort, lamb's quarters, ragweed, pigweed, Russian thistle), grasses (Bermuda grass, Timothy grass, rye), trees (several mesquites), and moulds, *Alternaria* and *Aspergillus* species. These studies on aeroallergens are mainly based on *Dermatophagoides farinae* and *D. pteronyssinus*, because most commercial test kits used in literature cover only these two mite species (Khurram et al. 2025). The diversity of the actual house dust mites has been little researched. Studies have been limited to two locations on the coast only, Dammam (Al-Qurashi 2006) and Jeddah (Edrees 2008, 2009, 2010, 2012). There are no records of house dust mites in the desert regions or in the highlands. In this work we explored the diversity and abundance of the House Dust Mite (HDM) fauna in three climatic regions of Saudi Arabia.

## Methods

### General information and sampling

House dust samples were collected over a period of one year and two months, starting in December 2016 and finishing in January 2018. The dust samples were provided by 5 households, each of them from each of the 5 Saudi cities: Al-Kharj, Jeddah, Riyadh, Shafa, and



**Fig. 1** Five cities in the three climatic regions, the Desert composed of the Arabian desert and East Sahero-Arabian xeric shrublands belonging to the Palearctic realm; the Coast being the Arabian Peninsula coastal fog desert; and the Highlands made up of the Southwestern Arabian foothills savanna, and the Southwestern Arabian montane woodlands (mountain woodlands), all belonging to the Afrotropical realm

Taif; therefore, a total of 25 sampling sites. For *Dermatophagoides* spp. repeated samplings were conducted (x2) totalling 50 samples (Table S2); a 1st sampling in 2017 was only on *Dermatophagoides* spp while 2nd sampling attempt in 2018 included all mites.

**Desert:** The city of Riyadh,  $24^{\circ}38'N$   $46^{\circ}43'E$ , is at an elevation of 600 m situated in the an-Nafud desert on the eastern part of the Najd plateau, and Al-Kharj ( $24^{\circ}08'54.0''N$   $47^{\circ}18'18.0''E$ ) is an oasis southeast of Riyadh, both in Riyadh Province. Climate is hot dry with desert subzone. They belong to the Arabian desert in the Palearctic realm. Low relative humidity with a trough as low as 10% in June and a peak of up to 50% in December to January (Meteo Portal 2017).

**Coast:** Jeddah,  $21^{\circ}32'36''N$   $39^{\circ}10'22''E$ , Makkah Province, is a port city at the Red Sea. Climate is hot dry maritime desert. It belongs to the tip of the Northern tip of Arabian Peninsula coastal fog desert or xeric shrub ecoregion in the Afrotropical realm although it does not experience so much fog. Relative humidity around Jeddah can be very high, fluctuating between low 40s in May to July to over 80% in October.

**Highlands:** Ash-Shafa,  $21^{\circ}04'10''N$   $40^{\circ}18'43''E$ , is at an elevation of 2,240 m and Taif,  $21^{\circ}16'30.34''N$   $40^{\circ}24'22.16''E$ , is at 1,880 m are on the slopes of the Hijaz Mountains that belong to the Sarat Mountains; they both are also in Makkah Province. These two cities are close together, yet, Shafa, despite its elevation, is considered part of the foothills savanna while Taif is part of the montane woodlands ecoregion in the Afrotropical realm. Climate is sub-tropical with a Mediterranean subzone and a mountainous subtype. Relative humidity ranges between 20% in June and 60% in January.

**Table 1** Importance of house dust mite allergens in Saudi Arabia and in the Arabian Peninsula

Condition	Importance	Location
Allergic rhinitis		
Coast	high	Western SA (Badran et al. 2016); Bahrain (Hasan and Rizwan 2015) Oman (Al-Tamemi et al. 2008; Al-Abri et al. 2015); Qatar (Sattar et al. 2003; Zahraldin et al. 2021; Thalappil et al. 2023); UAE (Sultan and Khalil 2012; Mahboub et al. 2014; Asad et al. 2015), children (Adham and Tawfik 2012; Al Sharif et al. 2017; Sharif et al. 2018)
Desert	low to moderate	Jordan (Aburuz et al. 2011), Kuwait (Dowaisan et al. 2000; Ezeamuzie et al. 2000a; Al-Dowaisan et al. 2004); children (al Mousawi et al. 2009; Abal et al. 2010), UAE (Bener et al. 2002)
	high	Riyadh (Almogren 2009)
	low to moderate	Riyadh (Al-Shalan et al. 1989; Zakzouk and Gad-El-Rab 1996), Najran (Alqahtani 2020), children (Alqahtani 2016); Qassim (Alra-sheedi et al. 2023a)
Highland	high	Abda children (Asseri et al. 2025)
	moderate	Yemen (Al-Mehdar et al. 2025)
	low	Jazan (Zaki et al. 2019)
	highest	Abha (Al-Ghamdi et al. 2022)
Nation		Saudi Arabia (Emara et al. 2012; Almehezia et al. 2019)
Asthma		
Coast	high	Jazan (Alhazmi et al. 2025); Jeddah (Al-Frayh et al. 1997; Koshak 2006), Al-Khobar (Al-Nahdi and Al-Quorain 1987), children (Al Khater 2017), Makkah (Tayeb 2015), children (Al-Frayh et al. 1992); Oman (Al-Tamemi et al. 2008); Qatar (Sattar et al. 2003; Zahraldin et al. 2021; Thalappil et al. 2023); UAE (Sultan and Khalil 2012; Asad et al. 2015), children (Al-Aani and Alhammadi 2015; Al Sharif et al. 2017; Sharif et al. 2018)
	low to moderate	Kuwait (Ezeamuzie et al. 2000a, b; Khadadah et al. 2000a, b; al Mousawi et al. 2001); children (Hijazi et al. 2002; al Mousawi et al. 2009; Abal et al. 2010); UAE (Bener et al. 2002)
Desert	high	Riyadh (Almogren 2009)
	low to moderate	Riyadh (Al-Frayh et al. 1997; Gad-El-Rab 1999), children (Al-Frayh et al. 1992); Najran (Alqahtani 2020), children (Alqahtani 2016)

Table 1 (continued)

Condition	Importance	Location
Highland	high	Abhar (Al-Frayh et al. 1997), Aseer (Al-Ghamdi et al. 2019), Jazan (Zaki et al. 2019) Yemen (Al-Mehdar et al. 2025)
Nation	moderate	Tabuk (Al-Dusari et al. 1997)
Asthma flair ups	highest	Saudi Arabia (Emara et al. 2012)
Desert	highest	Qassim (Alrasheedi et al. 2023b)
Cat allergen sensitization	high	Jeddah (Tayeb 2020)
Coast	high	
Atopic dermatitis		
Coast	highest	UAE children (Adham et al. 2011)
Desert	little	Najran (Alqahtani 2016, 2020)
Highland	moderate	Tabuk (Al-Dusari et al. 1997)

**Table 2** Mite allergens and mites in the Arabian Peninsula

	Mite allergens in literature Saudi Arabia [% of sensitizations]a	Mite allergens in literature Arabian Peninsula [% of sensitizations]a	# Mites collected here in Saudi Arabia 1st - 2nd*
<b>Dermatophagoides</b>			
<i>D. farinae</i>	84% (Koshak 2006), 56% (Al-Frayh et al. 1997), 48% (Tayeb 2020), 36% (Hassan et al. 2013), 33% (Badran et al. 2016), 31% (Hasnain et al. 2012), 29% (Hasnain et al. 2012), 11% (Tayeb 2015); children 50% (Al-Frayh et al. 1992), 48% (Al Khater 2017).	Bahrain (Hasan and Rizwan 2015) Jordan 22% (Aburuz et al. 2011), 11% (Khasawneh et al. 2019), 8% (Al-Zayadneh et al. 2019), 2% (Khreesha et al. 2020); children 88% (Khreesha et al. 2020). Kuwait 39% (Ezeamuzie et al. 2000b), 32% (Dowaisan et al. 2000), 30% (Al-Dowaisan et al. 2004), 16% (Tubailah and Luqman 1987). Oman 48% (Al-Tamemi et al. 2008). Qatar 39% (Thalappil et al. 2023), 37% (Sattar et al. 2003), children 29% (Zahraidin et al. 2021). UAE 42% (Hasnain et al. 2012), children and adults (Yousif et al. 2025); children 89% (Adham and Tawfik 2012), 75% (Adham et al. 2011), 37% (Al Sharif et al. 2017; Sharif et al. 2018), 18% (Lestringant et al. 1999), 16% (Al-Aani and Alhammadi 2015)	19-43
Coast			
Desert	38% (Hasnain et al. 2012), 13% (Al-Frayh et al. 1997), 20% (Al-Shalan et al. 1989), 11% (Zakzouk and Gad-El-Rab 1996), 7% (Alqahtani 2020); 10% children (Al-Frayh et al. 1992; Alqahtani 2016).		5-11
Highlands	40% (Hasnain et al. 2012) 19% (Al-Frayh et al. 1997), 16% (Al-Ghamdi et al. 2019), 12% (Al-Ghamdi et al. 2022).	Yemen children and adults 10% (Al-Mehdar et al. 2025)	33-90
<b>D. microceras</b>			
Coast		Kuwait 45% (Ezeamuzie et al. 2000b), 39% (Dowaisan et al. 2000).	-
<b>D. pteronyssinus</b>			

Table 2 (continued)

	Mite allergens in literature Saudi Arabia [% of sensitizations]a	Mite allergens in literature Arabian Peninsula [% of sensitizations]a	# Mites collected here in Saudi Arabia 1st - 2nd*
Coast	87% (Koshak 2006), 47% (Tayeb 2020), 39% (Hasnain et al. 2012), 37% (Badran et al. 2016), 36% (Sullaman et al. 1997), 28% (Hasnain et al. 2012), 20% (Hassan et al. 2013), 18% (Tayeb 2015); children 45% (Al Khater 2017).	Bahrain 42% (Hasan and Rizwan 2015) Jordan 33% (Aburuz et al. 2011), 18% (Khreesha et al. 2020), 14% (Khasawneh et al. 2019), 12% (Al-Zayadneh et al. 2019), children 66% (Khreesha et al. 2020). Kuwait 54% (Hijazi et al. 2002), 53% (Ezeamuzie et al. 1997), 46% (Ezeamuzie et al. 2000b), 38% (Ezeamuzie et al. 2000a; Al-Dowaisan et al. 2004), 36% (Dowaisan et al. 2000), 29% (Khadadah et al. 2000b), 28% (Khadadah et al. 2000a), 23% (Ezeamuzie et al. 1997), 20% (Tubailah and Luqman 1987), children 4% (al Mousawi et al. 2009). Oman 51% (Al-Tamemi et al. 2008). Qatar 50% (Thalappil et al. 2023), 42% (Sattar et al. 2003); children 38% (Zahraidin et al. 2021). UAE 46% (Hasnain et al. 2012), children and adults (Yousif et al. 2025), children 96% (Adham and Tawfik 2012), 75% (Adham et al. 2011), 37% (Al Sharif et al. 2017; Sharif et al. 2018), 18% (Lestringant et al. 1999), 14% (Al-Aani and Alhammedi 2015).	24-28
Desert	37% (Hasnain et al. 2012), 13% (Alqahtani 2020); children 15% (Alqahtani 2016).		5-7
Highlands	40% (Hasnain et al. 2012), 25% (Al-Frayh et al. 1997), 23% (Al-Ghamdi et al. 2019), 17% (Al-Ghamdi et al. 2022).	Yemen children and adults 10% (Al-Mehdar et al. 2025)	55-143
<i>Blomia tropicalis</i>			
Coast	17% (Hassan et al. 2013).	Yemen children and adults 4% (Al-Mehdar et al. 2025)	5 ---
Highlands			3 ---

<sup>a</sup> Percentages of sensitizations refer in most but not all cases to patient cohorts of allergy clinics; these percentages are only indicative; they only make sense in the specific context of each study, and in some comparisons between mite species. The last column enumerates the number of mites of these species found in the various climate regions in this survey. \*(2018) Second or repeated sampling of *D. farinae* *D. pteronyssinus* mites only

**Table 3** House Dust Mite Species collected in Saudi Arabia, with related literature

	House dust Saudi Arabia	Apart from house dust Saudi Arabia	House dust Arabian Peninsula
<i>Asigmata Pyroglyphidae</i>			
<i>Dermatophagoides farinae</i>	C 62 <sup>a</sup> ; Dammam (Al-Qurashi 2006), Jeddah (Edrees 2008, 2009, 2010, 2012) D 16, H 123 C 52; Jeddah (Edrees 2008, 2009, 2010, 2012) D 12, H 198	Storage Jeddah	Kuwait (Gamal-Eddin and Hamad 1992) Kuwait (Gamal-Eddin et al. 1985; Gamal-Eddin and Hamad 1992) New species
<i>D. pteronyssinus</i>			
<i>Dermatophagoides n. sp.</i>	C 104, D 0, H 0 New species	New species	
<i>Acaridae</i>			
<i>Aleuroglyphus ovatus</i>	C 11; Jeddah (Edrees 2008, 2009, 2012) D 1, H 3		
<i>Acarus siro</i>	C 1; Jeddah (Edrees 2009; Edrees 2012) D 2, H 26		
<i>Sancassania berlesei</i>	C 2, D 0, H 0 First record		First record
<i>Tyrophagus putrescentiae</i>	C 15; Jeddah (Edrees 2009; Edrees 2012) D 0, H 28	Hail (Al-Shemmary 2016)	Kuwait (Gamal-Eddin et al. 1985; Gamal-Eddin and Hamad 1992)
<i>Carpoglyphidae</i>			
<i>Carpoglyphus lactis</i>	C Jeddah (Edrees 2012)	Jeddah	
<i>Echimyopodidae</i>			
<i>Blomia freemani</i>	C Jeddah (Edrees 2009, 2012)		
<i>B. tropicalis</i>	C 5; Jeddah (Edrees 2008, 2009, 2010, 2012) D 0, H 3		Kuwait (Gamal-Eddin and Hamad 1992)

Table 3 (continued)

	House dust, Saudi Arabia	Apart from house dust Saudi Arabia	House dust Arabian Peninsula
Glycyphagidae			
Glycyphagus domesticus	C 2, D 1, H 5 First record	Riyadh (Al-Khalifa and Bayoumi 1983)	First record
Histiostomatidae			
Histiostoma feroniarum	C 6, D 0, H 1 First record	Likely first record	First record
Histiostoma sp.	C 0, D 0, H 5 First record	Majmaah (Mashaly et al. 2011)	First record
Suidasiidae			
Suidasia pontifica	C 7, D 0, H 6 First record		First record
S. nesbitti	C Jeddah (Edrees 2008, 2009, 2012)		
Prostigmata			
Acarophenacidae			
Acarophenax tribolii		Storage Jeddah (Al-Nasser 2011)	
Caligonellidae		Storage Riyadh (Rostom 1993)	
Cheyletidae			
Acaropsella volgini		Storage Dammam, Riyadh (Rostom 1993)	
Acaropsis sollers		Storage Dammam Riyadh (Rostom 1993)	
Cheyletus eruditus	C 29, D 51, H 37 First record	Riyadh (Al-Youssif and Soliman 1979; El-Bahrawy and Al-Dakhil 1993)	First record
C. malaccensis	C Jeddah (Edrees 2008, 2009, 2012)	Storage Dammam (Rostom 1993), Jeddah; Hail (Al-Shermary 2016), Riyadh (Al-Youssif and Soliman 1979)	Kuwait (Gamal- Eddin and Hamad 1992) First record
Cheletopsis sp.	C 0, D 10, H 0 First record		
Erytraeidae			
Abrolophus spp.		Storage Dammam, Riyadh (Rostom 1993)	

**Table 3** (continued)

	House dust, Saudi Arabia	Apart from house dust Saudi Arabia	House dust Arabian Peninsula
Tetranychidae			
Bryobia sp.	C 1, D 0, H 6 First record	Tabuk, Riyadh, Qassim, Baha, Wadi e Hamifa (Alatawi and Kamran 2018)	First record
Eutetranychus sp.	C 4, D 0, H 12 First record	Throughout (Kamran et al. 2018)	First record
Tetranychus spp.	C 5, D 1, H 9 First record	Throughout (Alatawi and Kamran 2018)	Kuwait (Gamal- Eddin and Hamad 1992)
Mesostigmata			
Blattisociidae		Storage Dammam (Rostom 1993), Jeddah; Riyadh (Rostom 1993)	
Blattisocius keegani		Storage Jeddah	
B. tarsalis			
Laelapidae			
Stratiolaelaps miles	C 4, D 4, H 9 First record		Kuwait (Gamal- Eddin and Hamad 1992)
Macrochelidae			
Macrocheles muscaedomesticae	C 5, D 2, H 6 First record	Makkah Riyadh (Samsřinák 1979; Alatawi et al. 2018), Qassim (Fouly and Al-Rehiayani 2011)	First record
Macronyssidae			
Ornithonyssus bacoti	C Dammam (Al-Qurashi 2006)		
Melichariidae			
Proctolaelaps pygmaeus	C 5, D 3, H 11 First record	Makkah	First record
Oribatida			
Ceratozetidae			

Table 3 (continued)

	House dust Saudi Arabia	Apart from house dust Saudi Arabia	House dust Arabian Peninsula
Ceratozetes sp.	C 0, D 0, H 6 First record		First record
Oribatulidae			
Oribatula tibialis	C 16, D 2, H 28 First record	(Bayoumi and Al-Khalifa 1986)	First record
Schelorbaitidae			
Schelorbaites sp.	C 3, D 0, H 10 First record	(Bayoumi and Al-Khalifa 1986)	First record

<sup>a</sup>The table lists the mite species found in this study and any other species found in house dust in Saudi Arabia and Arabian Peninsula. C #, D #, H #: total number of mites collected from Coast, Desert, Highland samples in this study

## Processing of samples for their arthropod content

Dust samples were collected from human dwellings by their owners or inhabitants with the aid of their own vacuum cleaner devices, extracted from floors, carpets and rugs at a rate of 1 m<sup>2</sup> per minute and for 5 min. Once the vacuuming process was completed, the dust collected from the 5 households (of each city) were transferred into jars containing 96% ethanol, followed by alcohol evaporation afterwards, all done by the Saudi Arabian 1st author in its home institution (see authors affiliations). The samples were poured from the jars into open sealing bags contained in trays (so widely open) allowing slow evaporation for up to 3 days (to avoid shrinking and loss). The preserved/fixed arthropods into labelled sealed plastic bags were transported to the laboratory in UK for further analysis.

Once in the laboratory at the University of Reading, UK, the preserved dry specimens mixed with some dust particles were rehydrated in 70% alcohol (v/v, water/alcohol). For this, the bags were re-filled with 70% alcohol and allowed rehydration of specimens over night to avoid shrinking and potential loss. Then they were divided in subsamples to ease the examination under the stereomicroscope (Motic SMZ-171). All micro-arthropods and/or arthropod parts were separated for further identification. The extracted mite specimens were then permanently mounted on glass slides in Hoyer's medium for microscopic examination (Krantz 1978). The material was deposited in the Forensic Acarology Collection at the University of Reading.

A Nikon Optiphot phase contrast microscope was utilised for identification of the specimens. Images were captured using a Motic Moticam 3 Plus (microscope digital camera). Accordingly, the following keys and descriptions were used for mite families and species identification (Fain 1967; Van Bronswijk et al. 1973; Van Bronswijk and De Cock 1973; Hughes 1976; Wharton 1976; Perez 1993; Colloff 1998; Mašán 2003; Krantz and Walter 2009; Michaud et al. 2012; Solarz 2012; Alatawi et al. 2018; Kamran et al. 2018). For the separation of *Dermatophagoides* species, the key of Hart and Fain (1988) was used, considering the epimeral plates (ventral), hysteronotal shield and middle dorsum, bursa copulatrix (females) and size (thickness) of legs 1 and 3 (males).

## Data analyses

Data were counts of individual mites per species, family, by city and region. Boxplots were built for mite counts allowing visualization of families or species and totals (StataCorp 2014). Kruskal-Wallis rank tests were used to compare species of Pyroglyphidae by region and in more detail, by city. In order to check for dependency between the mite species in the Pyroglyphidae (*Dermatophagoides*) and their locations, Pearson pairwise correlations and the Kendall rank correlation coefficient (Kendall's  $\tau_b$  but no Effect size) were also estimated with an alpha of 0.05. These are non-parametric tests and do not depend on assumptions founded on the type of distribution.

A Generalized Linear Model (GLM) Poisson Regression based, Log likelihood, was applied to study the predator-prey interaction by region using (1) *Cheyletus eruditus* (HDM predator) as dependent variable, and *D. farinae*, *D. pteronyssinus*, *Dermatophagoides* n. sp. and *Blomia tropicalis* as independent variables; and (2) *Cheyletus eruditus* vs. Pyroglyphidae and Acaridae. In addition, a *t*-test for paired samples was conducted between Pyroglyphidae and *Cheyletus eruditus* to account for dependency of the data. Data from the

remaining families were not statistically analysed due to the low counts in each family and the great number of zeros (0). In addition, indicator species analysis (IndVal) was applied on the three regions; for which  $p$ -values of indicators were run by 9999 random permutations (with an alpha of 0.05). The  $p$ -values were Bonferroni corrected. Graphs and statistical analyses were performed with StataCorp (2014) and PAST4 (Hammer et al. 2001; Hammer and Harper 2024).

Articles published in journals considered to be predatory are not referenced.

## Results and Discussion

### Diversity of house-dust mites

Five hundred and seventy four specimens of allergy-causing *Dermatophagoides* and *Blomia* species (Pyroglyphidae, Echimyopodidae) have been collected (Tables S1 (sampling of 2018) and S2 (*Dermatophagoides* repeated sampling)). While mite allergens have been widely reported, no mites had been collected before in the desert and highlands.

A total of 998 mites were identified from 4 mite orders, Astigmata (considered a cohort within the Oribatida), Oribatida, Mesostigmata, and Prostigmata, 14 families, and 23 identified morpho-species (Tables S1 and S2).

A new species of *Dermatophagoides* (Pyroglyphidae) has been collected.

*Sancassania berlesei* (Acaridae), *Histiostoma feroniarum* (Histiostomatidae), *Suidasia pontifica* (Suidasiidae), *Cheletopsis* sp. (Cheyletidae) and *Ceratozetes* sp. (Ceratozetidae) are first records for Saudi Arabia and the Arabian Peninsula (Table 3).

*Glycyphagus domesticus* (Glycyphagidae), *Histiostoma* sp. not *H. feroniarum* (Histiostomatidae), *Cheyletus eruditus* (Cheyletidae), *Bryobia* sp. and *Eutetranychus* sp. (Tetranychidae), *Macrocheles muscaedomesticae* (Macrochelidae), *Proctolaelaps pygmaeus* (Melicharidae), *Scheloribates* sp. (Scheloribatidae), *Oribatula tibialis* (Oribatulidae), and *Scheloribates* sp. (Scheloribatidae) are first records for house dust in Saudi Arabia and the Arabian Peninsula (Table 3).

*Tetranychus* spp. (Tetranychidae) and *Stratiolaelaps miles* (Laelapididae) are new records for house dust in Saudi Arabia; they have been previously described for house dust in Kuwait (Table 3).

Among these, 5 species are new to the dust mite fauna worldwide: Prostigmata: *Cheletopsis* sp. (Cheyletidae); Mesostigmata: *Proctolaelaps pygmaeus* (Melicharidae), *Macrocheles muscaedomesticae* (Macrochelidae); and Oribatida: *Oribatula tibialis* (Oribatulidae), *Ceratozetes* sp. (Ceratozetidae) (Table 3).

A detailed discussion of the mites collected and the nature of the findings is presented after Table 3.

### Astigmata, Sarcoptiformes, Acariformes

excluding the results and analysis of *Dermatophagoides* spp, which are presented later.

*Aleuroglyphus ovatus* (Troupeau, 1879) (Acaridae, Acaroidea).

*A. ovatus* or the brown-legged grain mite (basionym *Tyroglyphus ovatus*).

There are some twenty reports of *A. ovatus* in house dust in Europe, Africa, and Asia (Somorin et al. 1978; Maurya et al. 1983; Suto et al. 1992; Yu et al. 2015). *A. ovatus* is present in house dust in Jeddah (Edrees 2008, 2009, 2010). Interestingly, the mite has not yet been collected from natural habitats in Saudi Arabia. The mite has also been retrieved from chicken houses in Israel (Mumcoughlu and Lutsky 1990). In workers on poultry farms, the mite can enter the ear canal and cause tinnitus (Kato et al. 2011). It is a well-known stored product mite of some 35 commodities dominated by grains and flour but also found associated with food items like bread, candied fruit, bean curd, cheese, dried fish and shrimps, pasta, and processed food (Saleh et al. 1985; Hagstrum et al. 2013a). Patients allergic to *Dermatophagoides* species show crossreactivity to *A. ovatus* and *Chortoglyphus arcuatus* (Troupeau, 1879) (Puerta et al. 1993).

*Acarus siro* Linnaeus, 1758 (Acaridae, Acaroidea).

The grain mite *A. siro* (synonyms *Acaris farris* (Oudemans, 1905), *Acarus farinae* Linnaeus, 1758, *Acarus farinae* De Geer, 1778, *Acarus farris* (Oudemans, 1905), *Aleurobius farinae* Canestrini, 1888, *Aleurobius farris* Oudemans, 1902, *Tyroglyphus farinae* (De Geer, 1778) has been detected in house dust in Jeddah (Edrees 2009, 2012). It is a well-known house dust mite with reports from many countries like Türkiye (Aykut 2021), India (Gill and Dhaliwal 2018; Saw et al. 2024), Egypt (Heikal 2015a). It has been reported from litter in chicken houses in Israel some 35 years ago (Mumcoughlu and Lutsky 1990), yet, it seems still to spread. The first confirmed sighting from Brazil, for example, is only from 2022 (Barbosa et al. 2022), however, sensitization has been detected earlier (Ferraroni et al. 2013). The species is known as a stored product mite, *siro* referring to a grain-storage pit, and Linnaeus finding it in flour, the species is now known from some 88 commodities including baby food, biscuits, cheese, bread, dried soup, and halva. The mite is deliberately added to the rind of the cheese mimolette to increase its flavour. In rare cases, it can cause oral mite anaphylaxis (Preiser-Funke and Bergmann 2023).

*Sancassania berlesei* (Michael, 1903) (Acaridae, Acaroidea).

This is a first record for the wet grain or mushroom musty mite, *S. berlesei* (basionym *Tyroglyphus berlesei*, synonyms *Caloglyphus berlesei* (Michael, 1903), *C. rodionovi* Zachvatkin, 1941, *Cologlyphus berleis lapsus duplex*), in house dust for Saudi Arabia and the Arabian Peninsula. It highly depends on wet environments, explaining why it was only found in Jeddah. It is a house dust mite recorded from many countries. It might contribute to atopy (Saxena et al. 1980). The species occurs in occupational ear acariasis of a worker in a waste processing plant in Spain (Poggioli and Taboada 2021) and caused tinnitus in a worker on a chicken farm in Japan. The mite is common in litter from chicken houses in Israel (Mumcoughlu and Lutsky 1990). It is a stored product mite especially of some twenty commodities under slightly damp or mouldy conditions (Hagstrum et al. 2013a; EL-Sayed 2021). Egypt features several sister species as stored product mites, *S. betae* (Attiah, 1969), *S. krameri* (Berlese, 1881), *S. mycophagus* (Mégnin, 1874), and *S. rhicoglyphoides* (Zachvatkin, 1937).

*Tyrophagus putrescentiae* (Schrank 1781) (Acaridae, Acaroidea).

The copra itch, ham or mould mite, *T. putrescentiae* (basionym *Acarus putrescentiae*), has been isolated from house dust in Jeddah (Edrees 2009, 2012). It feeds on moulds including entomopathogenic fungi (Ou et al. 2024). It is a formidable stored product mite affecting some 142 different commodities (Hagstrum et al. 2013a; EL-Sayed 2021; Klimov et al. 2024). It is also a threat to the poultry industry (Sulzbach et al. 2024). In Egypt, *T. zachvat-*

*kini* Volgin, 1948 serves as a stored product mite as well. There is cross-reactivity between the allergens of *Dermatophagoides farinae*, *D. pteronyssinus*, and *T. putrescentiae* (Chen et al. 2025).

*Carpoglyphus lactis* (Linnaeus, 1767) (Carpoglyphidae, Hemisarcoptidea).

*C. lactis*, basonym *Acarus lactis*, has been collected from house dust in Jeddah (Edrees 2012). It is known as a house dust mite in Europe, North and South America, and Asia.

It is a storage mite too. Linnaeus described it as living in milk, and one of the German vernacular names is milk mite (Milchmilbe). However, it is better known as the dried fruit mite for its prevalence on sugar-rich substrates; some 40 commodities including wine corks carry this mite (Stejskal et al. 2015). Many beehives in Europe and the USA bear *C. lactis*, where they consume stored pollen; the mite also shows up in honey. In the production of sherry, some oak casks are colonised by such amounts of *C. lactis* that they lead to a higher floral and refreshing sensation, but also to a slightly increased bitterness in the final wine (Marín et al. 2009).

*C. lactis* and the acaroid species of concern here, *Aleuroglyphus ovatus*, *Acarus siro*, *Dermatophagoides farinae*, *D. pteronyssinus*, *Glycyphagus domesticus*, *Sancassania berlessei*, *Suidasia nesbitti*, and *Tyrophagus putrescentiae* can sometimes occur in such numbers in food products that they might cause urinary and intestinal acariasis; they have been recovered from human urine and feces (Li et al. 2003).

Some phytoseeid predatory species used in biocontrol are commercial reared on *C. lactis* as prey species (Cao et al. 2025).

The sister species, *C. ganzhouensis* Jiang, 1991 is found in house dust and on brown sugar in slaughter houses in China, while *C. munroi* (Hughes, 1952) in barn dust in Sweden, in bee hives in Czechia, and on spider webs in the UK.

*Blomia freemani* Hughes, 1948 (Echimyopodidae, Glycyphagoidea).

Edrees (2009); (Edrees 2012) found *B. freemani* in dust of houses, motels, and hotels in Jeddah. In Alexandria, Egypt, *B. freemani* is one of the dominant species in house dust (Rezk et al. 1996). Although originally described from the Northern Ireland, reports from house dust in Europe (Switzerland, Wales) and South America (Colombia) are rare, while it is quite common in house dust in China and India. In China, *B. freemani* together with *Dermatophagoides farinae* and *D. pteronyssinus* were more abundant in mattress dust than in house dust (Lai 1988). The breeding rate of acaroid mites like *B. freemani* was the highest in taxis, followed by private cars, but less so in buses (Wang and Li 2012).

Like several other mite species contaminating stored flour, *B. freemani* has caused oral mite anaphylaxis (pancake syndrome) (Wen et al. 2005). *B. freeman* has originally been described as a mite of stored wheat, it is a storage mite of rice, millet, sweet potato chips, and Chinese medicine.

In addition to *B. tropicalis*, sister species *B. kulagine* Zakhvatkin 1942, *B. thori* Zakhvatkin 1942, and *B. tjibodas* (Oudemans, 1910) are also associated with house dust.

*Blomia tropicalis* Van Bronswijk, De Cock and Oshima 1973 (Echimyopodidae, Glycyphagoidea).

*B. tropicalis* has been repeatedly described from house dust in Jeddah (Edrees 2008, 2009, 2010, 2012). It sometimes occurs together with *B. freemani* (Edrees 2009, 2012). It has a rich medical history (Caraballo et al. 2024). Severe cases of allergic rhinitis can result from dual sensitization to *D. pteronyssinus* and *B. tropicalis* (González-Pérez et al. 2020). Patients with allergic rhinitis sensitized to *B. tropicalis* might be most vulnerable to air

pollutants (Quek et al. 2019). There are several vaccine candidates based on *B. tropicalis* (Castro-Almarales et al. 2020; da Silva et al. 2020). It is also a stored product mite of rice and tobacco.

*Glycyphagus domesticus* (de Geer, 1778) (Glycyphagidae, Glycyphagoidea).

This is a first record for *Glycyphagus* and *G. domesticus* (basonym *Acarus domesticus*) in house dust for Saudi Arabia and the Arabian Peninsula. As the species epithet suggests, *G. domesticus* is known as the house mite. Already in 1983, *G. domesticus* has been recovered from soil in the Riyadh region (Al-Khalifa and Bayoumi 1983). *G. domesticus* in house dust, for example, in Egypt has been published twice (Heikal 2015a, b), in Türkiye (Mutlu et al. 2019), France (Reboux et al. 2019), or Poland (Solarz et al. 2021). It can build massive colonies on furniture stuffed with not thoroughly cleaned horse hair (Oudemans 1897). It feeds on the fat still adherend to the hair. Together with *Acarus siro* and *Lepidoglyphus destructor*, *G. domesticus* can cause oral mite anaphylaxis (Mulhall and Conlon 2019). It is a stored product mite of some 44 commodities. In addition, in Egypt, *G. aegyptiacus* Attiah, 1971, and *G. oryzae* Attiah, 1971 are known as stored product mites.

*Histiostoma feroniarum* (Dufour, 1839) (Histiostomatidae, Histiostomatoidea).

This is a first record for *Histiostoma* and *H. feroniarum* (basonym *Hypopus feroniarum*), synonyms *Tyroglyphus rostroserratus*, *Acarus mamillaris*, in house dust for Saudi Arabia and the Arabian Peninsula. There might be three specimens of *H. feroniarum* collected in Saudi Arabia in the Royal Belgian Institute of Natural Sciences, Brussels, Belgium according to the Global Biodiversity Information Facility. They are not obvious in the museums catalogue. The only species published from Saudi Arabia is *H. sammari* Eraky & Shoker, 1994 phoretic on ant species (Mashaly et al. 2011). *H. feroniarum* is known from house dust in several countries, for example, China (Li et al. 2005), island of Tenerife, Spain (Sanchez-Covisa et al. 1999), Venezuela (Hurtado and Parini 1987) or Australia (Colloff et al. 1991). *H. humidatus* (Vitzthum, 1927) and *H. sapromyzae* (Dufour, 1839) are fellow house dust mites (Lee and Cho 1984; Colloff et al. 1991). *H. feroniarum* is a stored product mite of bread, wet grain, dried fruit, dried vegetables, and Chinese medicine. *H. feroniarum* is common in mushroom houses (Szafranek and Lewandowski 2017; Qu et al. 2023). *H. onioni* Eraky & Shoker, 1994 and *H. sarrai* are stored product mites of onions in Egypt.

*Suidasia pontifica* Oudemans, 1905 (Suidasiidae, Acaroidea).

This is a first record for *S. pontifica*, the scaly grain mite, in house dust for Saudi Arabia and the Arabian Peninsula. There is some confusion about the naming of the species. In 1978, *S. medanensis* Oudemans 1924 was synonymized with *S. pontifica* Oudemans 1906, retaining the original and therefore the correct name, *S. pontifica* (Fain and Philips 1978). This was never opposed; yet, *S. medanensis* is now more often used in the understanding that they are the same species. *S. melanensis* in the literature is a spelling error. *S. pontifica* is widespread as a house dust mite (Somorin et al. 1978; Colloff et al. 1991; Mariana et al. 2000; Modak et al. 2004). Some commercial skin prick test solutions in India for *Dermatophagoides pteronyssinus* or *Blomia tropicalis* contained instead proteins of *S. pontifica* (Huber et al. 2021). The importance of *S. pontifica* as a stored product mite is even greater. Together with its sister species, *S. nesbittii* Hughes, 1948, it is found in some 55 commodities (Hagstrum et al. 2013a; Bakr et al. 2022; Ta-Phaisach et al. 2023). It is associated with a variety of food products. The large occurrence in a pancake mix in Panama lead to the pancake syndrome in the form of oral mite anaphylaxis (Barrera et al. 2015; Sánchez-Borges et al. 2020). The pancake syndrome can also be induced by other mite species like

*Dermatophagoides farinae* (Erben et al. 1993; Kano and Murata 2025), *D. pteronyssinus*, *Tyrophagus putrescentiae* (Takahashi et al. 2014), *Acarus siro* (Preiser-Funke and Bergmann 2023), and the booklouse *Liposcelis bostrychophila* Badonnel, 1931 (Matsumoto et al. 2025). Oral mite anaphylaxis might also look like an acute asthma attack (Garcia et al. 2016). *S. pontifica* was found together with *Tyrophagus putrescentiae*, *Blattisocius tarsalis* (Berlese, 1918), and *Blomia tropicalis* in peanut bars and milk sweets in Brazil (Franzolin et al. 1994).

*Suidasia nesbitti* Hughes, 1948 (Suidasiidae, Acaroidea).

*S. nesbitti*, the scaly grain or wheat pollard itch mite, has been found in house dust of Jeddah several times (Edrees 2008, 2009, 2012). It is reported from Europe, the Middle East, and Asia as a dust mite. It is also a cosmopolitan pest of grains and feed high in protein and fat (Devi et al. 2022). *S. nesbitti* has been collected from as a soil mite from soil of a moss-turf area and the wet soil from a kitchen liquid waste collection area of the Schirmacher Oasis in East Antarctica (Sanyal 2004).

## Prostigmata, Trombidiformes, Acariformes

*Cheyletus eruditus* (Schrank 1781) (Cheyletidae, Cheyletoidea).

This is a first record for *C. eruditus* (basonym *Acarus eruditus*, syn. *C. butleri* Hughes, 1948, *C. desitus* Qayyum & Chaudhri, 1977, *C. ferox* Trouessart, 1889, *Cheyletus ferox* Banks, 1906, *C. seminivorus* Packard, 1878, and others) in house dust for Saudi Arabia and the Arabian Peninsula. It is also known under vernacular names like grain itch mite, mattress itch mite, prairie itch mite, and so on, however, Schrank named it a Büchermilbe, book mite, when he described it from the bookbinder's glue of humid books, making it a true domestic mite (Schrank 1781). This predator mite has been associated with maize grains and rodents in Saudi Arabia (Al-Youssif and Soliman 1979; El-Bahrawy and Al-Dakhil 1993; Alatawi and Kamran Ahmad 2017). *C. eruditus* or its sister species, *C. malaccensis* (Oudemans, 1903) are almost always found in house dust samples (Liu et al. 2010; Saw et al. 2018; Zhou et al. 2022; Karuthedath and Raghavan 2024). *C. malaccensis* has been found in house dust in Jeddah city (Edrees 2008). The two species are the main predators of the *Dermatophagoides* species in house dust. Although *Dermatophagoides* species have been extensively associated with atopic diseases, *C. eruditus* can also be of clinical relevance for persistent, non-occupational allergic rhinitis and skin symptoms (Poza Guedes et al. 2016; Hsu et al. 2024; Castromil-Benito et al. 2025). As a stored product mite, it is also associated with more than 53 commodities ranging from grains, spices, seeds, dried fruits to Chinese medicine (Hagstrum et al. 2013a). The species is commercially used as a biocontrol agent against a number of grain mites, in particular *Acarus siro* and *Lepidoglyphus destructor* (Schrank 1781) in agricultural warehouses.

*Cheyletus malaccensis* (Oudemans, 1903) (Cheyletidae, Cheyletoidea).

*C. malaccensis*, basonym *Cheletes malaccensis*, synonyms *Cheletes fortis* Oudemans, 1904, *C. vorax* Oudemans, 1903; *Cheyletus avidus* Qayyum & Chaudhri, 1977, *C. ayyazi* Akbar, Aheer & Ishtiaq, 1993, *C. baridos* Akbar, Rahi & Chaudhri, 1988, *C. caucasicus* Zachvatkin, 1949, *C. egypticus* Elbadry, 1969, and some ten more.

What applies to *C. eruditus* also applies to *C. malaccensis*. As a predator species it is found in house dust as well as in stored products. Although it does not carry a commonly used vernacular name, it has been encountered in more countries with a truly worldwide

distribution, and it affects even a wider diversity of commodities than *C. ereditus*. As a biocontrol agent, it is newer. *C. malaccensis* prefers a slightly warmer climate; *malaccensis* refers to Malacca (Melaka), Malaysia. It predate better on booklice (Hu et al. 2025) and it is considered for biocontrol in poultry farms. The expression of heterotrophic males is much more pronounced in *C. malaccensis*. Both species engage in cannibalism if prey is scarce. The colour of *C. malaccensis* changes with the prey it consumes, yellowish with *Aleuroglyphus ovatus* and *Carpoglyphus lactis*, blackish with *Lasioderma serricorne* Fabricius, 1792, and brown spots with *Cryptolestes pusillus* (Schönherr, 1817) (Han et al. 2025). When *C. malaccensis* bites, meaning stabbing human skin, it tries to feed and causes papules (Yoshikawa 1987).

*Cheletopsis* sp. Oudemans, 1904 (Cheyletidae, Cheyletoidea).

This is a first record for *Cheletopsis* in house dust worldwide. It is also a first record for this genus for Saudi Arabia, the Arabian Peninsula, and Asia. *Cheletopsis* species are mostly known from feather quills and bird's nests (Bochkov et al. 2002; Bochkov and Skoracki 2012). They predate on other mite species.

*Tetranychus* spp. Dufour, 1832 (Tetranychidae, Tetranychoidae).

This is a first record for *Tetranychus* in house dust for Saudi Arabia. The red spider mites, *T. evansi* Baker & Pritchard, 1960, *T. neocaledonicus* André, 1933, *T. salicornicus* Alatawi and Kamran 2018; *T. turkestanii* (Ugarov & Nikolskii, 1937), and *T. urticae* Koch, 1836 (Alatawi and Kamran 2018) are known from Saudi Arabia; *T. salicornicus* is a new species described from Riyadh. Tetranychidae have been reported early on as part of the house dust fauna (Solomon 1961), and they are often mentioned in the house dust literature but rarely identified to the genus or species level. *Tetranychus* species occur in houses worldwide, for example, in Australia, Colombia, Iceland, Türkiye (Colloff et al. 1991; Hallas et al. 2004; Navarro et al. 2008; Zeytun et al. 2015, 2016) and Kuwait (Gamal-Eddin and Hamad 1992).

*Eutetranychus* sp. Banks, 1917 (Tetranychidae, Tetranychoidae).

This is a first record for the genus *Eutetranychus* in house dust for Saudi Arabia and the Arabian Peninsula. *E. africanus* (Tucker, 1926), *E. banksi* (McGregor, 1914), the citrus brown mite, *E. orientalis* (Klein, 1936), and *E. palmatus* Attiah, 1967 are known from Saudi Arabia (Kamran et al. 2018). *E. orientalis* has been found in house dust in India (Modak et al. 2004).

*Bryobia* sp. C.L.Koch, 1835 (Tetranychidae, Tetranychoidae).

This is a first record for *Bryobia* in house dust for Saudi Arabia and the Arabian Peninsula. Four species are known from Saudi Arabia, *B. alveolata* Auger & Flechtmann, 2009 originating from Tunisia, the clover mite, *B. praetiosa* Koch, 1836, *B. rubrioculus* (Scheuten, 1857), and *B. tuttlei* Smiley & Baker, 1995 originating from Yemen (Alatawi and Kamran 2018). There are some 149 named species in this genus (Mirza et al. 2025). *B. praetiosa* is known from house dust in Italy and Israel (Castagnoli et al. 1983; Mumcuoglu and Shalom 2013). While numbers of *Bryobia* mites in house dust are normally in the single digits, mainly in the past in Europe and North America, often new houses had been invaded in the spring by hundreds of thousands, of grass mites, *B. praetiosa* or *B. graminum* (Schrank 1781) (syn. *B. cristata* (Dugès, 1834), leading, for example, to a quarter of a million on the floor of a bedroom in the USA (English and Snetsinger 1957; Anderson and Morgan 1958; Rack 1984; Mumcuoglu and Shalom 2013). In England, new police headquarters and Magistrates' courts were overrun by such high numbers of mites that they had to be closed (Parkin 1958). The buildup of large populations in just one or two years is facilitated by

parthenogenetic reproduction in many species. *B. graminum* has been described from house dust of rural farms in Iceland (Gígjá 1944; Guðmundsson et al. 2008). Unidentified species of *Bryobia* have been collected from house dust in Czechia, in the living and bed rooms of houses in Türkiye, mattresses in Denmark, and in a house in Israel (Costa 1978; Vobrůzková et al. 1979; Andersen 1984; Zeytun et al. 2016). These mites often develop in sunny places where grass reaches directly the wall of houses and buildings, which was often the case in new urban or suburban developments of the 1950's to the 1980's. The mites are very mobile and enter the houses mostly through open windows by accident in search for food, mainly grass, which they will not find indoors; eventually starving to death notwithstanding any house plants. Large amounts of eggs might be deposited on outside window frames. The mites climb up the walls of buildings up to the fourth floor. When crushed, they leave green stains behind. They do not attack humans or animals.

### Mesostigmata, Parasitiformes

*Proctolaelaps pygmaeus* (Müller, 1859) (Melicharidae, Ascoidea).

This is a first record for *P. pygmaeus* (basonym *Gamasus pygmaeus*, and some 20 synonyms including *Lasioseius ventritrichosus* Schweizer, 1949, *Garmania alpina* (Schweizer, 1949) in house dust worldwide. It has been found in wheat floor in Jeddah and in manure of chicken farms in the Makkah region. At other places, it is associated with mouldy barley, bulbs, and timber, hay, rice and other grains. The mite feeds on fungi and predaes on small arthropods. It stands out for being able to feed on solid food. It can bite humans and cause extensive papular dermatosis; this has been reported from timber workers in Australia and New Zealand (Halliday et al. 1998). Damp construction wood possibly introduced it into a newly constructed office building in Berlin, Germany.

*Macrocheles muscaedomesticae* (Scopoli, 1772) (Macrochelidae, Eviphidoidea).

Surprisingly, this is a first record for *M. muscaedomesticae* (basonym *Acarus muscae domesticae*), synonym *Macrocheles lundae* Krantz, 1970, for house dust worldwide. Wherever there is the house fly *Musca domestica* Linnaeus, 1758, vinegar flies like *Drosophila* Fallen, 1823 species, stable flies such as *Stomoxys calcitrans* (Linnaeus, 1758) or scuttle flies like *Megaselia* Rondani, 1856 species, one might expect to find *M. muscaedomesticae* as well (Perotti 2001; Perotti and Braig 2009). The manure of cattle sheep and poultry farms in the Makkah region all contained this species. It has been associated with dried fish, grain, maize trash, onion bulbs, mouldy peanuts, and rotting wheat residues elsewhere. *M. muscaedomesticae* have also been associated with human corpses (Braig and Perotti 2009; Che Kamaruzaman et al. 2018).

*Ornithonyssus bacoti* (Hirst, 1913) (Macronyssidae, Dermansossoidea).

*O. bacoti*, basonym *Leioznathus bacoti*, has a long list of synonyms including *Bdellonyssus nagayoi* (Yamada, 1930), *Hirstionyssus musculi* Domrow, 1963, *Lipionyssus lutzi* Fonseca, 1942, *Liponyssus meprai* Manso Soto & Pletneff, 1951, *Macronyssus bacoti* (Hirst, 1913). It is known as the tropical rat mite. The adjective tropical is slightly misleading as it now can be found worldwide. It is a blood-feeding mite that causes a non-specific dermatitis in humans and is often un- or misdiagnosed. Bites might occur in a cluster. The mite leaves its host after bloodfeeding. Many pathogens and parasites have been transmitted in laboratory experiments, but its role as a vector to humans is still doubted with the exception of the bacterium *Bartonella henselae* causing cat-scratch disease (Bradley et al.

2014). Unfed protonymphs have survived up to 43 days (Sudd 1952), Kelaher et al. (2005) claim survival of the mite for up to 63 days without a bloodmeal. *O. bacoti* is able move over longer distances, which means it can be found in houses and workplaces that are not infested by rodents (Theis et al. 1981). This implies that an infestation with this mite is not necessarily associated with run-down homes or workplaces but can occur also on modern, well-maintend premises (Kelaher et al. 2005).

The mite shows little host specificity, even humans can act as an alternative host. It can associate with a wide diversity of pet, companion, and domesticated animals, examples are African pygmy hedgehog in Germany (van Well and Beck 2022), donkey in France (Dumitrache et al. 2023), hamsters, gerbils, guinea pigs, rabbits, degus, kangaroo mice, hedgehogs, tricolour squirrels, and sugar gliders in Italy (d'Ovidio et al. 2018), dogs in the USA (Bradley et al. 2014), in Japan (Ogata et al. 1978), cats (Baker 1999), and chicken (Ponnudurai et al. 2015; Villanueva et al. 2020). The owners of the animals might become infested as well.

On this description, one would not expect finding this mite in house dust or in large numbers. However, in the port city of Dammam in Saudi Arabia, from the dust of houses of children suffering allergenic manifestations, *D. farinae* and *O. bacoti* were recovered (Al-Qurashi 2006). Although *O. bacoti* has not been reported from house dust for any other country of the Arabian Peninsula, the mite has been found on rats in Qatar (Islam et al. 2023) and should be assumed to be present over the entire Arabian Peninsula. Outside Saudi Arabia, dermatitis patients in Egypt had *D. pteronyssinus*, *O. bacoti* and *Haemogamasus pontiger* (Berlese, 1904) in their house dust (Morsy et al. 1994). In rural houses in Egypt, *O. bacoti* was the most numerous mite in house dust, more abundant even than pyroglyphid species (El-Shazly et al. 2006). In Japan, the density of *O. bacoti* is between 16 and 530 mites per gram of house dust, with little difference between houses built from wood or concrete (Toma et al. 1993b). Infestations have been reported from older student accommodation in Edinburgh, Scotland (Sargison et al. 2025). In ten kindergardens in Iran, 20% ( $N=342$ ) of the mites collected from dust were *O. bacoti*, only 18% *D. farinae* (Soleimani-Ahmadi et al. 2017).

*Stratiolaelaps miles* (Berlese, 1892) (Laelapididae, Dermanyssoidea).

This is a first record for *S. miles* (basionym *Laelaps miles*, syn. *Hypoaspis miles* (Berlese, 1892) for Saudi Arabia. It has been reported from house dust in Kuwait and India (Gamal-Eddin and Hamad 1992; Modak et al. 2004). It is a stored product mite of decaying oatmeal spillage and wheat residue. The mite predate on flies, thrips, springtails, and other mites. Owners of pet reptiles, amphibians and invertebrates like spiders use this species to prevent or treat infestations by other mites. In the snail breeding industry, it is used against the parasitic while snail mite *Riccardoella limacum* (Schrank, 1776) (Ereynetidae). Mushroom farms use it against the *Lycoriella* Frey, 1942 sciarid fly and fungus gnats. The widespread use as a biocontrol agent has led to IgE sensitization against this species in greenhouse workers (Kronqvist et al., 2005). There has been some confusion in the biocontrol industry about the identity of *S. miles* and *S. scimitus* (Womersley, 1956) (Walter and Campbell 2003).

## **Oribatida (Sarcoptiformes, Acariformes)**

*Oribatula tibialis* (Nicolet, 1855) (Oribatulidae, Oripodoidea).

This is a first record for *O. tibialis* (basonym *Notaspis tibialis*, and some 16 synonyms) in house dust worldwide. *O. tibialis* has been recorded from sandy soil under ficus trees in Samitah, Saudi Arabia (Bayoumi and Al-Khalifa 1986). There is also one report from Oman (Vasar et al. 2022). The high abundance soil mites (Oribatida) of the family Oribatulidae seems unusual for indoors. However, Oribatida highest prevalence in the highland region matches with their natural habitat, areas with tree cover, more vegetation that can sustain Oribatida communities (Galal et al. 2021). In fact, the Oribatida is a poorly known group of mites from the Arabian Peninsula. Aside from being a soil mite it is also known as a ‘stored timber mite’. It can be found in wood chips (Klimek and Chachaj 2018). It has been implicated in dermatitis found on the trunk and arms of wood workers (Liguori et al. 1989). The species stands out as one of a few animals releasing hydrogen cyanate as a chemical defence against predators (Brückner et al. 2017). Lichens and bird nests are additional habitats (Melekhina 2023; Melekhina et al. 2023). Because of its small size, a subspecies is living amidst the adhesive traps on leaves of a carnivorous plant (Antor and Garcia 1995). *O. tibialis* is becoming common in European cities (Mangová et al. 2019). What we see here in Saudi Arabia is the transition of this species to a house dust mite. *O. tibialis* is a stored product mite of dried apricots in Türkiye (Çobanoğlu 2008). One *Oribatula* species has been reported from house dust in Korea and China, *O. sakamorii*, now *Phauloppia sakamorii* (Aoki, 1970) (Ree et al. 1997; Chen et al. 2023). Unidentified species of Oribatulidae have been seen in house dust of Japan and Colombia (Oshima 1970; Charlet et al. 1977; Suto et al. 1991; Hatsushika and Miyoshi 1992; Toma et al. 1993a).

*Scheloribates* Berlese, 1908 sp. (Scheloribatidae, Oripodoidea)

This is a first record for *Scheloribates* in house dust for Saudi Arabia and the Arabian Peninsula. *Andeszetes* Hammer, 1961, *Megascheloribates* Lee & Pajak, 1990, *Neoscheloribates* Hammer, 1973, *Paraschelobates* Jacot, 1935, *Protoschelobates* Jacot, 1934, *Semischeloribates* Hammer, 1973, and *Storkania* Jacot, 1929 are synonyms of the genus *Scheloribates*. Four to six species of *Scheloribates* (Bayoumi and Al-Khalifa 1986; Kardar 1988) are known from soil and leaf litter in various places of Saudi Arabia: *S. fimbriatus* subsp. *africanus* Wallwork, 1964 (Al-Khalil, Khamsin), *S. laevigatus* (Koch, 1835) (Riyadh, Durma, Jalajil, Al-Zulfi, Al-Qatif, Al-Sayl Al-Kabir, Samitah), *S. pallidulus* (Koch, 1841) (Al-‘Uwaynah, Al-Surrah), *S. riyadhiensis* Kardar 1988 (Riyadh), now considered a *species inquirenda* by Subías (2022), *S. saudicus* Bayoumi & Al-Khalifa, 1985 (I-Kharj, I-Dir‘iyah) and *S. saudiensis* Kardar 1988 (Riyadh), now also considered a *species inquirenda* by Subías (2022). *S. sacculipunctatus* Mahunka 2009 is a new species from leaf litter near Al-Hayer in the UAE (Kardar 1988; Mahunka 2009). Species of *Scheloribates* have been found in the dust of several houses in Brazil (Rosa and Flechtmann 1979) and Colombia (Charlet et al. 1977), *S. laevigatus* has been found in house dust in China (Chen et al. 2023), *S. latipes* (Koch, 1844) now a subspecies of *S. pallidulus*, the same species that is the most common *Scheloribates* in Saudi Arabia, has been found in house dust in Korea (Ree et al. 1997) and *rufafulvus*, now *Protoribates* (*P.*) *rufafulvus* (Kardar, 1977) and considered a *species inquirenda* by Subías (2022); and *S. translamellaris* (Kadar, 1977), now *S. praeincisus* subsp. *sandvicensis* (Jacot, 1934), in house dust in India (Maurya and Jamil 1982). *S. laevigatus* is also a pest of stored barley, rice, and wheat grains and mouldy grains, potatoes and floor debris (dust) in Iran, Japan, USA, and Europe (Ostovan 1993; Hagstrum et al. 2013a).


*Ceratozetes* Berlese, 1908 sp. (Ceratozetidae, Ceratozetoidea)

This is a first record for *Ceratozetes* in house dust worldwide. It is also a first record for this genus and family for Saudi Arabia and the Arabian Peninsula. The mites inhabit soil, litter, and canopy habitats. Many are adapted to colder and montane areas. There are currently some 48 species in the genus *Ceratozetes* (Behan-Pelletier and Eamer 2009; Subías 2022). Species geographically closest to the Arabian Peninsula are *C. baleensis* Ermilov et al. 2011 and *gracilis* (Michael, 1884) in forest soil of mountains in Ethiopia (Ermilov et al. 2011) and *conjunctus* Mihelcic, 1956 and *C. macromediocris* Shaldybina, 1970 in Iran (Akrami 2015). *C. conjunctus* has also been found in soil of industrial sites of the Basque Country (Corral-Hernández and Iturrondobeitia 2012). *Ceratozetes* has not been associated with any stored products.

### Analysis of mite families and association with main predators

Our study on all families of house dust mites found in Saudi Arabia, which corresponds to the 2nd sampling attempt (2018), shows a diverse mite composition and variation between regions (Fig. 2; Table S1). From a general view, the most abundant mites belong to the families Pyroglyphidae ( $N_{total} = 400$ ), followed by mites from the families Cheyletidae ( $N_{total} = 127$ ) and Acaridae ( $N_{total} = 115$ ). Oribatid soil mites ( $N_{total} = 65$ ) were unexpectedly numerous in the house-dust (families Oribatulidae, Ceratozetidae and Scheloribatidae). The remaining families did not reach a 50 counts threshold for their inclusion in the analysis.

Considering the total of HDM found, their abundance significantly differs in numbers between the three main regions ( $X^2: 16.302, df: 2, P < 0.001$ ). In particular while Pyro-



**Fig. 2** Boxplots of counts of mites within families by region. Boxplot description: Median, central line; box, lower and upper quartiles; external lines, lower and upper extremes; dots, outliers.

glyphidae had borderline variation in abundance (significance) between the three climatic regions ( $X^2$ : 5.692, df: 2,  $P=0.0551$ ), their main predator species, *C. eruditus* showed consistent differences comparing regions ( $X^2$ : 17.579, df: 2,  $P<0.001$ ). When compensating for data dependency, by pairing Pyroglyphidae with *C. eruditus*, the two populations varied significantly in each region (by equal means,  $P<0.001$  for all regions), and *C. eruditus* had a significant effect on populations of *D. farinae* in the desert region (GLM,  $P<0.01$ ). Using families Pyroglyphidae and Acaridae as independent variables, there was interaction between *C. eruditus* and Pyroglyphidae (GLM,  $P<0.01$ ) in the desert, while no interaction with Acaridae was observed in any region.

## The Indicator Species Analysis (IndVal)


The bird/mammal skin cheyletid, *Cheletopsis* sp. resulted as a significant marker for the desert. *D. pteronyssinus*, *Histiostoma* sp. and *Ceratozetes* sp. were found significantly associated with the highlands, while *Dermatophagoides* n. sp. resulted in the highest, significant association with the coastal region, followed by *Aleuroglyphus* sp. and *Histiostoma feroniarum*.

## Detailed analysis of *Dermatophagoides* species

From the data of the repeated sampling,  $2 \times 25$  samples a total of 566 specimens within 3 species were collected from the 3 regions: *Dermatophagoides pteronyssinus* ( $N_{\text{total}} = 262$ ), *D. farinae* ( $N_{\text{total}} = 200$ ) and *Dermatophagoides* n. sp. ( $N_{\text{total}} = 104$ ) (Fig. 3, Table S2). For none of the three species the data follow a linear distribution, even by pooling numbers of mites in regions (Shapiro-Wilk W test,  $P<0.0001$ , in all 3 cases). Between regions, pairwise correlations were different between *D. farinae* and *Dermatophagoides* n. sp. in the coast ( $P=0.05$ , borderline), and *D. farinae* and *D. pteronyssinus* in the highlands area ( $P<0.001$ ).

Kendall correlation coefficient suggests a positive correlation (concordant ranking) between *D. farinae* and *D. pteronyssinus* ( $P<0.01$ ) regardless of region, while *Dermatophagoides* n. sp. shows low or no correlation (is discordant) with the other two species, suggesting an independent distribution pattern. Within regions, there was a highly positive correlation between *D. farinae* and *D. pteronyssinus* in the highlands ( $P<0.01$ ). In other words, there is an association between these species. They occur together in the dust samples collected from different Saudi Arabian households.

Our studies of species interaction showed that within Saudi Arabian houses the common predator of Pyroglyphidae, *C. eruditus*, follows the same regional distribution pattern of Pyroglyphidae, specifically targeting populations of *D. farinae* in the desert. Interestingly, Saudi populations of *C. eruditus* seem not to significantly affect Acaridae species; but this could be due to the smaller samples of Acaridae in Saudi houses, compared to other regions of the world (Smiley 1991; Bertone et al. 2016). Predatory cheyletids are common in house dust, preying on other micro-arthropods (such as pyroglyphid mites), moving indoor via phoresy and/or entering associated with pets, as well as on domestic plants or on the humans themselves (Colloff 1998, 2009). Cheyletidae were abundantly found in the house-dust from the desert area, thus may be associated with a dry hot climate that might only sustain Pyroglyphidae due to the humidity produced by air conditioning (Van Strien et al. 2004;




**Fig. 3** Boxplots of counts of *Dermatophagoides* mites by species, *D. pteronyssinus*, *D. farinae* and *Dermatophagoides* sp. n. (new species), by region and sampling (A; 1st sampling and B: 2nd or repeated sampling). Boxplot description: median, central line; box, lower and upper quartiles; external lines, lower and upper extremes; dots, outliers

Thomas 2010). Acaridae, on the other hand, are quite prevalent in temperate regions of the world. Acaridae species have lower temperature thresholds.

Pyroglyphidae, Acaridae, and Oribatulidae are positively correlated. The three groups can utilise same food resources such as fungi and detritus, given the right environmental conditions, especially humidity (Arlan 1992; Arlian and Dippold 1996; Arlian and Morgan 2003; Colloff 2009; Klien and Walzl 2010). For Pyroglyphidae, this is perhaps due to the frequent humidity bursts combined with the right temperatures in these climatic regions (Hughes 1976; Zhang 1998; Colloff 2009; Thomas 2010); they favour indoor environments with high levels of humidity (Klien and Walzl 2010). This correlation can also be linked to the confirmed (negative) interaction with their main predator, *C. eruditus*.

The other unidentified Cheyletidae species found in this study, *Cheletopsis* sp., a bird mite, was only recorded from desert samples. Its occurrence in the desert helps explain that desert habitation is perhaps more exposed to the visit of wild birds and/or mammals (Hughes 1976; Bochkov 2002; Walter et al. 2009).

Of particular interest is the occurrence of a third, new species of *Dermatophagoides*, restricted only to the coastal region (*Dermatophagoides* n. sp.). The differences of the three *Dermatophagoides* species are shown in Fig. 4. This new species, which is currently being described, plus the prevalent *Aleuroglyphus* and *Histioglyphus feroniarum* could become reliable biomarkers for location, as traces of potential utility in forensic investigations (Hani et al. 2018).



**Fig. 4** *Dermatophagoides* mite species from Saudi Arabia, ventral view of males of (a) *D. pteronyssinus*, (b) *D. farinae* and (c) *Dermatophagoides* n. sp. . Bar, 100 µm, arrows showing epimeral plates

## Conclusions

Dust mites in Saudi Arabia are much more divers than expected with first records, species for the first time associated with house dust, and mite species likely new to science. This work shows that some dust mite species are specific for climate regions. It differentiates the importance of HDMs for allergies in coastal, desert, and highland areas; and it reviews all known HDMs for Saudi Arabia.

**Acknowledgements** We want to thank the house owners of the Kingdom of Saudi Arabia, who donated the vacuumed samples of house-dust and send them to us. We are thankful to Dr Marta Saloña-Bordas for helping confirming mite genera of the Oribatida.

**Author contributions** RHSA and MAP designed this study. RHSA collected the mites. RHSA, MH, and MAP identified the mites. RHSA, MAP and HRB analysed the data. RHSA, MAP, and HRB wrote the manuscript. All authors read and approved the final manuscript.

**Funding** This study was funded by UKRI EPSRC EP/Y036743/1 (to MAP) and the Saudi Arabian Cultural Bureau in London, SACB Reference Number I1124: University of Reading REF F3345756 (to RHSA and MAP).

**Competing Interests.**

The authors have no relevant financial or non-financial interests to disclose.

**Data availability** All data generated or analysed during this study are included in this published article.

## Declarations

**Competing interests** The authors declare no competing interests.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Abal AT, Ayed A, Nair PCMG, Mosawi M, Behbehani N (2010) Factors responsible for asthma and rhinitis among Kuwaiti schoolchildren. *Med Princ Pract* 19:295–298
- Abdel-Rahman EH, Abdelgadir M, AlRashidi M (2020) Ectoparasites burden of house mouse (*Mus musculus* Linnaeus, 1758) from Hai'1 of Saudi Arabia. *Saudi J Biol Sci* 27:2238–2244
- Aburuz S, Bulatova N, Tawalbeh M (2011) Skin prick test reactivity to aeroallergens in Jordanian allergic rhinitis patients. *East Mediterr Health J* 17:604–610
- Acuna-Cantillo L, Maldonado-Santana H, Calderón-Llosa Ó (2024) Revision de registros historicos de los acaros del polvo domestico en paises del Pacifico y Caribe continental en Latinoamerica [Review of the historical records of house dust mites in Pacific and Latin America Continental countries]. *Rev Alerg Mex* 71:70
- Adham TM, Tawfik SA (2012) *Dermatophagoides* in childhood asthma. Allergy to *Dermatophagoides* associates more severe childhood asthma with a potential role for acaricides. *Saudi Med J* 33:292–297
- Adham TM, Tawfik SA, Abdo NM (2011) House dust mites in pediatric atopic dermatitis. *Saudi Med J* 32:177–182
- Akdemir C, Karaman Ü, Cebeci Güler N, Direkel S, Uzunoğlu E, Şenturk H, Ayhan U (2023) Ordu, Giresun, Trabzon ve Rize İllerinde Ev Tozu Akarlarının Yaygınlığı ve Der p 1 ve Der f 1 Varlığı [Prevalence of house dust mites and presence of Der p 1 and Der f 1 in Ordu, Giresun, Trabzon and Rize provinces]. *Türk Parazitol Derg* 47:179–183
- Akrami MA (2015) An annotated checklist of oribatid mites (Acari: Oribatida) of Iran. *Zootaxa* 3963:451–501
- Al Khater SA (2017) Sensitization to common aeroallergens in asthmatic children in the Eastern region of Saudi Arabia. *Saudi J Med Med Sci* 5:136–141
- al Mousawi M, Behbehani N, Arifhodzic N, Lovel H, Woodcock A, Custovic A (2001) Environmental allergens in Kuwait. *Allergy* 56:1237–1238
- al Mousawi M, Ramadan MAG, Taher MM, Behbehani N (2009) Allergic sensitization in healthy school children in Kuwait: an emerging public health concern. *Kuwait Med J* 41:317–321
- Al Sharif H, Hassan E, Albarazi R (2017) Common aeroallergens and food allergens seen among allergic children in the United Arab Emirates. *Allergy* 72:398–399
- Al-Aani S, Alhammadi S (2015) Uncontrolled asthma in children. *Allergy* 70:422–422
- Al-Abri, R., D. Bharghava, M. Kurien, V. Chaly, Y. Al-Badaai, K. Bharghava (2015) Allergic rhinitis and associated comorbidities: Prevalence in Oman with knowledge gaps in literature. *Oman Med J* 29, 414–418
- Al-Dowaisan A, Fakim N, Khan MR, Arifhodzic N, Panicker R, Hanoon A, Khan I (2004) *Salsola* pollen as a predominant cause of respiratory allergies in Kuwait. *Ann Allergy Asthma Immunol* 92:262–267
- Al-Dusari SN, Somorin AO, Zaman AU (1997) IgE antibodies to mites in Saudi Arabia. *Allergy* 52:234–235
- Al-Frayh A, Gad-El-Rab MO, Alnajjar A, Hasnain SM (1992) A comparative study of immediate skin-test reactivity to inhalant allergens in asthmatic-children of 2 different regions in Saudi Arabia. *Ann Saudi Med* 12:468–471
- Al-Frayh AS, Hasnain SM, Gad-El-Rab MO, Schwartz B, AlMobairek K, AlSedairy ST (1997) House dust mite allergens in Saudi Arabia: regional variations and immune response. *Ann Saudi Med* 17:156–160
- Al-Ghamdi BR, Koshak EA, Omer FM, Awadalla NJ, Mahfouz AA, Ageely HM (2019) Immunological factors associated with adult asthma in the Aseer region, southwestern Saudi Arabia. *Int J Environ Res Public Health* 16:2495
- Al-Ghamdi BR, Omer FM, Awadalla NJ, Mahfouz AA (2022) Specific aeroallergen sensitization associated with current rhinitis among adults in southwestern Saudi Arabia. *Medicine (Baltimore)* 101:e30384
- Al-Khalifa MS, Bayoumi BM (1983) New records of soil Acarina and Collembola in Riyadh region. *J Coll Sci King Saud Univ* 14:57–61
- Al-Mehdar AA, Almorish MAW, Aldomini M, Alshoabi SA, Elkhalifa AME, Elhadi FEM, Bazie EA, Alhasan MM, Al-Sayaghi KM, Albadrani MS (2025) Aeroallergen sensitization patterns among allergic respiratory diseases patients living in Sana'a City, Yemen. *Aerobiologia* 2025, s10453-025-09878-6
- Al-Nahdi M, Al-Quorain AA (1987) Sex distribution and common allergens of bronchial asthma in a Saudi Arabian (Eastern Province) population. *Allergol Immunopathol (Madr)* 15:389–391
- Al-Nasser AS (2011) Quantitative survey of stored products mites infesting wheat flour in Jeddah Governorate. *predatory* 3:78–84
- Al-Qurashi AM (2006) House dust mites and allergic manifestations among some children in Dammam, Saudi Arabia. *J Egypt Soc Parasitol* 36:283–288
- Al-Shalan AA, Al-Frayh A, Reilly H, Al-Hussein KA, Wilson JD (1989) Inhalant allergens in patients with allergic rhinitis in Riyadh. *Ann Saudi Med* 9:331–336

- Al-Shemmary KA (2016) Thermal requirements and life table parameters of the predatory mite *Cheyletus malaccensis* Oudemans (Acari: Cheyletidae) in Hail region, Saudi Arabia. *Egypt J Biol Pest Control* 26:741–746
- Al-Tamemi S, Al-Shidhani A, Al-Abri R, Jothi B, Al-Rawas O, Al-Riyami B (2008) The pattern of sensitization to inhaled allergens in Omani patients with asthma, allergic rhinitis and rhinoconjunctivitis. *Allergy* 63:222–222
- Al-Youssif MS, Soliman ZR (1979) Cheyletid mites of Saudi Arabia with a key to species (Acari: Acariformes: Prostigmata). *Bull Entomol Soc Egypt* 62:213–218
- Al-Zayadneh EM, Alnawaiseh NA, Altarawneh AH, Aldmour IH, Albataineh EM, Al-Shagahin H, Alharazneh A, Alzayadneh E (2019) Sensitization to inhaled allergens in asthmatic children in southern Jordan: a cross-sectional study. *Multidiscip Respir Med* 14:37
- Alatawi FJ, Kamran M (2018) Spider mites (Acari: Tetranychidae) of Saudi Arabia: two new species, new records and a key to all known species. *J Nat Hist* 52:429–455
- Alatawi FJ, Kamran Ahmad S (2017) Predatory prostigmatid mite (Acari: Trombidiformes) fauna of the date palm agro-ecosystem in Saudi Arabia. *Syst Appl Acarol* 22:1444–1475
- Alatawi FJ, Kamran M, Mirza JH (2018) Mesostigmatic mites (Acari: Mesostigmata) of Saudi Arabia (excluding Phytoseioidea), new records and a key to the known species. *Zootaxa* 4388:373–394
- Alhazmi L, Oraibi O, Areeshi A, Alhazmi N, Hakami A, Kariri T, Akish E, Alamer A, Sayed A, Madkhali MA, Somaili M, Altherwi T, Alhazmi A, Madkhali MA, Hakami A (2025) Awareness of environmental risk factors contributing to asthma exacerbations among asthmatic patients in the Jazan region, Saudi Arabia. *Medicine (Baltimore)* 104:e41598
- Aljohani A, Clarke D, Byrne M, Gormally M, Fleming G (2024) Diversity and abundance of house dust mites in Irish homes, their exoskeleton-associated bacteria and susceptibility to clinically relevant antibiotics. *Biol Environ Proc R Ir Acad* 124B:29–41
- Almehizia AA, AlEsa RK, Alwusaidi KM, Alzamil KA, AlJumah M, Aljohani S, Almutairi AF, Salam M (2019) Allergic rhinitis: disease characteristics and coping measures in Saudi Arabia. *PLoS One* 14:e0217182
- Almogren A (2009) Airway allergy and skin reactivity to aeroallergens in Riyadh. *Saudi Med J* 30:392–396
- Alqahtani JM (2016) Asthma and other allergic diseases among Saudi schoolchildren in Najran: the need for a comprehensive intervention program. *Ann Saudi Med* 36:379–385
- Alqahtani JM (2020) Atopy and allergic diseases among Saudi young adults: a cross-sectional study. *J Int Med Res* 48:300060519899760
- Alrasheedi SM, Alkhalifah KM, Alnasyan S, Alwattban RR, Alsubhi RA, Alsamani RI, Alfouzan YA (2023a) Asthma exacerbations in asthmatic adult patients in the Qassim region of Saudi Arabia: a cross-sectional study. *Cureus* 15:e44997
- Alrasheedi SM, Alrashdi MN, Alkhdairi A, Alrashidi HM, Alrasheedi A, Alharbi EB, Almutairi FA, Alrashdi KS, Allabun FS, Ijtaitly GA (2023b) Awareness of asthmatic patients to the association between the environmental risk factors and asthma exacerbation in Al-Qassim region, Saudi Arabia. *Med Sci* 27:10.54905/diss/v27i133/e155ms2711
- Andersen A (1984) Abundance and spatial distribution of house-dust mites in their natural environment (Acari: Sarcoptiformes & Trombidiformes). *Entomol Medd* 52:25–32
- Anderson NH, Morgan CVG (1958) Life histories and habits of the clover mite, *Bryobia praetiosa* Koch and the brown mite, *B. arborea* M. and A., in British Columbia (Acarina: Tetranychidae). *Can Entomol* 90:23–42
- Antor RJ, Garcia MB (1995) A new mite plant association – mites living amidst the adhesive traps of a carnivorous plant. *Oecologia* 101:51–54
- Arlian LG (1992) Water balance and humidity requirements of house dust mites. *Exp Appl Acarol* 16:15–35
- Arlian LG, Dippold JS (1996) Development and fecundity of *Dermatophagoides farinae* (Acari: Pyroglyphidae). *J Med Entomol* 33:257–60. <https://doi.org/10.1093/jmedent/33.2.257>
- Arlian LG, Morgan MS (2003) Biology, ecology, and prevalence of dust mites. *Immunol Allergy Clin North Am* 23:443–468
- Asad F, Fernandez AR, Kumar A, Shoaib MA (2015) Co-existence of asthma with allergic rhinitis and common allergens in Al Ain, UAE: a cross-sectional study. *Ann Abbasi Shaheed Hosp Karachi Med Dent Coll* 20:137–141
- Asseri AA, Abuqail MA, Alotaibi AS, Abuqail WA, Alqahtani AS, Asiri LA, Alkhayri M, Moshebah AY, Elabd FM (2025) Sensitization patterns to aeroallergens and food allergens among children with atopic asthma in southwestern Saudi Arabia. *Children* 12:573
- Aykut M (2021) Comparison of goose down and woolen pillows in terms of house dust mites. *Syst Appl Acarol* 26:2070–2077
- Aykut M (2024) An annotated checklist of house dust mites (Acari: Acariformes) of Türkiye. *Persian J Acarol* 13:797–811

- Badran HS, Hussein A, Salah M, Lotfi WT (2016) Identification and prevalence of allergic, nonallergic, and local allergic rhinitis patients in Western Area, Saudi Arabia. *Ann Otol Rhinol Laryngol* 125:634–643
- Baker AS (1999) Mites and Ticks of Domestic Animals, an identification guide and information source. Natural History Museum, London
- Bakr AA, Saad MMG, Abdelgaleil SAM (2022) Mite incidence in Egyptian storage facilities and acaricidal activity of selected monoterpenes, phenylpropenes, and sesquiterpenes against *Suidasia medanensis* Oudemans (Astigmata: Suidasiidae), a formidable storage mite pest. *Persian J Acarol* 11:101–113
- Barbosa MFC, Corrêa AS, Moraes GJ (2022) *Acarus siro* L. (Astigmatina: Acaridae): first confirmed record for Brazil based on morphological and molecular characterization. *J Stored Prod Res* 97:101958
- Barrera OM, Murgas IL, Bermudez S, Miranda RJ (2015) Anafilaxia oral por ingestión de alimentos contaminados con acaros en Ciudad de Panamá, 2011–2014 [Oral anaphylaxis by ingestion of mite contaminated food in Panama City, 2011–2014]. *Rev Alerg Mex* 62:112–117
- Bayoumi BM, Al-Khalifa MS (1986) Oribatid Mites (Acari) of Saudi Arabia. In: Büttiker W, Krupp F (eds) *Fauna Saudi Arab*, vol 7. Karger, p 66–92, Basel, pp 66–92
- Behan-Pelletier VM, Eamer B (2009) *Ceratozetes* and *Ceratozetoides* (Acari: Oribatida: Ceratozetidae) of North America. *Can Entomol* 141:246–308
- Bener A, Safa W, Abdulhalik S, Lestringant GG (2002) An analysis of skin prick test reactions in asthmatics in a hot climate and desert environment. *Allerg Immunol (Paris)* 34:281–286
- Bertone MA, Leong M, Bayless KM, Malow TLF, Dunn RR, Trautwein MD (2016) Arthropods of the great indoors: characterizing diversity inside urban and suburban homes. *PeerJ* 4:e1582
- Bochkov AV (2002) The classification and phylogeny of the mite superfamily Cheyletoidea (Acari, Prostigmata). *Entomol Rev* 82:643–664
- Bochkov AV, Skoracki M (2012) New and poorly known predaceous cheyletids (Acariformes: Cheyletidae) from feather quills. *Int J Acarol* 38:486–496
- Bochkov AV, Fain A, Dabert J (2002) A revision of the genus *Cheyletopsis* (Acari Cheyletidae). *Bull de l'Institut Royal des Sci Nat de Belgique Entomologie* 72:5–26
- Bradley JM, Mascarelli PE, Trull CL, Maggi RG, Breitschwerdt EB (2014) *Bartonella henselae* infections in an owner and two papillon dogs exposed to tropical rat mites (*Ornithonyssus bacoti*). *Vector-Borne Zoonotic Dis* 14:703–709
- Braig HR, Perotti MA (2009) Carcasses and mites. *Exp Appl Acarol* 49:45–84
- Brückner A, Raspošnič G, Wehner K, Meusinger R, Norton RA, Heethoff M (2017) Storage and release of hydrogen cyanide in a chelicerate (*Oribatula tibialis*). *Proc Natl Acad Sci U S A* 114:3469–3472
- Buczek W, Bartosik K, Buczek A (2023) Development of *Dermacentor reticulatus* ticks in human household conditions. *J Pest Sci*. <https://doi.org/10.1007/s10340-023-01695-5>
- Çakan H, Güven K, Çevik FE, Demirci M, Saribas S (2015) Investigation of human DNA profiles in house dust mites: implications in forensic acarology. *Rom J Legal Med* 23:187–192
- Cao JL, Zhang KS, Li XT, Zhang ZQ (2025) Individual development and population growth of four phytoseiid predators feeding on *Carpoglyphus lactis* (Acari: Phytoseiidae, Carpoqlyphidae). *Syst Appl Acarol* 30:1534–1538
- Caraballo L, Lockey R, Puerta L, Zakzuk J, Acevedo N, Fernandez-Caldas E (2024) *Blomia tropicalis*: A 50-year history. *J Allergy Clin Immunol Pract* 2024, 11.007
- Castagnoli M, Liguori M (1983) N. Nannelli Gli acari della polvere delle case in Italia [House dust mites in Italy]. In: *Atti XIII del Congresso Nazionale Italiano di Entomologia, Sestriere, Torino*, p 577–582
- Castro-Almarales RL, Ronquillo-Díaz M, Alvarez-Castelló M, Rodríguez-Canosa J, González-León M, Enriquez-Domínguez I, Navarro-Viltre BI, Mateo-Morejón M, Oliva-Díaz Y, Ramirez-González W, Cox L, Labrada-Rosado A (2020) Subcutaneous allergen immunotherapy for asthma: a randomized, double-blind, placebo -controlled study with a standardized *Blomia* vaccine. *World Allergy Organ J* 13:100098
- Castromil-Benito ES, Betancor D, Parron-Ballesteros J, Gutierrez-Diaz G, Gonzalez-Perez R, Poza-Guedes P, Bernalda J, Mederos-Luis E, Rial Prado MJ, Iraola V, Moran Morales M, Cuesta-Herranz J, Villalba M, Pastor-Vargas C (2025) Predatory mite *Cheyletus eruditus* as a novel allergenic source: Identification of its allergens and cross-reactivity. *J Allergy Clin Immunol Pract* 2025, 04.034
- Charlet LD, Mulla MS, Sanchez-Medina M (1977) Domestic Acari of Colombia: Acaridae, Glycyphagidae and Cryptostigmata recovered from house dust. *Int J Acarol* 3:55–64
- Che Kamaruzaman NA, Mašan P, Velásquez Y, González-Medina A, Lindström A, Braig HR, Perotti MA (2018) *Macrocheles* species (Acari: Macrochelidae) associated with human corpses in Europe. *Exp Appl Acarol*. <https://doi.org/10.1007/s10493-018-0321-4>
- Chen H, Zhu Y, Fan H, Zhao W, Liu L, Cao Y, Ye C, Sun E (2023) Comparative study on DNA barcoding of house oribatid mites based on the 18S and 28S ribosomal DNA gene. *Chin J Parasitol Parasit Dis* 41:163–169

- Chen J, Zhou Y, Ren Y, Wang K, Zhong L, Zhou D, Chen L, Cheng Q, Liao Y, Cui Y, Hao C (2025) Characterization and cross-reactivity assessment of group-29 allergens originating from *Dermatophagoides pteronyssinus*, *Dermatophagoides farinae*, and *Tyrophagus putrescentiae*. *Asian Pac J Allergy Immunol* 18:173–180
- Clarke D, Burke D, Gormally M, Byrne M (2015) Dynamics of house dust mite transfer in modern clothing fabrics. *Ann Allergy Asthma Immunol* 114:335–340
- Çobanoğlu S (2008) Mites (Acari) associated with stored apricots in Malatya, Elazığ and İzmir provinces of Turkey. *Turk Entomol Derg* 32:3–20
- Colloff MJ (1998) Taxonomy and identification of dust mites. *Allergy* 53:7–12
- Colloff MJ (2009) *Dust Mites*, Springer, Dordrecht
- Colloff MJ, Stewart GA, Thompson PJ (1991) House dust acarofauna and Der p I equivalent in Australia: the relative importance of *Dermatophagoides pteronyssinus* and *Euroglyphus maynei*. *Clin Exp Allergy* 21:225–230
- Corpuz-Raros LA (2022) An updated list of domestic mites (Acari) from the Philippines. *Philipp Entomol* 36:1–14
- Corral-Hernández E, Iturrondobeitia JC (2012) Effects of cattle and industries on oribatid mite communities of grassland soil in the Basque Country (Spain). *Int J Acarol* 38:217–229
- Costa M (1978) *Insects anti man [Insects against man]*. Hakibbutz Hameuchad Publ. House, Bnei Brak
- Crowther D, Horwood J, Baker N, Thomson D, Pretlove S, Ridley I, Oreszczyn T (2000) *House Dust Mites and the Built Environment: A Literature Review*. University College London, London
- Cui YB (2014) When mites attack: domestic mites are not just allergens. *Parasit Vectors* 7:411
- d'Ovidio D, Noviello E, Santoro D (2018) Prevalence and zoonotic risk of tropical rat mite (*Ornithonyssus bacoti*) in exotic companion mammals in southern Italy. *Vet Dermatol* 29:522–525
- da Silva ES, Aglas L, Pinheiro CS, de Andra Belitardo EMM, Silveira EF, Huber S, Torres RT, Wallner M, Briza P, Lackner P, Laimer J, Pacheco LGC, Cruz AA, Alcântara-Neves NM, Ferreira F (2020) A hybrid of two major *Blomia tropicalis* allergens as an allergy vaccine candidate. *Clin Exp Allergy* 50:835–847
- Devi P, Gulati R, Rani S (2022) *Uttar Pradesh J Zool*, 84–92
- Dowaisan A, Al-Ali S, Khan M, Hijazi Z, Thomson MS, Ezeamuzie CI (2000) Sensitization to aeroallergens among patients with allergic rhinitis in a desert environment. *Ann Allergy Asthma Immunol* 84:33–438
- Dumitrache MO, Györke A, Julien F, Kondratjeva J, Cadiergues MC (2023) Case report: identification of the tropical rat mite (*Ornithonyssus bacoti*) on a domestic donkey in France. *Front Vet Sci* 10:1141290
- Edrees NO (2008) Abundance, distribution, and species diversity of house dust mites in homes of patients with asthma in middle region of Jeddah City. *Egypt J Exp Biol (Zool)* 4:279–283
- Edrees NO (2009) Distribution dynamics of dust mites in two locations of patient homes with respect to the allergical kind. *Am-Eurasian J Agric Environ Sci* 6:680–688
- Edrees NO (2010) Influence of life styles and locations factors on mite in the dust of air conditioner filter, in three different regions in Jeddah Governorate. *Bio-Sci Res Bull* 26:131–142
- Edrees NO (2012) Prevalence of house dust mites in two levels of dorms (hotel and motel) of Jeddah District Western Saudi Arabia. *Life Sci J Acta Zengzhou Univ Overseas Ed* 9:3673–3683
- El-Bahrawy AAF, Al-Dakhil MA (1993) Studies on the interrelation between rodents and their ectoparasitic acarines in Riyadh Region, Saudi Arabia. *J Egypt Soc Parasitol* 23:675–685
- EL-Sayed MM (2021) The suitability of some storage products to *Tyrophagus putrescentiae* and *Sancassania berlesii* infestation accompanied with application some flavoring powders to reduce population. *Egypt Acad J Biol Sci B Zool* 13:339–351
- El-Shazly AM, El-Beshbishi SN, Azab MS, El-Nahas HA, Soliman ME, Fouad MAH, El-Salahy M, M.M. M (2006) Present situation of house dust mites in Dakahlia Governorate, Egypt. *J Egypt Soc Parasitol* 36:113–126
- Emara MM, Mansour HA, Shehata MT, Abu-Zahab Z (2012) Outcome of sublingual immunotherapy with multiple allergens in asthmatic patients with and without allergic rhinitis. *Life Sci J Acta Zengzhou Univ Overseas Ed* 9:817–829
- English LL, Snetsinger R (1957) The habits and control of the clover mite in dwellings. *J Econ Entomol* 50:135–141
- Erben AM, Rodriguez JL, McCullough J, Ownby DR (1993) Anaphylaxis after ingestion of beignets contaminated with *Dermatophagoides farinae*. *J Allergy Clin Immunol* 92:846–849
- Ermilov SG, Sidorchuk EA, Rybalov LB (2011) New oribatid mites of the superfamily Ceratozetoidea (Acari: Oribatida) from Ethiopia. *Afr Invertebr* 52:323–343
- Ezeamuzie CI, al-Mousawi M, Dashti H, Al-Bashir A, Al-Hage M, Al-Ali S (1997) Prevalence of allergic sensitization to inhalant allergens among blood donors in Kuwait - a desert country. *Allergy* 52:1194–1200
- Ezeamuzie CI, Al-Ali A, Khan M, Hijazi Z, Dowaisan A, Thomson MS, Georgi J (2000a) IgE-mediated sensitization to mould allergens among patients with allergic respiratory diseases in a desert environment. *Int Arch Allergy Appl Immunol* 121:300–307

- Ezeamuzie CI, Thomson MS, Al-Ali S, Dowaisan A, Khan M, Hijazi Z (2000b) Asthma in the desert: Spectrum of the sensitizing aeroallergens. *Allergy* 55:157–162
- Faccini JLH, Coutinho V (1987) Ocorrência de *Lynxacarus radovskyi* (Acari: Linstrophoridae) em gato doméstico no Brasil [Occurrence of *Lynxacarus radovskyi* (Acari: Linstrophoridae) in domestic cats in Brazil]. *Arq Univ Fed Rural Rio J* 9:91–93
- Fain A (1967) Le genre *Dermatophagoides* Bogdanov, 1864. Son importance dans les allergies respiratoires et cutanées chez l'homme (Psoroptidae: Sarcoptiformes) [The genus *Dermatophagoides* Bogdanov, 1864. Its importance in respiratory and skin allergies in humans (Psoroptidae: Sarcoptiformes)]. *Acarologia* 9:179–225
- Fain A, Philips JR (1978) Notes on the genus *Suidasia* Oudemans, 1905 with descriptions of a new species from Australia (Acari, Astigmata, Saprogllyphidae). *Int J Acarol* 4:115–123
- Fernández-Caldas E, Puerta L, Caraballo L (2014) Mites and allergies. In: Bergmann K-C, Ring J (eds) *History of Allergy*. Karger, Basel, pp 234–242
- Ferraroni NR, Oliveira S, Ferraroni JJ (2013) Prevalence of sensitization to the storage mite *Acarus siro* in middle-east of Brazil. *J Allergy Clin Immunol* 131 Suppl., AB163
- Fouly AH, Al-Rehiyani SM (2011) Predaceous mites in Al-Qassim Region, Saudi Arabia, with description of two new laelapid species (Acari: Gamasida: Laelapidae). *J Entomol* 8:139–151
- Franzolin MR, Baggio D, Correia M, Rodrigues RMMS (1994) Presence of mites in bars of peanut sweet and milk sweet, sold by street merchants in the city of Sao Paulo. *Rev Inst Adolfo Lutz* 54:11–15
- Frost CL, Braig HR, Amendt J, Perotti MA (2010) Indoor arthropods of forensic importance: Insects associated with indoor decomposition and mites as indoor markers. In: Amendt J, Campobasso CP, Goff ML, Grassberger M (eds) *Current Concepts in Forensic Entomology*. (Springer, Dordrecht, pp 93–108
- Gad-El-Rab MO (1999) The frequency of IgE antibodies specific to inhalant and food allergens in adult asthmatic patients in Riyadh, Saudi Arabia. *Ann Saudi Med* 19:447–449
- Galal TM, Al-Yasi HM, Fadl MA (2021) Vegetation zonation along the desert-wetland ecosystem of Taif Highland, Saudi Arabia. *Saudi J Biol Sci* 28:3374–3383. <https://doi.org/10.1016/j.sjbs.2021.02.086>
- Gamal-Eddin FM, Hamad MGM (1992) Present situation of house dust mites and its relation to currently recorded allergy in Kuwait state (the Gulf area). *J Egypt Soc Parasitol* 22(Suppl 3):1–101
- Gamal-Eddin FM, Aboul-Atta AM, Hamad MGM, Eraki A-SM, Turbeileh FMS (1985) House dust mites in Kuwait: Preliminary pilot survey to shed some light on diagnosis and treatment of mite allergy. *J Egypt Soc Parasitol* 15:313–322
- Garcia ME, Sanchez-Borges M, Capriles-Hulett A, Fernandez-Caldas E (2016) Oral mite anaphylaxis mimicking acute asthma. *Allergol Immunopathol (Madr)* 44:484–485
- Gígja G (1944) Meindyr í husum og grodri og varmir gegn heim [Pests in houses and plants and protection against them]. *Jens Gudbjörnsson, Reykjavik*
- Gill NK, Dhaliwal AK (2018) Seasonal variation of allergenic acarofauna from the homes of allergic rhinitis and asthmatic patients. *J Med Entomol* 55:262–268
- González-Pérez R, Pineda F, Poza-Guedes P, Castillo M, Matheu V, Sanchez-Machin I (2020) Molecular allergen profiling of dual mite sensitization in severe allergic rhinitis. *J Investig Allergol Clin Immunol* 30:421–429
- Guðmundsson G, Sigurðarson SP, Tómasson K, Gíslason D, Hallas TE (2008) Maurar í húsrýki á íslenskum bóndabæjum [Mites in house dust on Icelandic farms]. *Laeknabladid* 94:723–727
- Hagstrum DW, Klejdysz T, Subramanyam B, Nawrot J (2013a) *Atlas of Stored-Product Insects and Mites*. AACC International, St. Paul, Minnesota
- Hagstrum DW, Klejdysz T, Subramanyam B, Nawrot J (2013b) *Stored-product mites*. In: Hagstrum DW, Klejdysz T, Subramanyam B, Nawrot J (eds) *Atlas of Stored-Product Insects and Mites*. AACC International, St. Paul, Minnesota, pp 533–575
- Hallas TE, Gíslason D, Björnsdóttir US, Jörundsdóttir KB, Janson C, Luczynska CM, Gíslason T (2004) Sensitization to house dust mites in Reykjavik, Iceland, in the absence of domestic exposure to mites. *Allergy* 59:515–519
- Halliday RB, Walter DE, Lindquist EE (1998) Revision of the Australian Ascidae (Acarina: Mesostigmata). *Invertebr Taxon* 12:1–54
- Hammer Ø, Harper DAT (2024) *Paleontological data analysis*, 2nd edn. Wiley, Hoboken
- Hammer Ø, Harper DAT, Ryan PD (2001) *PAST: paleontological statistics software package for education and data analysis*. *Paleontol Electron* 4:1–9
- Han YB, Xue DR, Xia LY, Gao YF, Lü J, Su X, Liu ZQ, Wu Y (2025) Different species of stored-grain pest prey alter the demographic parameters of the predator *Cheyletus malaccensis* (Acari: Cheyletidae). *J Stored Prod Res* 112:102617
- Hani M, Thieven U, Perotti MA (2018) Soil bulb mites as trace evidence for the location of buried money. *Forensic Sci Int* 292:e25–e30. <https://doi.org/10.1016/j.forsciint.2018.09.016>

- Hart BJ, Fain A (1988) Morphological and biological studies of medically important house-dust mites. *Acarologia* 29:285–295
- Hasan HY, Rizwan MA (2015) Prevalence of inhalant and food allergies after skin prick testing in patients with resistant allergic rhinitis in warm humid weather. *Bahrain Med Bull*. <https://doi.org/10.12816/0014420>
- Hasnain SM, Al-Frayh AR, Subiza JL, Fernandez-Caldas E, Casanovas M, Geith T, Gad-El-Rab MO, Koshak E, Al-Mehdar H, Al-Sowaidi S, Al-Matar H, Khouqeer R, Al-Abbad K, Al-Yamani M, Alaqi E, Musa OA, Al-Sedairy S (2012) Sensitization to indigenous pollen and molds and other outdoor and indoor allergens in allergic patients from Saudi Arabia, United Arab Emirates, and Sudan. *World Allergy Organ J* 5:59–65
- Hassan MSM, Tayeb M, Amir EM, Wali AM, Mohamed FS (2013) Skin reactivity to allergens in Rabigh Area, Kingdom of Saudi Arabia. *J Egypt Soc Parasitol* 43:295–302
- Hatsushika R, Miyoshi K (1992) Studies on house-dust mites: the itching dermatitis in Okayama Prefecture, Japan. *Kawasaki Med J* 18:1–9
- Heikal HM (2015a) House dust mites: studies on the occurrence, identification and control in rural houses of Shebin El-Kom locality, Egypt. *Afr Entomol* 23:451–457
- Heikal HM (2015b) Studies on the occurrence, identification and control of house dust mites at rural houses of Shebin El-Kom locality, Egypt. *Pak J Biol Sci* 18:179–184
- Hijazi Z, Ezeamuzie CI, Khan M, Dowaisan AR (2002) Characteristics of asthmatic children in Kuwait. *J Allergy* 39:603–609
- Hsu C-H, Chien D-K, Tsai J-J, Yen C-Y, Liao E-C (2024) The prevalence of IgE positivity, environmental factors, and clinical implications associated with mite species in allergic patients in Taiwan. *Asian Pac J Allergy Immunol*. <https://doi.org/10.12932/ap-270623-1644>
- Hu Y, Xue D, Zhang P, Liu Z, Fang Z, Zhang Y, Wu Y (2025) Optimizing *Cheyletus malaccensis* releases for the biocontrol of *Liposcelis bostrychophila*: density-dependent efficacy in stored grains. *Exp Appl Acarol* 95:1–12
- Huber S, Gadermaier G, Bohle B, Ferreira F, Briza P (2021) Proteomic profiling of commercial dust mite skin prick test solutions and allergy vaccines from India. *World Allergy Organ J* 14:100516
- Hubert J, Stejskal V, Athanassiou CG, Throne JE (2018) Health hazards associated with arthropod infestation of stored products. *Annu Rev Entomol* 63:553–573
- Hubert J, Vrtala S, Sopko B, Dowd SE, He Q, Klimov PB, Harant K, Talacko P, Erban T (2023) Predicting *Blomia tropicalis* allergens using a multiomics approach. *Clin Transl Allergy* 13:clt2.12302
- Hughes AM (1976) *The Mites of Stored Food and Houses*, 2nd edition. Her Majesty's Stationary Office, London
- Hurtado I, Parini M (1987) House dust mites in Caracas, Venezuela. *Ann Allergy Asthma Immunol* 59:128–130
- Ibrahim RY, Eldeek HEM, Rezk HA, Othman RA, El-Tayeb AA, Elnazer MMA (2022) Species identification and seasonal prevalence of house dust mites in Assiut City, Egypt: a descriptive study in an urban area. *Persian J Acarol* 11:83–99
- Islam MM, Farag E, Hassan MM, Jaffrey SS, Atta M, Al-Marri AM, Al-Zeyara AM, Al Romaihi H, Bansal D, Mkhize-Kwitshana ZL (2023) Rodent-borne zoonoses in Qatar: a possible one-health framework for the intervention of future epidemic. *One Health* 16:100517
- Kamezaki H, Shoji K, Uemura S, Tsuda Y (2022) Fauna and spatial distribution of house-dust mites in Japanese mattress. *Med Entomol Zool* 73:137–143
- Kamezaki H, Sato T, Takeshita M, Hashimoto T, Tsuda Y (2024) Density and distribution of house dust mites found in the bedding and hairs of indoor-reared dogs. *Med Entomol Zool* 75:147–151
- Kamran M, Khan EM, Alatawi FJ (2018) The spider mites of the genus *Eutetranychus* Banks (Acari, Trombidiformes, Tetranychidae) from Saudi Arabia: two new species, a re-description, and a key to the world species. *Zookeys* 799:47–88
- Kano Y, Murata K (2025) What did the patient really eat? Pancake syndrome. *Am J Med press*, 138, e148-e149
- Kardar HMA (1988) Three new species of Oribatei (Acari: Cryptostigmata) from Saudi Arabia. In: Channabasavanna GP, Viraktamath CA (eds) *Progress in Acarology*, vol 1. E. J. Brill, Leiden, pp 291–294
- Karuthedath I, Raghavan ST (2024) Fauna and distribution of house dust mites in two northern districts of Kerala, India. *Persian J Acarol* 13:93–103
- Kato T, Ito Y, Nakao H, Kadosaka T (2011) Tinnitus caused by tiny mites – suspected cases of erratic migration of *Otodectes cynotis* and a case of *Aleuroglyphus ovatus* infestation. *Med Entomol Zool* 62:199–204
- Kelaher J, Jogi R, Katta R (2005) An outbreak of rat mite dermatitis in an animal research facility. *Cutis* 75:282–286
- Khadadah M, Onadeko BO, Ezeamuzie CI, Mustafa HT, Marouf R, Sugathan TN (2000a) The association of skin test reactivity, total serum IgE levels, and peripheral blood eosinophilia with asthma in Kuwait. *J Asthma* 37:481–488

- Khadadah ME, Onadeko BO, Ezeamuzie CI, Maroof R, Mustafa HS, Sugathan TN (2000b) Studies of the relationship between allergen-specific IgE antibodies and skin test reactivity in patients with asthma in Kuwait. *Med Princ Pract* 9:260–267
- Khasawneh R, Al-Hiary M, Al-Abadi B, Bani-Salameh A, Al-Momani S (2019) Total and specific immunoglobulin E for detection of most prevalent aeroallergens in a Jordanian cohort. *Med Arch* 73:272–275
- Khreesha L, Ghunaim M, Ramzown M-A, Alkhouljah M, Tawalbeh M, Al-Iede M, Kanaan T, Alrabayah M, Eid S (2020) Minimizing the number of aeroallergen extracts in skin prick test in IgE-mediated allergic disorders in both adults and children in Jordan. *J Asthma Allergy* 13:315–321
- Khurram T, Missous G, van Panhuys N, Karim MY (2025) Aeroallergen sensitivity patterns in Gulf countries: a systematic review. *Clin Transl Allergy* 15:e70033
- Kim E-M, Kim JY, Kwak YS, Yi M-H, Yong T-S (2024) The storage mite *Tyrophagus putrescentiae* induces greater lung inflammation than house dust mites in mice. *Parasites, Hosts and Diseases* 62:365–377
- Klien K, Walzl MG (2010) Private matters of *Sancassania berlesei* (Acaridida, Acaridae). Testes, receptaculum seminis, ovary and the location of sperm. In: Sabelis, M.W., J. Bruin (eds) *Trends in Acarology*. Dordrecht, Springer, pp 85–89
- Klimek A, Chachaj B (2018) Colonization of hardwood and pine wood chips by mites (Acari), with particular reference to oribatid mites (Oribatida). *Folia For Pol Ser For* 60:22–33
- Klimov PB, Hubert J, Erban T, Perotti MA, Braig HR, Flynt A, He Q, Cui Y (2024) Genomic and metagenomic analyses of the domestic mite *Tyrophagus putrescentiae* identify it as a widespread environmental contaminant and a host of a basal, mite-specific Wolbachia lineage (supergroup Q). *Int J Parasitol* 54:661–674
- Kontschán J, Hornok S (2025) First report of *Macrocheles robustulus* (Berlese, 1904) (Acari: Mesostigmata: Macrochelidae) on the pet beetle, *Pachnoda marginata peregrina* Kolbe (Coleoptera: Melolontidae: Cetoninae). *Acarologia* 65:717–720
- Kontschán J, Urszán J, Hornok S (2022) First record of two new pet-associated parasitic mites (Acari Mesostigmata Laelapidae and Macronyssidae) from Hungary. *Redia* 105:17–19
- Koshak EA (2006) Skin test reactivity to indoor allergens correlates with asthma severity in Jeddah, Saudi Arabia. *Allergy Asthma Clin Immunol* 2:11–19
- Krantz GW (1978) *A manual of acarology*, 2nd edn. Oregon State University Book Stores, Corvallis
- Krantz GW, Walter DE (2009) *A Manual of Acarology*, Third Edition, edn. Texas Tech University Press, Texas
- Kronqvist M, Johansson E, Kolmodin-Hedman B, Öman H, Svartengren M, van Hage-Hamsten M (2005) IgE-sensitization to predatory mites and respiratory symptoms in Swedish greenhouse workers. *Allergy* 60:521–526
- Lai NK (1988) Study of dust mite allergy in the Guangzhou area. *Zhonghua Yu Fang Yi Xue Za Zhi* 22:347–349
- Lee WK, Cho BK (1984) An ecological study on the house dust mite. *Korean J Dermatol* 22:286–294
- Lestringant GG, Bener A, Frossard PM, Abdulkhalik S, Bouix G (1999) A clinical study of airborne allergens in the United Arab Emirates. *Allerg Immunol (Paris)* 31:263–267
- Lezcano JJ, Murgas IL, Barrera OM, Miranda RJ (2020) House dust mites (Acari: Astigmata) from mattresses in Panama. *Acarologia* 60:576–586
- Li CP, Cui YB, Wang J, Yang QG, Tian Y (2003) Acaroid mite, intestinal and urinary acariasis. *World J Gastroenterol* 9:874–877
- Li C-p, He J, Jiang J-j, Wang H-y (2005) Composition and diversity of acaroid mite community in different environments in Huainan City. *Chin J Parasitol Parasit Dis* 23:460–462
- Liguori G, Ceccarelli MT, Mellino M, Marinelli P (1989) Recenti esperienze sugli acari dei prodotti immagazzinati [Recent experience with mites in stored products]. *Ann Ig* 1:237–246
- Liu X-y, Wu J, Wang B, Li M, Ran P-x, Liu Z-g (2010) Investigation on house dust mite in different geographical regions of China. *Chin J Zoonoses* 26:310–314
- Mahakittikun V, Boitano JJ, Ninsanit P, Wangapai T, Ralukruedek K (2011) Effects of high and low temperatures on development time and mortality of house dust mite eggs. *Exp Appl Acarol* 55:339–347
- Mahboub B, Al-Hammadi S, Prakash VP, Sulaiman N, Blaiss MS, Al Redha A, Vats DM (2014) Prevalence and triggers of allergic rhinitis in the United Arab Emirates. *World Allergy Organ J* 7:19
- Mahunka S (2009) Oribatid mites from the Arabian Peninsula, including further records from Socotra (Acari: Oribatida) (*Acarologica Genavensia CXII*). *Rev Suisse Zool* 116:257–274
- Malainual N, Vichyanond P, Phanurai P (1995) House-dust mite fauna in Thailand. *Clin Exp Allergy* 25:554–560
- Mangová B, Sládkovičová VH, Krumpál M, Kozánek M (2019) The impact of different urban conditions on structural characteristics of oribatid mite communities. *Biol (Bratisl)* 74:153–168

- Mariana A, Ho TM, Gendeh BS, Iskandar H, Zainuldin-Taib M (2000) First report on sensitization to allergens of a house dust mite, *Suidasia pontifica* (Acari: Saprogllyphidae). *Southeast Asian J Trop Med Public Health* 31:722–723
- Marín J, Ocete R, Pedroza M, Zalacain A, de Miguel C, López MA, Salinas MR (2009) Influence of the mite *Carpoglyphus lactis* (L) on the aroma of pale and dry wines aged under flor yeasts. *J Food Compos Anal* 22:745–750
- Mašán P (2003) Macrochelid Mites of Slovakia (Acari, Mesostigmata, Macrochelidae). Institute of Zoology, Slovak Academy of Sciences, Bratislava
- Mashaly AMA, Shoker NI, Negm MW (2011) Observations on mites (Arachnida: Acari) associated with three ant species (Hymenoptera: Formicidae) from Saudi Arabia. *Pak J Zool* 43:1095–1100
- Matsumoto C, Kawakami Y, Ishibashi O, Sagara I, Sakaguchi M, Harada Y, Takahashi M, Niiyama S, Inui T, Fukuda H (2025) A clinical case of anaphylaxis after eating oatmeal contaminated with booklice (*Liposcelis bostrychophila*). *J Dermatol* 52:146–149
- Maurya KR, Jamil Z (1982) Distribution of the cryptostigmatid, mesostigmatid and prostigmatid mites (Acarina) in the houses of Northeastern and Northern India. *Biol Membr* 7:131–138
- Maurya KR, Jamil Z, Dev B (1983) Prevalence of astigmatid mites Acarina in the domestic environment of northeastern and northern India. *Biol Membr* 8:207–215
- Melekhina EN (2023) Lichen-associated oribatid mites in the Taiga zone of northeast European Russia: taxonomical composition and geographical distribution of species. *Diversity* 15:599
- Melekhina EN, Korolev AN, Selivanova NP (2023) Oribatid mites (Oribatida) associated with nests of hollow-nesting birds, on the example of a model species, the European pied flycatcher (*Ficedula hypoleuca*), in the Taiga forests of the European north-east of Russia. *Diversity* 15:765
- Meteo Portal (2017) National Center for Meteorology (NCM). Saudi Arabia. <https://meteo.ncm.gov.sa/en/account/login/>
- Michaud JP, Schoenly KG, Moreau G (2012) Sampling flies or sampling flaws? Experimental design and inference strength in forensic entomology. *J Med Entomol* 49:1–10. <https://doi.org/10.1603/ME10229>
- Mirza JH, Elgoni NA, Kamran M, Alatawi FJ (2025) Mites of the genus *Bryobia* (Acari, Tetranychidae): taxonomic notes on some species and a diagnostic key to the world species. *Zookeys* 1239:51–73
- Modak A, Saha GK, Tandon N, Gupta SK (2004) Faunal diversity and habitat preference of house dust mites in West Bengal in relation to nasobronchial allergic disorders. *Rec Zool Surv India* 102:137–146
- Morsy TA, Zohdi HW, Abdalla KF, El Fakahani AF, Ibrahim AA, Khalil HT (1994) Isolation of three species of mites from house dust of atopic dermatitis patients in Qalyobia Governorate, Egypt. *J Egypt Soc Parasitol* 24:323–331
- Mulhall J, Conlon DN (2019) Oral mite anaphylaxis; a case series from an Irish centre. *Allergy* 74:821–821
- Mumcuoglu KY, Lutsky I (1990) A prevalence survey of poultry house mites in Israel. *Acarologia* 31:51–56
- Mumcuoglu KY, Shalom U (2013) The clover mite, *Bryobia praetiosa* Koch (Acari: Tetranychidae) as a nuisance inside a human habitation in Israel. *Acarina* 21:81–83
- Mutlu D, Akdemir C, Uzunoglu E, Direkci S, Guler NC (2019) Giresun Ilinde Ev Tozu Akarlarinin Yayginligi ve Epidemiyolojisi Uzerine Arastirmalar [The investigation of the prevalence and epidemiology of house dust mites in Giresun]. *Turk Parazitol Derg* 43:78–82
- Navarro JM, Mendoza Meza DL, Bermúdez DM (2008) Identificación de ácaros del polvo casero en colchones y almohadas de niños alérgicos de Santa Marta, Colombia [Identification of home dust mites in mattresses and pillows of allergic children from Santa Marta, Colombia]. *Rev Fac Cienc Salud* 5:24–31
- Ogata M, Suzuki T, Itagaki H, Ishida F, Nakai T (1978) Mites of household pet animals and their pathogenic effects on the owners. *Med Entomol Zool* 29:267–268
- Ogata M, Itagaki H, Suzuki T, Ishida F (1980) Mites of household pet animals and their pathogenic effects on the owners. *Jpn J Sanit Zool* 31:67–68
- Oshima S (1970) Studies on the mite fauna of the house-dust of Japan and Taiwan with special reference to house-dust allergy. *Jpn J Sanit Zool* 21:1–17
- Ostovan H (1993) [Faunistic study of stored product mites in Kazerun and biology of important species]. *Tarbiat Moda res University*
- Ou C, Chen Q, Hu X, Zeng Y, Zhang K, Hu Q, Weng Q (2024) Mycophagous mite, *Tyrophagus putrescentiae*, prefers to feed on entomopathogenic fungi, except *Metarhizium* generalists. *Microorganisms* 12:1042
- Ouchi T, Miyamoto J, Ishii A (1976) Mite fauna of room dust collected from newly constructed apartment houses. *Med Entomol Zool* 27:427–429
- Oudemans AC (1897) List of Dutch Acari, seventh part: Acaridae Latr. 1806, and Phytoptidae Pagenst. 1861, with synonymical remarks en description of new species etc. *Tijdschr Entomol* 40:250–260
- Parkin EA (1958) Some insect invaders of domestic premises. *Ann Appl Biol* 46:119–123
- Pei J, Gong J, Wang Z (2020) Risk prediction of household mite infestation based on machine learnings. *Build Environ* 183:107154
- Perez I Pérez-Iñigo C (1993) Acari: Oribatei, Poronota, Fauna Ibérica, vol 3. CSIC, Madrid, p 320

- Perotti MA (2001) Prey location and predation rates of predatory mites (Acari: Macrochelidae) on immature stages of pest flies (Diptera: Muscidae). *Syst Appl Acarol* 6:27–33
- Perotti MA, Braig HR (2009) Phoretic mites associated with animal and human decomposition. *Exp Appl Acarol* 49:85–124
- Perotti MA, Braig HR (2019) Acarology in crimino-legal investigations: The human acarofauna during life and death. In: Byrd JH, Tomberlin JK (eds) *Forensic Entomology: The utility of arthropods in legal investigations*, 3rd edn. Taylor & Francis, Boca Raton, pp 483–495
- Perotti MA, Goff ML, Baker AS, Turner BD, Braig HR (2009) Forensic acarology, an introduction. *Exp Appl Acarol* 49:3–13
- Petrova AD, Zheltikova TM (2000) Long-term dynamics and structure of house dust mite community in Moscow (Acariformes, Astigmata). *Zool Zh* 79:1402–1408
- Pike AJ, Wickens K (2008) The house dust mite and storage mite fauna of New Zealand dwellings. *N Z Entomol* 31:17–22
- Podder S, Biswas H, Saha GK (2021) A faunistic survey of house dust mites of Kolkata, West Bengal, India. *Acarol Stud* 3:22–31
- Poggioli NG, Taboada JL (2021) Occupational ear acariasis. *Eur Ann Otorhinolaryngol Head Neck Dis* 138:421–422
- Ponnudurai G, Rani N, Velusamy R, Anbarasi P (2015) A note on tropical rat mite (*Ornithonyssus bacoti*) infestation among workers in poultry farms of Namakkal, Tamil Nadu. *J Vet Parasitol* 29:124–126
- Poza Guedes P, Sanchez Machin I, Matheu V, Iraola V, Gonzalez R, Perez (2016) Role of predatory mites in persistent nonoccupational allergic rhinitis. *Can Respir J* 2016, 5782317
- Preiser-Funke T, Bergmann K-C (2023) Oral mite anaphylaxis (pancake syndrome) caused by storage mite *Acarus siro* and its treatment with allergen immunotherapy. *Allergol Sel* 7:113–115
- Puerta L, Fernández-Caldas E, Lockey RF, Caraballo LR (1993) Sensitization to *Chortoglyphus arcuatus* and *Aleuroglyphus ovatus* in *Dermatophagoides* spp allergic individuals. *Clin Exp Allergy* 23:117–123
- Qu SX, Li HP, Zhu JC, Liu JJ, Luo X, Ma L, Wang Q, Chen KP (2023) Growth, development, and host preference of *Histiostoma feroniarum* (Acaridida: Histiostomatidae): effects of temperature and types of mushroom cultivar. *Bull Entomol Res* 113:412–418
- Quek Y-W, Chuang C-Y, Pan H-H, Sun H-L, Lue K-H, Ku M-S (2019) Boys with allergic rhinitis who were sensitized to *Blomia tropicalis* are the most vulnerable to air pollutants. *Am J Rhinol Allergy* 33:730–736
- Rack G (1984) Systematik, Morphologie und Biologie von Milben (Acari) in Häusern und Vorräten sowie Milben von medizinischer Bedeutung. Teil IV: Die Grasmilbe *Bryobia cristata* (Dugès, 1834) und Grasmilbenplagen [Systematics, morphology and biology of mites (Acari) in houses and stores and mites of medical importance. Part IV: The grass mite *Bryobia cristata* (Dugès, 1834) and plagues of grass mites]. *Prakt Schadlbekämpfer* 36:133–136
- Reboux G, Valot B, Rocchi S, Scherer E, Roussel S, Millon L (2019) Storage mite concentrations are underestimated compared to house dust mite concentrations. *Exp Appl Acarol* 77:511–525
- Ree H-I, Jeon S-H, Lee I-Y, Hong C-S, Lee D-K (1997) Fauna and geographical distribution of house dust mites in Korea. *Korean J Parasitol* 35:9–17
- Rezk HA, Abd El-Hamid MM, Abd El-Latif MA (1996) House dust mites in Alexandria region, Egypt. *Alexandria J Agric Res* 41:209–216
- Rosa AE, Flechtmann CHW (1979) Mites in house dust from Brazil. *Int J Acarol* 5:195–198
- Rostom ZMF (1993) Survey of some granivorous and non-granivorous insects and mites of stores in Saudi Arabia. *J Stored Prod Res* 29:27–31
- Saleh SM, El-Helaly MS, El-Gayar FH (1985) Survey on stored product mites of Alexandria (Egypt). *Acarologia* 26:87–93
- Samšić K (1979) Acari from *Musca domestica*. In: Wittmer W, Büttiker W (eds) *Fauna Saudi Arab*, vol 1. Karger, Basel, pp 69–74
- Sánchez-Borges M, Capriles-Hulett A, Fernández-Caldas E (2020) Oral mite anaphylaxis: who, when, and how? *Curr Opin Allergy Clin Immunol* 20:242–247
- Sanchez-Covisa A, Rodríguez-Rodríguez JA, la De Torre F, Garcia-Robaina JC (1999) Faune d'acariens de la poussière domestique de l'île de Tenerife [Mite fauna of house dust of the island of Tenerife]. *Acarologia* 40:55–58
- Sanyal AK (2004) Notes on the ecology of soil mites (Acari) in two contrasting sites of Schirmacher Oasis, East Antarctica. *Acarina* 12:151–157
- Sargison ND, Chaudhry U, Costa-Junior L, Kutcher JR, Li K, Sargison FA, Zahid O (2025) The diagnosis and vector potential of *Ornithonyssus bacoti* tropical rat mites in northern Europe. *Vet Parasitol Reg Stud Rep* 58:101204


- Sattar HA, Mobayed H, Al-Mohammed AA, Ibrahim AS, Jufairi AA, Balamurugan P, Mary VP, Bener A (2003) The pattern of indoor and outdoor respiratory allergens in asthmatic adult patients in a humid and desert newly developed country. *Eur Ann Allergy Clin Immunol* 35:300–305
- Saw MK, Bharati SK, Sinha SK (2018) Faunal diversity of house dust mite in grocery shops in coal mining area of Dhanbad, India. *J Exp Zool India* 21:1261–1266
- Saw MK, Bharati SK, Sinha SK (2024) Assessment of house dust mites prevalence in the college classrooms of Dhanbad, India. *J Exp Zool India* 27:1701–1704
- Saxena RK, Jamil Z, Saxena KC, Saxena RP (1980) Contribution of *Caloglyphus berlesesi* in house dust atopy of Lucknow region, India. *Biol Membr* 5:122–128
- Schrank FvP (1781) *Envmratio Insectorvm Avstriae Indigenorvm* [A List of Insects Native to Austria]. Augsburg, Germany, Widow of E. Klett and Franck, p 513–514, 546, and foldout II
- Sharif HA, Salih E, Awad H, Albarazi R (2018) Common aeroallergens among children with allergies in the United Arab Emirates. *Indian J Allergy Asthma Immunol* 32:4–9
- Shinohara N, Hashimoto K, Kim H, Yoshida-Ohuchi H (2023) Fungi, mites/ticks, allergens, and endotoxins in different size fractions of house dust from long-term uninhabited houses and inhabited houses. *Build Environ* 229:109918
- Smiley RL (1991) Mites (Acari). In: Gorham, R.L. (ed) *Insect and Mite Pests in Food An Illustrated Key*. Agricultural Handbook, vol 1 and 2. Washington, D.C.: U.S. Government Printing Office, p 3–44, 483–526
- Solarz K (2009) Indoor mites and forensic acarology. *Exp Appl Acarol* 49:135–142
- Solarz K (2012) House Dust Mites and Storage Mites (Acari: Oribatida: Astigmata). Identification keys. Polish Academy of Sciences, Kraków
- Solarz K, Obuchowicz A, Asman M, Nowak W, Witecka J, Pietrzak J, Marek M, Lonak A, Stadnicka I, Hajduga-Stasko B (2021) Abundance of domestic mites in dwellings of children and adolescents with asthma in relation to environmental factors and allergy symptoms. *Sci Rep* 11:18453
- Solarz K, Pajak K, Pawelczyk O, Bobinski R, Ciecchacka M, Dutka M, Ulman-Wlodarz I (2022) Mites occurring in farm buildings as allergic agents and indicators in forensic analyses. *Acarologia* 62:36–47
- Soleimani-Ahmadi M, Zare M, Abtahi SM, Khazeni A (2017) Species identification and prevalence of house dust mites as respiratory allergen in kindergartens of the Bandar Abbas city. *Iran J Allergy Asthma Immunol* 16:133–139
- Solomon ME (1961) Mites in houses, shops and other occupied buildings. *Sanitarian Lond* 69:291–296
- Somorin AO, Hunponu-Wusu OO, Mumcuoglu Y, Heiner DC (1978) Mite allergens in Nigerians: studies on house dust mites in houses of allergic patients in Lagos. *Ir J Med Sci* 147:26–30
- StataCorp, L. (2014) Stata 13. StataCorp LP, College Station
- Stejskal V, Hubert J, Aulicky R, Kucerova Z (2015) Overview of present and past and pest-associated risks in stored food and feed products: European perspective. *J Stored Prod Res* 64:122–132
- Subias LS (2022) Listado sistemático, sinonímico y biogeográfico de los ácaros oribátidos (Acariformes, Oribatida) del mundo (excepto fósiles) [Systematic, synonymic and biogeographical list of oribatid mites (Acariformes, Oribatida) of the World (except fossils)]. *Monogr Electron S E A* 12:1–539
- Sudd JH (1952) Laboratory studies of adult female *Bdellonyssus bacoti* (Hirst, 1916) (Acarina, Parasitiformes). *Ann Trop Med Parasitol* 46:158–164
- Suliaman FA, Holmes WF, Kwick S, Khouri F, Ratard R (1997) Pattern of immediate type hypersensitivity reactions in the Eastern Province, Saudi Arabia. *Ann Allergy Asthma Immunol* 78:415–418
- Sultan DM, Khalil MM (2013) Dust mite allergens: A major risk factor in development of allergies in Dubai. In: Cozma V (ed) XI European Multicolloquium of Parasitology, Cluj-Napoca, Romania, 2012. Medimond, Bologna, pp 69–72
- Sulzbach A, Gedoz C, Quadri VD, Ohlweiler LM, Silva FR, da Silva GL, Johann L (2024) Impact of the mites *Dermanyssus gallinae* and *Tyrophagus putrescentiae* on production and health of laying hens. *Exp Appl Acarol* 92:61–85
- Suto C, Sakaki I, Itoh H (1991) Comparative studies on the population dynamics of house dust mites *Dermatophagoides farinae* and *D. pteronyssinus* (Acari: Pyroglyphidae) in homes in Nagoya. *Med Entomol Zool* 42:129–140
- Suto C, Sakaki I, Itoh H, Mitibata M (1992) Influence of floor levels on the prevalence of house dust mites in apartments in Nagoya, Japan. *Med Entomol Zool* 43:307–318
- Szafranek P, Lewandowski M (2017) Mite community on Polish mushroom farms. *Int J Acarol* 43:217–222
- Ta-Phaisach N, Konvipasruang P, Sringarm K, Chiu C-I, Attasopa K, Chanbang Y (2023) Mites (Acari) infesting stored animal feed products in northern Thailand. *Syst Appl Acarol* 28:1579–1596
- Takahashi K, Taniguchi M, Fukutomi Y, Sekiya K, Watai K, Mitsui C, Tanimoto H, Oshikata C, Tsuburai T, Tsurikisawa N, Minoguchi K, Nakajima H, Akiyama K (2014) Oral mite anaphylaxis caused by mite-contaminated okonomiyaki/ pancake-mix in Japan: 8 case reports and a review of 28 reported cases. *Allergol Int* 63:51–56

- Takaoka M (2000) Review on mite fauna of house dust in Japan. *J Acarol Soc Jpn* 9:93–103
- Tayeb MM (2015) Sensitization pattern to common aeroallergens at Makkah. *Allergy* 70:540–540
- Tayeb MMS (2020) Cat allergen sensitization which clinical allergies are commonly associated? A retrospective study (Saudi Arabia). *Afr J Respir Med* 15:38–42
- Thalappil S, Ibrahim T, Shibu S, Mobayed H, Al-Nesf M (2023) Aeroallergens sensitization pattern in atopic patients in Qatar: A retrospective study. *Qatar Med J* 2023, qmj.2023.sqac.2
- Theis J, Lavoipierre MM, La Perriere R, Kroese H (1981) Tropical rat mite dermatitis. Report of six cases and review of mite infestations. *Arch Dermatol* 117:341–343
- Thomas WR (2010) Geography of house dust mite allergens. *Asian Pac J Allergy Immunol* 28:211–224
- Toma T, Miyagi I, Kishimoto M, Nagama T, Tamanaha I (1993a) Fauna and seasonal appearances of mites in house dusts collected from the residences including asthmatic children in Okinawa Prefecture, Ryukyu Archipelago. *Medical Entomology and Zoology* 44:223–235
- Tubeilah FA, Luqman W (1987) Comparative evaluation for pollens and mite extract skin tests. *Trop Geogr Med* 39:233–237
- Vackova T, Pekar S, Klimov PB, Hubert J (2021) Sharing a bed with mites: preferences of the house dust mite *Dermatophagoides farinae* in a temperature gradient. *Exp Appl Acarol* 84:755–767
- Vackova T, Pekar S, Klimov PB, Hubert J (2023) Population growth and respiration in the dust mite *Dermatophagoides farinae* under different temperature and humidity regimes. *Exp Appl Acarol* 89:157–169
- van Bronswijk JEMH, AWAM De Cock (1973) The genus *Blomia* Oudemans (Acari: Glycyphagidae) II. Comparison of its species. *Acarologia* 15:490–505
- van Bronswijk JEMH, AWAM De Cock, Oshima S (1973) The genus *Blomia* Oudemans (Acari: Glycyphagidae). I. Description of *Blomia tropicalis* sp. n. from house dust in tropical and sub-tropical regions. *Acarologia* 15:477–489
- van Strien RT, Gehring U, Belanger K, Triche E, Gent J, Bracken MB, Leaderer BP (2004) The influence of air conditioning, humidity, temperature and other household characteristics on mite allergen concentrations in the northeastern United States. *Allergy* 59:645–652
- van Well C, Beck W (2022) Tropical rat mites (*Ornithonyssus bacoti*) in a pet African pygmy hedgehog - a case report. *Kleintierpraxis* 67:195–201
- Vasar M, Davison J, Sepp S-K, Mucina L, Oja J, Al-Quraishy S, Anslan S, Bahram M, Bueno CG, Cantero JJ, Decocq G, Fraser L, Hiesalu I, Hozzein WN, Koorem K, Meng Y, Moora M, Onipchenko V, Opik M, Partel M, Vahter T, Tedersoo L, Zobel M (2022) Global soil microbiomes: a new frontline of biomecology research. *Glob Ecol Biogeogr* 31:1120–1132
- Vidal-Quist JC, Ortego F, Lambrecht BN, Rombauts S, Hernandez-Crespo P (2025) Stage-specific transcriptomic analysis reveals insights into the development, reproduction and biological function of allergens in the European house dust mite *Dermatophagoides pteronyssinus*. *BMC Genomics* 26:527
- Villanueva AC, Astos EC, Temoche LC (2020) Determination of the presence of *Ornithonyssus* spp. in breeder hens and roosters in three genetic lines on a farm in Solano county, California, USA. *Rev Patol Trop* 49:55–62
- Vobrázková E, Samsňák K, Špičák V (1979) Allergogenous mites (Acari: Pyroglyphidae) in private recreation houses. *Folia Parasitol* 26:343–349
- Walter DE, Campbell NJH (2003) Exotic vs endemic biocontrol agents: would the real *Stratiolaelaps miles* (Berlese) (Acari: Mesostigmata: Laelapidae) please stand up? *Biol Control* 26:253–269
- Walter DE, Lindquist EE, Smith IM, Cook DR, Krantz GW (2009) Order Trombidiformes. In: Krantz GW, Walter DE (eds) *A Manual of Acarology*, 3rd edn. Texas Tech University, Lubbock, pp 233–420
- Wang X-C, Li C-P (2012) The breeding density and the major allergens of acaroid mites in different motor habitats. *Zhongguo Meijiieshengwuxue Ji Kongzhi Zazhi* 23:228–230
- Wang Q, He W, Zhou Y, Feng R, Wang Y, Liu L, Yuan Y, Dai J, Liu Y, Zhang X (2025) Polystyrene nanoplastics aggravate house dust mite induced allergic airway inflammation through EGFR/ERK-dependent lung epithelial barrier dysfunction. *Ecotoxicol Environ Saf* 298:118329
- Wen DC, Shyur SD, Ho CM, Chiang YC, Huang LH, Lin MT, Yang HC, Liang PH (2005) Systemic anaphylaxis after the ingestion of pancake contaminated with the storage mite *Blomia freemani*. *Ann Allergy Asthma Immunol* 95:612–614
- Wharton GW (1976) House dust mites. *J Med Entomol* 12:577–621
- Wu Y, Yang F (2025) Co-exposure to microplastics enhances the allergenic potentials of house dust mite allergen Der p 1. *Environ Res* 277:121613
- Xiong Q, Wan A-Y, Liu X, Fung C-H, Xiao X, Malainual N, Hou J, Wang L, Wang M, Yang KY, Cui Y, Leung E-H, Nong W, Shin S-K, Au S-N, Jeong KY, Chew F-T, Hui J-L, Leung T-F, Tungtrongchitr A, Zhong N, Liu Z, Tsui S-W (2022) Comparative genomics reveals insights into the divergent evolution of astigmatic mites and household pest adaptations. *Mol Biol Evol* 39:msac097
- Yoshikawa M (1987) Feeding of *Cheyletus malaccensis* (Acari, Cheyletidae) on human-body fluids. *J Med Entomol* 24:46–53

- Yousif IO, Rahamathullah N, Khan NS, Mesbahuzzaman M, Abdulrazzaq M, Alobeidli FA, Kandakurti PK (2025) Prevalence of immunoglobulin E sensitization to various inhaled allergens among the residents of the Northern United Arab Emirates. Automated evaluation of a membrane-based test system. *Saudi Med J* 46:282–291
- Yu J-M, Luo Q-H, Sun J-L, Shi C-L, Yin J, Zhou Y-L, Tang R, Zhang H, Yu Z, Chen M (2015) Diversity of house dust mite species in Xishuangbanna Dai, a tropical rainforest region in Southwest China. *BioMed Res Int* 2015, 421716
- Yu J-J, O Connor BM, Sarker S, Toledo A, Pan X, Wang C (2025) Prevalence and diversity of mite ecto-parasites of house mice (*Mus musculus* L.) in urban environments in New Jersey, USA. *Pest Manag Sci* 81:3293–3301
- Zahraldin K, Chandra P, Tuffaha A, Ehlayel M (2021) Sensitization to common allergens among children with asthma and allergic rhinitis in Qatar. *J Asthma Allergy* 14:287–292
- Zaki WM, Salem HA, El Azizy HM, Ajeebi ME (2019) CD 14 promoter polymorphisms associated with different allergic phenotypes and modulated by house dust mite allergy. *Parasitol United J* 12:26–33
- Zakzouk SM, Gad-El-Rab MO (1996) A study of clinical and allergic aspects of rhinitis patients in Riyadh. *Ann Saudi Med* 16:550–553
- Zare M, Hassani-Azad M, Soleimani-Ahmadi M, Majnoon R (2021) The influence of indoor environmental factors on the prevalence of house dust mites as aeroallergens in Bandar Abbas student dormitories, South of Iran, 2019. *J Med Entomol* 58:1865–1873
- Zeytun E, Doğan S, Aykut M, Özçiçek F, Ünver E, Özçiçek A (2015) House dust mites in Erzinean province. *Türk Parazitol Derg* 39:124–130
- Zeytun E, Doğan S, Özçiçek F, Ünver E, Dilkaraoğlu S (2016) Comparison of living and bedrooms in terms of house dust mites in the province of Erzincan, Turkey. *J Med Entomol* 53:26–30
- Zhang Z-Q (1998) Review biology and ecology of trombidid mites (Acari: Trombidioidea). *Exp Appl Acarol* 22:139–155
- Zhou Y, Zhu X, Ren Y, Hui Y (2022) Distribution of *Cheyletus* mites in dust samples. *Eur Zool J* 89(1):238–1,245

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Authors and Affiliations

Riyadh Hussain<sup>1,2</sup> · S. Aeban<sup>1,2</sup> · Medjedline Hani<sup>1</sup> · Henk R. Braig<sup>3</sup> ·  
M. Alejandra Perotti<sup>1</sup> 

✉ M. Alejandra Perotti  
m.a.perotti@reading.ac.uk

Riyadh Hussain  
riyad\_1430@yahoo.com

Medjedline Hani  
medjedlinehani@gmail.com

Henk R. Braig  
hrbraig@icloud.com

<sup>1</sup> Ecology and Evolutionary Biology, School of Biological Sciences, University of Reading, Whiteknights Campus, HLS Bld, Reading RG6 6AS, UK

<sup>2</sup> Department of Medical Molecular Laboratory, Security Forces Hospital Makkah, Makkah, Saudi Arabia

<sup>3</sup> Institute and Museum of Natural Sciences (IMCN), National University of San Juan (UNSJ), San Juan, Argentina