

# Gyaan Roots Classes

“From Roots to Result”

Tilmapur Chauraha, Ashapur, Varanasi

Assignment Sheet

Subject:-Physics

Class:-X

CHAPTER

## LIGHT – REFLECTION AND REFRACTION

### LIGHT

An object reflects light that falls on it. This reflected light when received by our eyes, enables us to see things.

### Reflection of light

Reflection of light is the phenomenon of bouncing back of light in the same medium on striking the surface of any object.

There are two types of reflection:

1. Regular reflection or Specular Reflection
2. Irregular reflection or Diffuse Reflection

### Regular Reflection


When the reflecting surface is smooth and well polished, the parallel rays falling on it are reflected parallel to one another, the reflected light goes in one particular direction. This is Regular reflection or Specular reflection see below... This is Regular reflection or Specular reflection see below figure.

### Irregular reflection

---

Prepared by: Pranjal Sir (☎ 9250471512)


When the reflecting surface is rough, the parallel rays falling on it reflected in different direction, as shown in below fig. Such a reflection is known as diffuse reflection or irregular reflection.



## LAWS OF REFLECTION OF LIGHT

According to the laws of Reflection of light,

- (i) The angle of incidence is equal to the angle of reflection, and
- (ii) The incident ray, the normal to the mirror at the point of incidence and the reflected ray, all lie in the same plane.



These laws of reflection are applicable to all types of reflecting surfaces including spherical surfaces.

## **OBJECTS**

Anything which gives out light rays either its own or reflected by it is called an object.

## **LUMINOUS OBJECTS**

The objects like the sun, other stars, electric bulb, tube light etc. which emit their own light are called luminous objects.

## **NON – LUMINOUS OBJECTS**

The objects which do not emit light themselves but only reflect or scatter the light which falls on them, are called non-luminous objects.

A flower, chair table, book, trees, etc are all non-luminous objects.

## **IMAGES**

Image is an optical appearance produced when light rays coming from an object are reflected from a mirror (or refracted through lens).

## **REAL IMAGE**

The image which can be obtained on a screen is called a real image. In a cinema hall, we see the images of actors and actress on the screen. So, the images formed on a cinema screen is an example of real images.


## **VIRTUAL IMAGE**

The image which cannot be obtained on a screen is called a virtual image. A virtual image can be seen only by looking into a mirror. The image of our face in a plane mirror is an example of virtual image.

## LATERAL INVERSION

When an object is placed in front of a plane mirror, then the right side of object appears to become the left side of image; and the left side of object appears to become the right side of image. This change of sides of an object and its mirror image is called lateral inversion.

The phenomenon of lateral inversion is due to the reflection of light.



## CLASSES

### CHARACTERISTICS OF IMAGES FORMED BY PLANE MIRROR

From Roots to Results

The characteristics of images formed by plane mirrors are:

1. The image of a real object is always virtual. Such an image cannot be taken on a screen.
2. The image formed in a plane mirror is always erect.
3. The size of the image in a plane mirror is always the same as the size of the object.
4. The image formed in a plane mirror is as far behind the mirror, as the object is in front of the mirror.


5. The image formed in a plane mirror is laterally inverted i.e. the left side of the objects becomes the right side of the image and vice-versa

## SPHERICAL MIRROR

A spherical mirror is that mirror whose reflecting surface is the part of a hollow sphere of glass. The spherical mirrors are of two types: Concave mirror and Convex mirror.


**CONCAVE MIRROR:** A concave mirror is that spherical mirror in which the reflection of light takes place at the concave surface (or bent-in surface).

**CONVEX MIRROR:** A convex mirror is that spherical mirror in which the reflection of light takes place at the convex surface (or bulging-out surface).




## PRINCIPAL FOCUS OF A SPHERICAL MIRROR

The principal focus of a concave mirror is a point on its principal axis to which all the light rays which are parallel and close to the axis, converge after reflection from the concave mirror. A concave mirror has a real focus. The focus of a concave mirror is in front of the mirror. Since a concave mirror converges a parallel beam of light rays, it is also called a converging mirror.



The principal focus of a convex mirror is a point on its principal axis from which a beam of light rays, initially parallel to the axis, appears to diverge after being reflected from the convex mirror. A convex mirror has a virtual focus. The focus of a convex mirror is situated behind the mirror. Since a convex mirror diverges a parallel beam of light rays, it is also called a diverging mirror.



**Focal Length:** The focal length of a spherical mirror is the distance between its pole and principal focus. It is denoted by the letter 'f'.

**Relation between Radius of curvature and focal length of a spherical mirror**

The focal length of a spherical mirror is equal to half of its radius of curvature.

$$f = \frac{R}{2}$$


In other words, for spherical mirrors of small apertures, the radius of curvature is found to be equal to twice the focal length.

$$R=2f$$


## RULES FOR OBTAINING IMAGES FORMED BY SPHERICAL MIRRORS

The intersection of at least two reflected rays gives the position of the image of the point object. Any two of the following rays can be considered for locating the image.


1. A ray parallel to the principal axis, after reflection, will pass through the principal focus in case of a concave mirror or appear to diverge from the principal focus in case of a convex mirror.



2. A ray passing through the principal focus of a concave mirror or a ray which is directed towards the principal focus of a convex mirror, after reflection, will emerge parallel to the principal axis.




3. A ray passing through the centre of curvature of a concave mirror or directed in the direction of the centre of curvature of a convex mirror, after reflection, is reflected back along the same path. The light rays come back along the same path because the incident rays fall on the mirror along the normal to the reflecting surface.



4. A ray incident obliquely to the principal axis, towards a point P (pole of the mirror), on the concave mirror or a convex mirror, is reflected obliquely. The incident and reflected rays follow the laws of reflection at the point of incidence (point P), making equal angles with the principal axis.

# GYANROOTS




## FORMATION OF DIFFERENT TYPES OF IMAGES BY A CONCAVE MIRROR

The type of image formed by a concave mirror depends on the position of the object in front of the mirror. There are six positions of the object:

### Case-1: Object is in between P and F

When an object is placed between the pole (P) and focus (F) of a concave mirror, the image formed is:


- (i) behind the mirror
- (ii) virtual and erect and
- (iii) larger than the object (or magnified)



### Case-2: Object is at the focus (F)

When an object is placed at the focus of a concave mirror, the image formed is:


- (i) at infinity
- (ii) real and inverted, and
- (iii) highly magnified (or highly enlarged)




**Case-3: Object is in between focus (F) and centre of curvature(C)**

When an object is placed between the focus (F) and centre of curvature(C) of a concave mirror, the image formed is

1. beyond the centre of curvature
2. real and inverted, and
3. larger than the object (or magnified)




**CYANOBOOTS**



#### Case-4: Object is at the centre of curvature(C)

When an object is placed at the centre of curvature of a concave mirror, the image formed is


1. at the centre of curvature
2. real and inverted, and
3. same size as the object



#### Case-5: Object is beyond the centre of curvature(C)

When an object is placed beyond the centre of curvature of a concave mirror, the image formed is


1. between the focus and centre of curvature
2. real and inverted, and
3. smaller than the object (or diminished)



### Case-6: Object is at infinity

When an object is placed at infinity of a concave mirror, the image formed is

1. between the focus and centre of curvature
2. real and inverted, and
3. much smaller than the object (or highly diminished)



### USES OF CONCAVE MIRRORS

1. Concave mirrors are commonly used in torches, search-lights and vehicles headlights to get powerful parallel beams of light.
2. Concave mirrors are used as shaving mirrors to see a larger image of the face.
3. The dentists use concave mirrors to see large images of the teeth of patients.

4. Concave mirrors are used as doctor's head mirrors to focus light coming from a lamp on to the body parts of a patient to be examined by the doctor.
5. Concave dishes are used in TV dish antennas to receive TV signals from the distant communications satellite.
6. Large concave mirrors are used to concentrate sunlight to produce heat in solar furnaces.


### FORMATION OF DIFFERENT TYPES OF IMAGES BY A CONVEX MIRROR

The type of image formed by a convex mirror depends on the position of object in front of the mirror. There are six positions of the object:

#### Case-1: Object is placed between P and infinity

When an object is placed between pole and infinity in front of a convex mirror, the image formed is


- i. between the pole and focus
- ii. virtual and erect, and
- iii. smaller than the object (or diminished)



## Case—2: Object is at infinity

When an object is placed at infinity of a convex mirror, the image formed is:

- i. behind the mirror at focus
- ii. virtual and erect, and
- iii. much smaller than the object (or highly diminished)



## USES OF CONVEX MIRRORS

Convex mirrors are commonly used as rear-view (wing) mirrors in vehicles. These mirrors are fitted on the sides of the vehicle, enabling the driver to see traffic behind him/her to facilitate safe driving. Convex mirrors are preferred because they always give an erect, though diminished, image. Also, they have a wider field of view as they are curved outwards. Thus, convex mirrors enable the driver to view much larger area than would be possible with a plane mirror.

**Q-1. Define the principal focus of a concave mirror.**

Ans. Light rays that are parallel to the principal axis of a concave mirror converge at a specific point on its principal axis after reflecting from the mirror. This point is known as the principal focus of the concave mirror.

**Q-2. The radius of curvature of a spherical mirror is 20 cm. What is its focal length?**

Ans. Here  $R=20$  cm

We know that  $f=\frac{R}{2} \Rightarrow f=\frac{20}{2}=10$  cm

**Q-3. Name a mirror that can give an erect and enlarged image of an object.**

Ans. When an object is placed between the pole and the principal focus of a concave mirror, the image formed is virtual, erect, and enlarged.

**Q-4. Why do we prefer a convex mirror as a rear-view mirror in vehicles?**

Ans. Convex mirrors give a virtual, erect, and diminished image of the objects placed in front of them. They are preferred as a rear-view mirror in vehicles because they give a wider field of view, which allows the driver to see most of the traffic behind him.

### **MIRROR FORMULA**

In a spherical mirror, the distance of the object from its pole is called the object distance ( $u$ ).

The distance of the image from the pole of the mirror is called the image distance ( $v$ ). The distance of the principal focus from the pole is called the focal length ( $f$ ). There is a relationship between these three quantities given by the mirror formula which is expressed as:

$$\frac{1}{f} = \frac{1}{v} + \frac{1}{u}$$

### **MAGNIFICATION**

Magnification produced by a spherical mirror gives the relative extent to which the image of an object is magnified with respect to the object size. It is expressed as the ratio of the height of the image to the height of the object. It is usually represented by the letter  $m$ . If  $h_1$  is the height

of the object and  $h_2$  is the height of the image, then the magnification  $m$  produced by a spherical mirror is given by:


$$m = \frac{\text{height of the image}}{\text{height of the object}} \Rightarrow m = \frac{h_2}{h_1}$$

The magnification  $m$  is also related to the object distance ( $u$ ) and image distance ( $v$ ). It can be expressed as:

$$m = \frac{h_2}{h_1} = -\frac{v}{u}$$

**Points to be remembered:**

- The height of the object is taken to be positive as the object is usually placed above the principal axis.
- The height of the image should be taken as positive for virtual images. However, it is to be taken as negative for real images.
- When the image is real, it is inverted so  $h_2$  is negative which results  $m$  is  $-ve$ . A negative sign in the value of the magnification indicates that the image is real.
- When the image is virtual, it is erect so  $h_2$  is positive which results  $m$  is  $+ve$ . A positive sign in the value of the magnification indicates that the image is virtual.




**SIGN CONVENTION FOR SPHERICAL MIRRORS**

The following sign convention is used for measuring various distances in the ray diagrams of spherical mirrors:

1. Object is always placed to the left of mirror.
2. All distances are measured from the pole of the mirror.

- Distances measured in the direction of the incident ray are positive and the distances measured in the direction opposite to that of the incident rays are negative.
- Distances measured above the principal axis are positive and that measured below the principal axis are negative.



# GYANROOTS

## CLASSES

### From Roots to Results

**Q-1. Find the focal length of a convex mirror whose radius of curvature is 32 cm.**

Ans. Here  $R=32$  cm

We know that  $f=\frac{R}{2} \Rightarrow f=\frac{32}{2} = 16$  cm

Hence, the focal length of the given convex mirror is 16 cm.

**Q-2. A concave mirror produces three times magnified (enlarged) real image of an object placed at 10 cm in front of it. Where is the image located?**

Ans. Here, magnification,  $m = -3$ ,

object distance,  $u = -10$  cm and

image distance,  $v = ?$

Putting these values in the magnification formula for a mirror, we get


$$m = -\frac{v}{u} \Rightarrow -3 = -\frac{v}{-10}$$

$$\Rightarrow v = -30 \text{ cm}$$


### NUMERICALS BASED ON CONVEX AND CONCAVE MIRROR

1. Find the focal length of a convex mirror of radius of curvature 1 m.
2. Focal length of a convex mirror is 50 cm. What is its radius of curvature?
3. Radius of curvature of a concave mirror is 25 cm. What is its focal length?
4. A concave mirror produces 10 cm long image of an object of height of 2 cm. What is the magnification produced?
5. An object 1 cm high is held near a concave mirror of magnification 10. How tall will be the image?
6. An object 4 cm in size is placed at a distance of 25 cm from a concave mirror of focal length 15 cm. Find the position, nature and height of the image.
7. A converging mirror forms a real image of height 4 cm, of an object of height 1 cm placed 20 cm away from the mirror. Calculate the image distance. What is the focal length of the mirror?
8. A 4.5 cm needle is placed 12 cm away from a convex mirror of focal length 15 cm. Give the location of the image and the magnification. Describe what happens as the needle is moved farther from the mirror.
9. An arrow 2.5 cm high is placed at a distance of 25 cm from a diverging mirror of focal length 20 cm. Find the nature, position and size of the image formed.

10. The image formed by a convex mirror of focal length 20 cm is a quarter of the object. What is the distance of the object from the mirror?
11. Find the size, nature and position of image formed by a concave mirror, when an object of size 1 cm is placed at a distance of 15 cm. Given focal length of mirror is 10 cm.
12. An object 2 cm high is placed at a distance of 16 cm from a concave mirror, which produces 3 cm high inverted image. What is the focal length of the mirror? Also, find the position of the image.
13. An erect image 3 times the size of the object is obtained with a concave mirror of radius of curvature 36 cm. What is the position of the object?
14. A 2.5 cm candle is placed 12 cm away from a convex mirror of focal length 30 cm. Give the location of the image and the magnification.
15. An object is placed in front of a concave mirror of focal length 20 cm. The image formed is 3 times the size of the object. Calculate two possible distances of the object from the mirror.
16. The image formed by a convex mirror is virtual, erect and smaller in size. Illustrate with figure.
17. A concave mirror produces a real image 10mm tall, of an object 2.5mm tall placed at 5cm from the mirror. Calculate focal length of the mirror and the position of the image.
18. An object is placed at a large distance in front of a convex mirror of radius of curvature 40cm. How far is the image behind the mirror?
19. An object is placed 15cm from a convex mirror of radius of curvature 90cm. Calculate position of the image and its magnification.
20. The image formed by a convex mirror of focal length 30cm is a quarter of the object. What is the distance of the object from the mirror?
21. When an object is placed at a distance of 60cm from a convex mirror, the magnification produced is  $\frac{1}{2}$ . Where the object should be placed to get a magnification of  $\frac{1}{3}$ ?




22. An object is placed 18cm front of a mirror. If the image is formed at 4cm to the right of the mirror, Calculate its focal length. Is the mirror convex or concave? What is the nature of the image? What is the radius of curvature of the mirror?
23. A convex mirror used for rear view on an automobile has a radius of curvature of 3m. If a bus is located at 5m from this mirror, find the position, nature and magnification of the image.
24. An object 3cm high is held at a distance of 50cm from a diverging mirror of focal length 25cm. Find the nature, position and size of the image formed.
25. A converging mirror of focal length 20cm forms an image which is two times the size of the object. Calculate two possible distances of the object from the mirror.
26. The linear magnification of a convex mirror of focal length 15cm is  $\frac{1}{3}$ . What is the distance of the object from the focus of the mirror?
27. The focal length of a convex mirror is 12.5 cm. How far is its centre of curvature (i) from the pole (ii) from the focus.
28. Find the focal length of a concave mirror that produces four times larger real image of an object held at 5cm from the mirror.
29. An object is held at 30cm in front of a convex mirror of focal length 15cm. At what distance from the convex mirror should a plane mirror be held so that images in the two images coincide with each other?
30. Draw any three ray diagrams to show how the size and nature of image of an object change when it move from centre of curvature of concave mirror towards the pole of the mirror.



## REFRACTION OF LIGHT

The change in direction of light when it passes from one medium to another obliquely, is called refraction of light. In other words, the bending of light when it goes from one medium to another obliquely is called refraction of light. The refraction takes place when light enters from air to water (see below figure).



The speed of light is different in different substances. The refraction of light is due to the change in the speed of light on going from one medium to another. Thus, when light goes from one medium to another, its speed changes. And this change in speed of light causes the refraction of light.

### MEDIUM

A transparent substance in which light travels is known as a medium. Medium can be divided into two types:


1. Optically rarer medium: A medium in which the speed of light is more is known as optically rarer medium (or less dense medium).

2. Optically denser medium: A medium in which the speed of light is less is known as o...  
 Optically denser medium: A medium in which the speed of light is less is known as optically rarer medium (or more dense medium).


Glass is an optically denser medium than air and water.

**RULES OF REFRACTION:**

Rule-1: When a light ray travels from a rarer medium to a denser medium, the light ray bends towards the normal.



Rule-2: When a light ray travels from a denser medium to a rarer medium, the light ray bends away from the normal.



## LAWS OF REFRACTION

According to laws of refraction of light:


- (i) The incident ray, the refracted ray and the normal to the interface of two transparent media at the point of incidence, all lie in the same plane.
- (ii) The ratio of sine of angle of incidence to the sine of angle of refraction is a constant, for the light of a given colour and for the given pair of media. This law is also known as Snell's law of refraction.

If  $i$  is the angle of incidence and  $r$  is the angle of refraction, then,  $\frac{\sin i}{\sin r} = \text{constant}$

This constant value is called the refractive index of the second medium with respect to the first.

## REFRACTIVE INDEX

The refractive index of a medium is defined as the ratio of speed of light in vacuum to the speed of light in the medium. It is represented by  $n$ .



## REFRACTIVE INDEX

The refractive index of a medium is defined as the ratio of speed of light in vacuum to the speed of light in the medium. It is represented by  $n$ .

➤ Refractive index of a medium,  $n = \frac{\text{speed of light in vacuum air}}{\text{speed of light in medium}} = \frac{c}{v}$

Both  $c$  and  $v$  are in m/s.

➤ Relative refractive index of medium 2 w.r.t. medium 1 is  ${}^1n_2 = \frac{n_2}{n_1} = \frac{v_1}{v_2}$

➤ Both  $v_1, v_2$  are in m/s,  $n_2, n_1$  have no units.

➤  ${}^1n_2 = \frac{1}{2n_1}$

➤ Snell's law of refraction: When light travels from medium 1 to medium 2, then

$${}^1n_2 = \frac{n_2}{n_1} = \frac{\sin i}{\sin r}$$

➤  ${}^a n_w = \frac{\text{real depth } x}{\text{apparent depth } a}$


Both  $x$  and  $y$  are in metre or in cm.

➤ Velocity of light in vacuum/air is  $c = 3 \times 10^8$  m/s.

## TWO REFRACTIONS THROUGH A RECTANGULAR GLASS SLAB

On passing through a rectangular glass slab, a ray of light suffers two refractions, one while going from air to glass and the other while going from glass to air. Light emerges from rectangular slab in a direction parallel to that in which it entered the glass slab. However the final emergent ray is slightly shifted sideways from the direction of original incident ray by a distance  $x$  called lateral shift.

The perpendicular distance between the original path of incident ray and the emergent ray coming out of the glass slab is called lateral displacement of the emergent ray of light. Lateral displacement depends mainly on three factors: angle of incidence, thickness of glass slab and refractive index of glass slab. Actually lateral displacement is directly proportional to (i) angle of incidence (ii) thickness of glass slab (iii) refractive index of glass slab. Higher the values of these factors, greater will be the lateral displacement. The angle which the emergent ray makes with the normal is called the angle of emergence.



**CLASSES**  
From Roots to Results

### CONDITION FOR NO REFRACTION

Refraction will not take place under the following two conditions:

**1-When light is incident normally on a boundary.**

A ray of light traveling in medium 1 falls normally. Therefore angle of incidence,  $i=0^\circ$ .

According to Snell's law:

$$\frac{\sin i}{\sin r} = \frac{n_2}{n_1}$$

$$\text{or } \sin r = \frac{n_1}{n_2} \sin i = \frac{n_1}{n_2} \sin 0^\circ = \frac{n_1}{n_2} \times 0 = 0$$

or  $r=0$

Thus, there is no deviation in the ray at the boundary. Hence, no refraction occurs when light is incident normally on a boundary of two media.

## 2. When the refractive indices of two media are equal.

When refractive index of medium 1 is equal to refractive index of medium 2 i.e.


$n_1 = n_2$ , then according to Snell's law:

$$\frac{\sin i}{\sin r} = \frac{n_2}{n_1} = 1$$

or  $\sin i = \sin r$

or  $i=r$

Hence no refraction occurs at the boundary that separates two media of equal refractive indices.



**1. A ray of light travelling in air enters obliquely into water. Does the light ray bend towards the normal or away from the normal? Why?**

The light ray bends towards the normal. When a ray of light travels from an optically rarer medium to an optically denser medium, it gets bent towards the normal. Since water is optically denser than air, a ray of light travelling from air into the water will bend towards the normal.

**2. Light enters from air to glass having refractive index 1.50. What is the speed of light in the glass? The speed of light in vacuum is  $3 \times 10^8$  m/s.**

Refractive index of a medium  $n_m$  is given by,

$$n_m = \frac{\text{speed of light in vacuum } c}{\text{speed of light in the medium } v}$$

Speed of light in vacuum,  $c = 3 \times 10^8$  m/s

Refractive index of glass,  $n_g = 1.50$

$$\text{Speed of light in the glass, } v = \frac{c}{n_g} = \frac{3 \times 10^8}{1.50} = 2 \times 10^8 \text{ m/s}$$

**3. Find out, from Table 10.3, the medium having highest optical density. Also find the medium with lowest optical density.**

Highest optical density = Diamond

Lowest optical density = Air

Optical density of a medium is directly related with the refractive index of that medium. A medium which has the highest refractive index will have the highest optical density and vice-versa. It can be observed from table 10.3 that diamond and air respectively have the highest and

lowest refractive index. Therefore, diamond has the highest optical density and air has the lowest optical density.

**4. You are given kerosene, turpentine and water. In which of these does the light travel fastest? Use the information given in Table 10.3.**

Speed of light in a medium is given by the relation for refractive index ( $n_m$ ). The relation is given as:

$$n_m = \frac{\text{speed of light in vacuum air} = c}{\text{speed of light in medium} = v}$$

It can be inferred from the relation that light will travel the slowest in the material which has the highest refractive index and travel the fastest in the material which has the lowest refractive index. It can be observed from table 10.3 that the refractive indices of kerosene, turpentine, and water are 1.44, 1.47, and 1.33 respectively. Therefore, light travels the fastest in water.

**5. The refractive index of diamond is 2.42. What is the meaning of this statement?**

Refractive index of a medium  $n_m$  is related to the speed of light in that medium  $v$  by the relation:

$$n_m = \frac{\text{speed of light in vacuum air} = c}{\text{speed of light in medium} = v}$$

Where,  $c$  is the speed of light in vacuum/air.

The refractive index of diamond is 2.42. This suggests that the speed of light in diamond will reduce by a factor 2.42 compared to its speed in air.

## NUMERICALS

1. Light travels through water with a speed of  $2.25 \times 10^8$  m/s. What is the refractive index of water?
2. Light travels from rarer medium 1 to a denser medium 2. The angle of incident and refraction are respectively  $45^\circ$  and  $30^\circ$ . Calculate the (i) refractive index of second medium with respect to the first medium and (ii) refractive index of medium 1 with respect to the medium 2.
3. A pond of depth 20cm is filled with water of refractive index  $4/3$ . Calculate apparent depth of the tank when viewed normally.
4. How much time will light take to cross 2mm thick glass pane if refractive index of glasses is  $3/2$ ?
5. Calculate speed of light in water of refractive index  $4/3$ .
6. A ray of light passes from air to glass ( $n=1.5$ ) at an angle of  $30^\circ$ . Calculate the angle of refraction.
7. A ray of light is incident on a glass slab at an angle of  $45^\circ$ . If refractive index of glass be 1.6, what is the angle of refraction?
8. The refractive index of diamond is 2.47 and that of glass is 1.51. How much faster does light travel in glass than in diamond?
9. The refractive index of glycerine is 1.46. What is the speed of light in air if its speed in glycerine is  $2.05 \times 10^8$  m/s?
10. The refractive index of glass is 1.6 and that of diamond is 2.4. Calculate (i) refractive index of diamond with respect to glass and (ii) refractive index of glass with respect to diamond.
11. A ray of light is travelling from glass to air. The angle of incidence in glass is  $30^\circ$  and angle of refraction in air is  $60^\circ$ . What is the refractive index of glass w.r.t air?
12. A ray of light is travelling from air to water. What is the angle of incidence in air, if angle of refraction in water is  $45^\circ$ ? Take refractive index of water = 1.32
13. A water tank appears to be 4 m deep when viewed from the top. If refractive index of water is  $4/3$ , what is the actual depth of the tank?

14. What is the real depth of a swimming pool when its bottom appears to be raised by 1m?

Given refractive index of water is  $\frac{4}{3}$ .

15. A jar 15 cm long is filled with a transparent liquid. When viewed from the top, its bottom appears to be 12cm below. What is the refractive index of the liquid?

## SPHERICAL LENSES


A lens is any transparent material (e.g. glass) of an appropriate shape that can take parallel rays of incident light and either converge the rays to a point or diverge the rays from a point.

A transparent material bound by two surfaces, of which one or both surfaces are spherical, forms a lens.

Some lenses will focus light rays to a single point. These lenses are called converging or concave lenses. Other lenses spread out the light rays so that it looks like they all come from the same point. These lenses are called diverging or convex lenses. Lenses change the direction of light rays by refraction. They are designed so that the image appears in a certain place or as a certain size. Lenses are used in eyeglasses, cameras, microscopes, and telescopes.


## CONVEX LENS

A lens may have two spherical surfaces, bulging outwards. Such a lens is called a double convex lens. It is simply called a convex lens. It is thicker at the middle as compared to the edges. Convex lens converges light rays. Hence it is called converging lens.



## CONCAVE LENS


A double concave lens is bounded by two spherical surfaces, curved inwards. It is thicker at the edges than at the middle. Such lenses diverge light rays and are called diverging lenses. A double concave lens is simply called a concave lens.



FROM ROOTS TO RESULTS


## TERMS RELATED TO SPHERICAL LENS

**Principal Axis:** The principal axis is the line which runs horizontally straight through the optical centre of the lens. It is also sometimes called the optic axis. In other words, an imaginary straight line passing through the two centres of the curvature of a lens is called its principal axis.



**Optical Centre:** The optical centre (O) of a convex lens is usually the centre point of the lens. The direction of all light rays which pass through the optical centre, remains unchanged.

**Centre of Curvature:** A lens has two spherical surfaces. Each of these surfaces forms a part of a sphere. The centers of these spheres are called **centres of curvature of the lens**. The centre of curvature of a lens is usually represented by the letter C. Since there are two centres of curvature, we may represent them as  $C_1$  and  $C_2$ .



**Aperture:** The effective diameter of the circular outline of a spherical lens is called its aperture. Lenses whose aperture is much less than its radius of curvature are called thin lenses with small aperture.


**Focus:** The focus or focal point of the lens is the position on the principal axis where all light rays that run parallel to the principal axis through the lens converge (come together) at a point. Since light can pass through the lens either from right to left or left to right, there is a focal point on each side of the lens ( $F_1$  and  $F_2$ ), at the same distance from the optical centre in each direction. (Note: the plural form of the word focus is foci.)

**Focal Length:** The focal length ( $f$ ) is the distance between the optical centre and the focal point.


### **RULES FOR OBTAINING IMAGES FORMED BY SPHERICAL LENSES**

The intersection of at least two refracted rays give the position of image of the point object. Any two of the following rays can be considered for locating the image.


1. A ray of light from the object, parallel to the principal axis, after refraction from a convex lens, passes through the principal focus on the other side of the lens, as shown in below figure. In case of a concave lens, the ray appears to diverge from the principal focus located on the same side of the lens, as shown in below figure.



2-A ray of light passing through a principal focus, after refraction from a convex lens, will emerge parallel to the principal axis. This is shown in below figure. A ray of light appearing to meet at the principal focus of a concave lens, after refraction, will emerge parallel to the principal axis. This is shown in below figure.



3-A ray of light passing through the optical centre of a lens will emerge without any deviation. This is illustrated in below figure.



**CLASSES**  
From Roots to Results


### FORMATION OF DIFFERENT TYPES OF IMAGES BY A CONVEX LENS

The type of image formed by a convex lens depends on the position of object in front of the lens. There are six positions of the object:

**Case-1: Object is in between optical centre(O) and focus (F1)**

When the object is placed between optical centre(O) and focus(F1), the image formed is:


- (i) behind the object (on the left side of lens)
- (ii) virtual and erect, and
- (iii) larger than the object (enlarged or magnified)



**Case-2: Object is at the focus (F1)**

When the object is placed at the focus (F<sub>1</sub>), the image formed is:


- (i) at infinity
- (ii) real and inverted, and
- (iii) highly enlarged



### Case-3: Object is in between F1 and 2F1

When the object is placed between  $F_1$  and  $2F_1$  in front of a convex lens, the image formed is:


- (i) beyond  $2F_2$ ,
- (ii) real and inverted, and
- (iii) larger than the object (or magnified).



### Case-4: Object is at $2F_1$

When the object is placed at a distance  $2f$  in front of convex lens, the image formed is:


- (i) at  $2F_2$  on the other side of the lens,
- (ii) real and inverted, and
- (iii) of the same size as the object.



### Case-5: Object is at beyond $2F_1$

When the object is placed beyond  $2F_1$  in front of the convex lens, the image formed is:

- (i) between  $F_2$  and  $2F_2$  on the other side of the lens,
- (ii) real and inverted, and
- (iii) smaller than the object (or diminished).




### Case-6: Object is at infinity

When the object is placed at the infinity, the image formed is:

- (i) at the focus  $F_2$ .
- (ii) real and inverted, and
- (iii) much smaller than the object (or highly diminished or point sized).

**GYANROOTS**  
CLASSES  
From Roots to Results




**Formation of Different Types of Images by a Concave Lens**

The type of image formed by a concave lens depends on the position of object in front of the lens. There are two positions of the object.

**Case-1: Object is at infinity**

When the object is placed at the infinity, the image formed is:

- (i) at the focus F1
- (ii) virtual and erect, and
- (iii) much smaller than the object (or highly diminished or point sized).




ults

## Case-2: Object is in between optical centre(O) and infinity

When the object is placed in between optical centre (O) and infinity, the image formed is

- (i) Between optical centre(O) and focus  $F_1$ .
- (ii) Virtual and erect, and
- (iii) Smaller than the object (or diminished)




### SIGN CONVENTION FOR SPHERICAL LENSES

While using the lens formula we must make use of proper sign convention while taking the values of object (u), image distance (v), focal length (f), object height (h) and image height (h'). The sign conventions are as follows:


1. All distances are measured from the optical centre of the lens.
2. The distances measured in the same direction as the incident light are taken positive.
3. The distances measured in the direction opposite to the direction of incident light are taken negative.
4. Heights measured upwards and perpendicular to the principal axis are taken positive.

5. Heights measured downwards and perpendicular to the principal axis are taken negative.



Consequences of new Cartesian sign convention:

- The focal length of a convex lens is positive and that of a concave lens is negative.
- Object distance  $u$  is always negative.
- The distance of real image is positive and that of virtual image is negative.
- The object height  $h$  is always positive. Height  $h'$  of virtual erect image is positive and that of real inverted image is negative.
- The linear magnification,  $m = h'/h$  is positive for a virtual image and negative for a real image.



### LENS FORMULA

Lens formula gives the relationship between object distance ( $u$ ), image-distance ( $v$ ) and the focal length ( $f$ ). The lens formula is expressed as

$$\frac{1}{f} = \frac{1}{v} - \frac{1}{u}$$

where 'u' is the distance of the object from the optical centre (O), 'v' is the distance of the image from the optical centre (O) and 'f' is the distance of the principal focus from the optical centre (O).

## MAGNIFICATION

The magnification produced by a lens, similar to that for spherical mirrors, is defined as the ratio of the height of the image and the height of the object. It is represented by the letter m. If h is the height of the object and h' is the height of the image given by a lens, then the magnification produced by the lens is given by

$$m = \frac{\text{Height of the image } h'}{\text{Height of the object } h} = \frac{h'}{h}$$

Magnification produced by a lens is also related to the object-distance u, and the image-distance v. This relationship is given by

$$\text{(magnification) } m = \frac{h'}{h} = \frac{v}{u}$$

### Points to be remembered

- If the magnification 'm' has a positive value, the image is virtual and erect. And if the magnification 'm' has a negative value, the image will be real and inverted.
- A convex lens can form virtual images as well as real images, therefore, the magnification produced by a convex lens can be either positive or negative.
- A convex lens can form images which are smaller than the object, equal to the object or bigger than the object, therefore magnification 'm' produced by a convex lens can be less than 1, equal to 1 or more than 1.

- A concave lens, however, forms only virtual images, so the magnification produced by a concave lens is always positive.
- A concave lens forms image which are always smaller than the object, so the magnification 'm' produced by a concave lens is always less than 1.

### NUMERICALS BASED ON CONVEX LENS

1. A convex lens of focal length 10 cm is placed at a distance of 12 cm from a wall. How far from the lens should an object be placed so as to form its real image on the wall?
2. If an object of 7 cm height is placed at a distance of 12 cm from a convex lens of focal length 8 cm, find the position, nature and height of the image.
3. An object 4 cm high is placed at a distance of 10 cm from a convex lens of focal length 20 cm. Find the position, nature and size of the image.
4. A small object is so placed in front of a convex lens of 5 cm focal length that a virtual image is formed at a distance of 25 cm. Find the magnification.
5. Find the position and nature of the image of an object 5 cm high and 10 cm in front of a convex lens of focal length 6 cm.
6. Calculate the focal length of a convex lens, which produces a virtual image at a distance of 50 cm of an object placed 20 cm in front of it.
7. An object is placed at a distance of 100 cm from a converging lens of focal length 40 cm. What is the nature and position of the image?
8. A convex lens produces an inverted image magnified three times of an object at a distance of 15 cm from it. Calculate focal length of the lens.
9. An object placed 4 cm in front of a converging lens produces a real image 12 cm from the lens. What is the magnification of the image? What is the focal length of the lens? Also draw the ray diagram to show the formation of the image.
10. A lens of focal length 20 cm is used to produce a ten times magnified image of a film slide on a screen. How far must the slide be placed from the lens?

11. Determine how far an object must be placed in front of a converging lens of focal length 10 cm in order to produce an erect image of linear magnification 4.
12. A convex lens of focal length 6 cm is held 4 cm from a newspaper, which has print 0.5 cm high. By calculation, determine the size and nature of the image produced.
13. A convex lens of focal length 0.10 m is used to form a magnified image of an object of height 5 mm placed at a... A convex lens of focal length 0.10 m is used to form a magnified image of an object of height 5 mm placed at a distance of 0.08 m from the lens. Find the position, nature and size of the image.
14. An erect image 2 cm high is formed 12 cm from a lens, the object being 0.5 cm high. Find the focal length of the lens.
15. The filament of a lamp is 80 cm from a screen and a converging lens forms an image of it on a screen, magnified three times. Find the distance of the lens from the filament and the focal length of the lens.
16. An object 2 cm tall is placed on the axis of a convex lens of focal length 5 cm at a distance of 10 cm from the optical centre of the lens. Find the nature, position and size of the image formed. Which case of image formation by convex lenses is illustrated by this example?
17. A converging lens of focal length 5 cm is placed at a distance of 20 cm from a screen. How far from the lens should an object be placed so as to form its real image on the screen?
18. An object 5 cm high is held 25 cm away from a converging lens of focal length 10 cm. Find the position, size and nature of the image formed. Also draw the ray diagram.
19. At what distance should an object be placed from a convex lens of focal length 18 cm to obtain an image at 24 cm from it on the other side? What will be the magnification produced in this case?
20. The magnification produced by a spherical lens is +2.5. What is the nature of image and lens?

21. What is the nature of the image formed by a convex lens if the magnification produced by a convex lens is +3?
22. What is the nature of the image formed by a convex lens if the magnification produced by a convex lens is  $-0.5$ ?
23. What is the position of image when an object is placed at a distance of 10 cm from a convex lens of focal length 10 cm?
24. Describe the nature of the image formed when an object is placed at a distance of 30 cm from a convex lens of focal length 15 cm.
25. At what distance from a converging lens of focal length 12 cm must an object be placed in order that an image of magnification 1 will be produced?

#### NUMERICALS BASED ON CONCAVE LENS

1. A concave lens produces an image 20 cm from the lens of an object placed 30 cm from the lens. Calculate the focal length of the lens.
2. The magnification of a spherical lens is  $+0.5$ . What is the nature of lens and image?
3. If an object is placed at a distance of 50 cm from a concave lens of focal length 20 cm, find the position, nature and height of the image.
4. An object is placed at a distance of 4 cm from a concave lens of focal length 12 cm. Find the position and nature of the image.
5. An object is placed at a distance of 50 cm from a concave lens produces a virtual image at a distance of 10 cm in front of the lens. Draw a diagram to show the formation of image. Calculate focal length of the lens and magnification produced.

6. A 50 cm tall object is at a very large distance from a diverging lens. A virtual, erect and diminished image of the object is formed at a distance of 20 cm in front of the lens. How much is the focal length of the lens?

7. A concave lens of focal length 15 cm forms an image 10 cm from the lens. How far is the object placed from the lens? Draw the ray diagram.

8. An object 60 cm from a lens gives a virtual image at a distance of 20 cm in front of the lens. What is the focal length of the lens? Is the lens converging or diverging? Give reasons for your answer.

9. A concave lens of 20 cm focal length forms an image 15 cm from the lens. Compute the object distance.


10. A concave lens has focal length 15 cm. At what distance should the object from the lens be placed so that it forms an image at 10 cm from the lens? Also find the magnification produced by the lens.

11. Calculate the image distance for an object of height 12 mm at a distance of 0.20 m from a concave lens of focal length 0.30 m and state the nature and size of the image.

12. A concave lens has focal length of 20 cm. At what distance from the lens a 5 cm tall object be placed so that it forms an image at 15 cm from the lens? Also calculate the size of the image formed.

13. An object is placed 20 cm from (a) a converging lens and (b) a diverging lens of focal length 15 cm. Calculate the image position and magnification in each case.

14. A 2.0 cm tall object is placed 40 cm from a diverging lens of focal length 15 cm. Find the position and size of the image.



15. Find the position and size of the virtual image formed when an object 2 cm tall is placed 20 cm from (a) diverging lens of focal length 40 cm and (b) converging lens of focal length 40 cm.

16. The magnification produced by a spherical lens is +0.75. What is the nature of image and lens?

17. The magnification produced by a spherical lens and a spherical mirror is +0.8. What is the nature of lens and mirror?

18. The magnification produced by a spherical lens and a spherical mirror is +2.0. What is the nature of lens and mirror?

19. The lens A produces a magnification of -0.6 whereas lens B produces magnification of +0.6. What is the nature of lens A and B.

20. An object is 2m from a lens which forms an erect image one-fourth (exactly) the size of the object. Determine the focal length of the lens. What type of the lens is this?

### POWER OF A LENS

The power of a lens is defined as the reciprocal of its focal length. It is represented by the letter P. The power P of a lens of focal length f is given by

$$P = \frac{1}{f}$$

The SI unit of power of a lens is 'diopetre'. It is denoted by the letter D. If f is expressed in metres, then, power is expressed in diopetres. Thus, 1 diopetre is the power of a lens whose focal length is 1 metre.  $1D = 1m^{-1}$ . The power of a convex lens is positive and that of a concave lens is negative.

Many optical instruments consist of a number of lenses. They are combined to increase the magnification and sharpness of the image. The net power (P) of the lenses placed in contact is given by the algebraic sum of the individual powers P<sub>1</sub>, P<sub>2</sub>, P<sub>3</sub>, ... as  $P = P_1 + P_2 + P_3 + \dots$

## NUMERICALS ON POWER OF LENS

1. A concave lens produces an image 20cm from the lens of an object placed 30cm from the lens. Calculate the power of the lens.
2. A convex lens is of focal length 10 cm. What is its power?
3. A person having a myopia eye uses a concave lens of focal length 50cm. What is the power of the lens?
4. A thin lens has a focal length of  $-25\text{cm}$ . What is the power of the lens and what is its nature?
5. A lens has a power of  $-2.5\text{ D}$ . What is the focal length and nature of the lens?
6. Find the power of a concave lens of focal length 2 m.
7. A convex lens forms a real and inverted image of needle at a distance of 50cm from the lens. If the image is of the same size as the needle, where is the needle placed in front of the lens? Also, find the power of the lens.
8. Two thin lenses of power  $+3.5\text{ D}$  and  $-2.5\text{ D}$  are placed in contact. Find the power and focal length of the lens combination.
9. A doctor has prescribed a corrective lens of power  $-1.5\text{ D}$ . Find the focal length of the lens. Is the prescribed lens is diverging or converging?
10. A concave lens of focal length 25 cm and a convex lens of focal length 20 cm are placed in contact with each other. What is the power of this combination? Also, calculate focal length of the combination.
11. A convex lens of focal length 20 cm is placed in contact with a concave lens of focal length 10cm. What is the focal length and power of the combination?
12. An object is placed at a distance of 50cm from a concave lens of focal length 30cm. Find the nature and position of the image.
13. An object of height 2 cm is placed at a distance of 15cm in front of a concave lens of power  $-10\text{D}$ . Find the size of the image.

14. A convergent lens of power 8D is combined with a divergent lens of power  $-10D$ . Calculate focal length of the combination.

15. A concave lens is kept in contact with a convex lens of focal length 20cm. The combination works as a converging lens of focal length 100cm. Calculate power of concave lens.

16. Find the focal length and nature of lens which should be placed in contact with a lens of focal length 10 cm so that the power of the combination becomes 5D.

17. A convex lens of power 3D is held in contact with a concave lens of power  $-1 D$ . A parallel beam of light is made to fall on the combination. At what distance from the combination will the beam get focussed?

18. A convex lens of focal length 25cm and a concave lens of focal length 10cm are placed in close contact with one another.

- a). What is the power of the combination?
- b). What is the focal length of the combination?
- c). Is this combination converging or diverging?


19. The power of a combination of two lenses X and Y is 5D. If the focal length of lens X be 15 cm, then

- a). calculate the focal length of lens Y.
- b). State the nature of the lens Y.

20. Two lenses A and B have focal lengths of +20cm and  $-10$  cm, respectively.

- a). What is the nature of lens A and lens B?
- b). What is the power of lens A and lens B?

What is the power of the combination if lenses A and B are held close together?



### 1. Define 1 dioptre of power of a lens.

Power of lens is defined as the reciprocal of its focal length. If  $P$  is the power of a lens of focal length  $F$  in metres, then

The S.I. unit of power of a lens is Dioptre. It is denoted by  $D$ .


$$P = \frac{1}{f(\text{in meters})}$$

1 dioptre is defined as the power of a lens of focal length 1 metre.

Hence,  $1 D = 1 \text{ m}^{-1}$

2. A convex lens forms a real and inverted image of a needle at a distance of 50 cm from it. Where is the needle placed in front of the convex lens if the image is equal to the size of the object? Also, find the power of the lens.

When an object is placed at the centre of curvature,  $2F_1$ , of a convex lens, its image is formed at the centre of curvature,  $2F_2$ , on the other side of the lens. The image formed is inverted and of the same size as the object, as shown in the given figure.



It is given that the image of the needle is formed at a distance of 50 cm from the convex lens. Hence, the needle is placed in front of the lens at a distance of 50 cm.

Object distance,  $u = -50$  cm

Image distance,  $v = 50$  cm

Focal length =  $f$

According to the lens formula,

$$\frac{1}{v} - \frac{1}{u} = \frac{1}{f}$$

$$\frac{1}{f} = \frac{1}{50} - \frac{1}{-50} = \frac{1}{50} + \frac{1}{50} = \frac{1}{25}$$

$$f = 25 \text{ cm} = 0.25 \text{ m}$$

$$\text{Power of the lens, } P = \frac{1}{f(\text{in meters})} = \frac{1}{0.25} = +4\text{D}$$


**3. Find the power of a concave lens of focal length 2 m.**

Focal length of concave lens,  $f = -2$  m

$$\text{Power of the lens, } P = \frac{1}{f(\text{in meters})} = \frac{1}{-2} = -0.5\text{D}$$

Here, negative sign arises due to the divergent nature of concave lens.


Hence, the power of the given concave lens is  $-0.5$  D.




**GYANROOTS**  
**CLASSES**  
From Roots to Results

# ASSIGNMENT QUESTIONS SET – 1

1. Define refraction.
2. Define refractive index.
3. What is the unit of refractive index?
4. List out the factors on which the refractive index of a medium depends.
5. Define angle of incidence.
6. What is the angle of incidence if a ray of light is incident normal to the surface separating the two media?
7. What is a lens?
8. What is a concave lens?
9. What is the nature of the focus of a concave lens?
10. What type of image is formed by a concave lens?
11. A thin lens has a focal length  $f = -12$  cm. Is it convex or concave lens?
12. A lens forms an erect image for all positions of the object in front of it. Is the lens convex or concave?
13. Where should an object be placed so that a real and inverted image of same size is obtained using a convex lens?
14. Write the relation between  $u$ ,  $v$  and  $f$  of a thin lens.
15. What is the sign of  $u$ ,  $v$  and  $f$  for a convex lens according to Cartesian sign convention?
16. An object of height  $1$  m is placed at a distance of  $2f$  from a convex lens. What is the height of the image formed?
17. Define power.
18. What is least distance of distinct vision?
19. What happens when a ray of light passes through the optical centre of a lens?
20. State the laws of refraction.
21. Diagrammatically represent the refraction of light through a rectangular glass slab.
22. Define convex lens. Why is it referred to as converging lens?



23. Draw a diagram to show the second principal focus of a convex lens.
24. Distinguish between a convex and a concave lens.
25. Draw a ray diagram to show the refraction of light when it passes through the optic centre of a convex lens.
26. List out the uses of convex lenses.
27. With the help of a ray diagram show how an object gets magnified in a simple microscope.
28. Define the power of a lens. What is its unit?
29. With the help of a diagram explain how light gets refracted when it passes through a rectangular glass slab.
30. An object is placed at a distance of 50 cm from a concave lens of focal length 20 cm. Find the nature and position of the image.
31. An object is placed 50 cm from a lens which produces a virtual image at a distance of 10 cm in front of the lens. Draw a diagram to show the formation of image and calculate the focal length of the lens.
32. An object of height 4 cm is placed at a distance of 10 cm from a convex lens of focal length 20 cm. Find the position, nature and size of the image.
33. What is the power of a lens having a focal length of a) 50 cm b) -50 cm
34. Draw a ray diagram to show the position and nature of the image formed by a convex lens when the object is placed
- a) at  $2F_1$
  - b) between  $F_1$  and  $2F_1$
  - c) beyond  $2F_1$
- 

35.State and verify Snell’s law.

36.The bending of a beam of light when it passes obliquely from one medium to another is known as \_\_\_.

1. reflection
2. refraction
3. dispersion
4. deviation

37.The part of the lens through which the ray of light passes without suffering deviation is called \_\_\_.

1. optical centre
2. focus
3. centre of curvature
4. pole


38.Convex lens always gives a real image if the object is situated beyond \_\_\_.

1. optical centre
2. centre of curvature
3. Focus
4. radius of curvature

39. Parallel rays of light entering a convex lens always converge at \_\_\_.

1. centre of curvature
2. the principal focus
3. optical centre
4. the focal plane

40. Where should an object be placed so that a real and inverted image of the same size is obtained, using a convex lens?



1. Between O and F
2. At F
3. At 2F
4. At infinity

41. SI unit of the power of a lens is \_\_\_.

1. diopetre
2. cm
3. metre
4. watt

42. 1 D is the power of the lens of focal length of \_\_\_ cm.

1. 100
2. 10
3. 1/100
4. 1/10

43. In a simple microscope lens used is \_\_\_.


1. biconvex
2. biconcave
3. plano convex
4. cylindrical

44. Reciprocal of focal length in metres is known as the \_\_\_ of a lens.

1. focus
2. power
3. power of accommodation
4. far point

45. A convex lens is called \_\_\_.

1. converging lens
2. diverging lens
3. both converging and diverging lens
4. refracting lens



46. A positive magnification greater than unity indicates \_\_\_.

1. real image
2. virtual image
3. neither real nor virtual image
4. distorted image

47. The power of a convex lens of focal length 50 cm is \_\_\_.

1. +20 D
2. +2 D
3. +50 D
4. -20 D

48. The focal length of a lens whose power is -1.5 D is \_\_\_.

1. -66.66 cm
2. +1.5 m
3. +66.66 cm
4. -1.5 m

49. Real images formed by single convex lenses are always \_\_\_.


1. on the same side of the lens as the object
2. inverted
3. erect
4. smaller than the object

50. An object is placed 12 cm from a convex lens whose focal length is 10 cm. The image must be

1. virtual and enlarged
2. virtual and reduced in size
3. real and reduced in size
4. real and enlarged

51. When a person uses a convex lens as a simple magnifying glass, the object must be placed at a distance \_\_\_.

1. less than one focal length
2. more than one focal length



3. less than twice the focal length
  4. more than twice the focal length
52. The image produced by a concave lens is \_\_\_.

1. always virtual and enlarged
2. always virtual and reduced in size
3. always real
4. sometimes real, sometimes virtual

53. A virtual image is formed by \_\_\_.

1. a slide projector in a cinema hall
2. the ordinary camera
3. a simple microscope
4. Telescope

54. An object is placed 25 cm from a convex lens whose focal length is 10 cm. The image distance is \_\_\_ cm.


1. 50 cm
2. 16.66 cm
3. 6.66 cm
4. 10 cm

55. The least distance of distinct vision is \_\_\_

1. 25 cm
2. 25 m
3. 0.25 cm
4. 10 cm

56. A convex lens has a focal length of 20 cm. Its power in dioptries is \_\_\_

1. 2
2. 5
3. 0.5
4. 0.2



57. An object is placed before a concave lens. The image formed \_\_\_

1. is always erect
2. may be erect or inverted
3. is always inverted
4. is always real

58. A ray of light travels from a medium of refractive index  $n_1$  to a medium of refractive index  $n_2$ . If angle of incidence is  $i$  and the angle of refraction is  $r$ , then  $\frac{\sin i}{\sin r}$  is equal to

1.  $n_1$
2.  $n_2$
3.  $n_2 n_1$
4.  $n_1 n_2$

59. Two thin lenses of power  $+5\text{ D}$  and  $-2\text{ D}$  are placed in contact with each other. Focal length of the combination is \_\_\_


1.  $+3\text{ m}$
2.  $-3\text{ m}$
3.  $0.33\text{ m}$
4.  $-0.33\text{ m}$

60. An object is placed between two plane mirrors inclined at an angle of  $45^\circ$  to each other. How many images do you expect to see?

How can an inverted image of an object be obtained with a plane mirror?

62. How many images of himself can an observer see in a room whose ceiling and two adjacent walls are mirrors?

63. A convex mirror is held in water. What should be the change in its focal length?



64. If the light from the Sun subtend an angle  $Q$  at the pole of a concave mirror of focal length  $f$ , where will it converge? Also find the diameter of the image. Draw the necessary ray diagram.
65. When does Snell's law of refraction fail?
66. How does the frequency of a beam of ultraviolet light changes when it goes from air to glass?
67. State two conditions under which a ray of light suffering refraction from medium 1 to medium 2 does not undergo any change in direction.
68. Name the factor on which lateral shift produced by glass slab depends?
69. A glass lens is immersed in water. How is the power of the lens affected?
70. Why is the power of a lens is measured as reciprocal of focal length? Give reason.
71. A convex lens is made of a material of refractive index  $m_1$  when placed in a medium of refractive index  $m_2$  , behaves as a diverging lens. How are  $m_1$  and  $m_2$  related?
73. An object is held at a distance of 60 cm from a convex mirror of focal length 20 cm. At what distance from the mirror, should a plane mirror be held so that images in the two mirrors coincide? (ans. 22.5 cm)
74. How do you find the rough focal length of a convex lens? Is the same method applicable to a concave lens?
75. Which factors determines the focal length of a lens?

