



PATIENT INSTRUCTIONS

Colonoscopy & Upper Endoscopy (BiPegLyte Preparation)

Patient Name: _____

Physician: _____

Appointment Date: _____

Arrival Time: _____

LOCATION:

**London Digestive Disease Institute
785 Wonderland Road South, Suite 253A (Second Floor)
London, ON N6K 1M6**

If you have questions regarding your procedure or cannot attend, please contact the office immediately. Wait lists for this procedure are very long and it is important to give at least one week's notice to cancel or reschedule so that your appointment may be offered to another patient.

ENDOSCOPIC PROCEDURE - IMPORTANT INSTRUCTIONS

- Propofol sedation is used for endoscopic procedures.
 - **You must have an adult (family member or friend) who will stay in the waiting room throughout the procedure and drive you home afterwards. You cannot drive yourself home.**
- If you choose not to receive sedation, you may go home alone, but the procedure may be uncomfortable and incomplete.
- Do not drive, operate machinery, or drink alcohol for 12 hours after your procedure.

Please bring the following items to your appointment:

- Your health card
- A current list of all medications you take and allergies
- A copy of your completed LDDI Pre-Endoscopy Anesthesia Patient Questionnaire
- A covered water bottle
- A reusable bag for your belongings

Please avoid wearing jewelry, perfume, or cologne.

MEDICATIONS & MEDICAL CONDITIONS

- Stop iron supplements 7 days before the procedure.
- If you take blood thinners (other than aspirin), contact the office immediately for specific instructions.

- You may be instructed to alter your medication several days prior to your procedure.
 - Examples: Plavix (clopidogrel), Ticlid (ticlopidine), Brilinta (ticagrelor), Coumadin (warfarin), Pradaxa (dabigatran), Eliquis (apixaban), Xarelto (rivaroxaban).
- If you have diabetes and take insulin or other medications, contact the office immediately for specific instructions. You may be instructed to alter your medication several days prior to your procedure. Please ensure you bring your insulin or medication on the day of your procedure.
- Stop GLP-1 medications 14 days before your appointment.
 - Examples: Ozempic (semaglutide), Saxenda (liraglutide)
- Please notify our office if you have kidney disease or congestive heart failure. You may be instructed to use an alternative bowel cleansing solution.
- Continue all other medications as usual, including the day of the procedure.

HOW TO PREPARE

The following medication can be purchased over the counter at your pharmacy:

One (1) Package of Bi-PegLyte.

One Week Prior To Your Procedures:

- Stop eating nuts, seeds (poppy seeds, flax seeds and sesame seeds), corn, popcorn, fruits containing seeds (kiwi, raspberries, strawberries, tomatoes).

One Day Before Your Procedures:

- Consume clear fluids only all day. NO SOLID FOOD.
 - Examples: water, apple juice, sports drinks, clear broth, popsicles, Jell-O, black coffee or tea (no milk/cream). *Please avoid RED, BLUE, PURPLE coloured drinks.*
- 3:00 PM:
 - Take 3 Bisacodyl tablets (swallow whole; do not chew or crush).
- 7:00 PM:
 - Mix 1 packet of PEG 3350 with 1L of water. Chill for better taste.
 - Drink the full liter by 8:00 PM.
 - Continue drinking clear fluids.

Procedure Day:

- NO SOLID FOOD.
- 6 Hours Before Your Arrival Time:
 - Mix the second PEG 3350 packet with 1L water.
 - Drink the entire liter within 1 hour.
 - Continue drinking clear fluids after this but you must stop all fluids 2 hours before your appointment time.
 - Do not chew gum or suck on candy.



UNDERSTANDING YOUR PROCEDURES

What is a Colonoscopy and Upper Endoscopy?

These tests use long, thin, flexible tubes with cameras to examine:

- Your colon and rectum (via the rectum)
- Your upper digestive tract (via the mouth)

They help investigate symptoms such as:

- Heartburn, trouble swallowing, vomiting, bleeding, diarrhea or constipation, abdominal pain and screening for colon polyps and cancer.

During the procedures, the doctor may:

- Take a biopsy (a small tissue sample)
- Remove polyps (abnormal growths that can become cancerous)
 - You will not feel the biopsy or polyp removal.

What to Expect During and After Your Procedures

- You will receive propofol anesthesia for your sedative. This will be administered by an Anesthesiologist.
- A spray may be given to the back of your throat to make it feel numb.
- There may be some discomfort during the procedures, however, it is usually well-tolerated.
- You may feel bloating or need to pass gas/burp due to air inflated during the procedure.
- You will stay in the recovery room until the anesthesia wears off.
- Afterward, you can eat and drink as usual unless specified by your physician.
- Avoid alcohol for at least 12 hours.

RISKS AND COMPLICATIONS

These procedures are generally very safe. However, potential complications include:

- Sedation-related issues: low blood pressure, low oxygen, or difficulty waking.
- Bleeding from biopsy or polyp sites — usually minor.
- Perforation (a tear in the digestive tract wall):
 - Occurs in about 1 in 5000 endoscopies and 1 in 1000 colonoscopies.
 - May require hospitalization, IV fluids, blood transfusion, antibiotics, surgery, or an ostomy bag.

Limitations: These tests are not perfect — in rare cases; polyps, lesions or other serious problems in the colon may not be seen and can be missed.

***If you experience any problems after your procedure, please contact the office immediately
(519) 204-6333.***

