

High precision optical polarimetry of black hole X-ray binaries

Vadim Kravtsov, University of Turku

Andrei Berdyugin

Alexandra Veledina

Juri Poutanen

Astrophysical Polarimetry in the Time-Domain Era


Politecnico di Milano

Lecco, Italy

29 August -1 September 2022


Motivation

Geometry and spectral states of low-mass black hole X-ray binaries (BHXRBs)



[Image produced with BinSim by Rob Hynes]

Spectrum of BHXRb – product of a complex interplay between contribution of several components.





The identification of the different spectral components is essential for understanding the mechanisms that trigger the outbursts.

Motivation

Choice of a tool

There are **several methods** that can be used to identify the **contribution of different components**: *photometry, spectroscopy, imaging, timing, and **polarimetry***.

Why **polarimetry**?





Motivation

Choice of a tool

There are **several methods** that can be used to identify the **contribution of different components**: *photometry, spectroscopy, imaging, timing, and **polarimetry***.

Why **polarimetry**?

- **Polarized radiation can be produced by several** physical processes taking place in the BHXRBs, such as **synchrotron radiation** in the presence of an ordered magnetic field of the jet (hot accretion flow) or **scattering** of the accretion disk radiation **by electrons or dust**.





Motivation

Choice of a tool

There are **several methods** that can be used to identify the **contribution of different components**: *photometry, spectroscopy, imaging, timing, and **polarimetry***.

Why **polarimetry**?

- **Polarized radiation can be produced by several** physical processes taking place in the BHXRBs, such as **synchrotron radiation** in the presence of an ordered magnetic field of the jet (hot accretion flow) or **scattering** of the accretion disk radiation **by electrons or dust**.
- **Polarization carries information about the geometrical properties** of the emitting/scattering media.





Motivation

Choice of a tool

There are **several methods** that can be used to identify the **contribution of different components**: *photometry, spectroscopy, imaging, timing, and polarimetry*.

Why **polarimetry**?

- **Polarized radiation can be produced by several** physical processes taking place in the BHXRBs, such as **synchrotron radiation** in the presence of an ordered magnetic field of the jet (hot accretion flow) or **scattering** of the accretion disk radiation **by electrons or dust**.
- **Polarization carries information about the geometrical properties** of the emitting/scattering media.
- **Each component has different polarization properties** (or no polarization at all).



BHXRBs: previous polarimetric studies


Quick review of recent results

Polarization in the **outburst**:


- Shahbaz et al., 2016 – **V404 Cyg**, optical and NIR polarimetry, $P_{\text{int}} \sim 4\%$, jet origin
- Tanaka et al., 2016 – **V404 Cyg**, optical and NIR polarimetry, no intrinsic polarization
- Kosenkov et al., 2017 – **V404 Cyg**, optical polarimetry, $P_{\text{int}} \sim 1\%$, scattering origin
- Veledina et al., 2019 – optical polarimetry of **MAXI J1820+070**, $P_{\text{int}} \sim 0.7\%$, scattering origin

Polarization in **quiescence**:

- Dolan & Tapia, 1989; Dubus et al., 2008, **1A 0620–00**, optical polarimetry, $P_{\text{int}} \sim 3\%$, scattering origin
- Russel et al., 2016 – **1A 0620–00**, **Swift J1357.2**, NIR polarimetry, $P_{\text{int}} \sim 1\text{--}8\%$, jet origin
- Poutanen et al., 2022 – **MAXI J1820+070**, optical polarimetry, blue polarization up to $P_{\text{int}} \sim 5\%$, non-synchrotron origin, jet-spin misalignment(!)





Poutanen et al., 2022



Kosenkov et al., 2020

Our sample of BHXRBs

Galactic distribution and main parameters



Object	Companion	V mag	i deg	P_{orb}
XTE J1118+480	K7 V – M1 V	19.6 ± 0.2^1	68 ± 2	4.07841(5) h
Swift J1357.2–0933	M5 V	17.27 ± 0.02^1	> 70	2.8 ± 0.3 h
4U 1957+115	...	$\approx 19.0^3$	20 – 70	9.33 ± 0.01 h
V404 Cyg	K3 III	$\approx 18.7^2$	67 ± 3	6.473 ± 0.001 d
V4641 Sgr	B9 III	$\approx 13.5^2$	72 ± 4	2.817 ± 0.001 d
XTE J2012+381	-	21.3 ± 0.1^4
MAXI J1820+070	K4 V	-	73 ± 8	16.87 ± 0.07 h
MAXI J0637–430	...	$\approx 16.5^2$

Our sample of 8 BHXRBs

Observations


The telescopes and instruments we used

Nordic Optical Telescope (NOT)



La Palma, Canary Islands, Spain

DIPoI-UF polarimeter, mounted on NOT



Simultaneous Three-color (*BVR*) polarimeter

Why DIPoI-UF/2?

- Precision up to 10^{-5} ($\Delta P \sim 0.001\%$)
- **Simultaneous** three-band (*BVR*) polarimetry
- **Sky polarization is optically eliminated!**

Pirola et al., 2020

Observations

Detailed description

BHXRB	State	Observing nights
XTE J1118+480	Q	1 (NOT)
Swift J1357.2–0933	RH*	1 (NOT)
4U 1957+115	S	1 (NOT)
V4641 Sgr	RH*, Q	11 (5 – NOT, 6 – T60)
V404 Cyg	RH, Q	2 (NOT)
MAXI J1820+070	RH*, S, DH, Q	10 (T60)
MAXI J0637–430	S	3 (T60)

States: Q – quiescence, RH – rising hard, S – soft, DH – decaying hard

*Failed outburst


$$q_{\text{obs}} = q_{\text{int}} + q_{\text{is}} \Rightarrow \begin{cases} q_{\text{int}} = q_{\text{obs}} - q_{\text{is}} \\ u_{\text{int}} = u_{\text{obs}} - u_{\text{is}} \end{cases}$$

$$P_{\text{int}} = \sqrt{q_{\text{int}}^2 + u_{\text{int}}^2} \quad \theta_{\text{int}} = \frac{1}{2} \arctan u_{\text{int}}/q_{\text{int}}$$

When $P/\sigma < 5$, the polarization degree is biased towards higher values!

$$P_0 = \sqrt{P^2 - 2\sigma^2}, \text{ Simmons \& Stewart (1985)}$$

unbiased maximum-likelihood estimator



Results: Swift J1357.2–0933, 4U 1957+115 and XTE J1118+480

Observed and intrinsic polarization




Figure 1. Normalized observed Stokes parameters (q , u) for Swift J1357.2–0933, 4U 1957+115, and XTE J1118+480 (from left to the right). The blue, green, and red squares with 1σ errors correspond to the B , V , and R optical polarimetric measurements of the targets and the circles correspond to nearby stars.

Results: V4641 Sgr


Observed and intrinsic polarization

We measured the polarization of **six** nearby field stars with similar *Gaia* parallaxes.


The observed polarization of **V4641 Sgr** is consistent with the polarization pattern of nearby stars and shows **no dependence on orbital phase**.

From these two facts we can conclude, that the **polarization of V4641 Sgr has an interstellar origin**.

Polarization vs *Gaia* parallax



Polarization vs Orbital phase



Results: MAXI J1820+070, V404 Cyg and MAXI J0637–430


Observed and intrinsic polarization

MAXI J1820+070 shows complex behaviour of intrinsic polarization:

- $P_{\text{int}} \sim 0.5\%$ with the red spectrum in the **rising hard state**
- $P_{\text{int}} \sim 0.1\%$ in the **soft state**
- No polarization in **decaying hard state**
- **Substantial** (up to $P_{\text{int}} \sim 5\%$) **intrinsic polarization** with the blue spectrum in the **quiescent state** (Poutanen et al., 2022)

V404 Cyg shows small ($P_{\text{int}} < 0.5\%$) intrinsic polarization during the **quiescent state**, consistent with the previous measurements

Unbiased maximum-likelihood estimation of **soft state** polarization of **MAXI J0637–430** is **zero** with 3σ upper limit of $P_{\text{int}} < 1.1\%$



Summary

- Most of the sources show **no evidences** of significant intrinsic polarization
- The absence of intrinsic polarization at the optical wavelengths **puts constraints** on the potential contribution of non-stellar (jet, hot flow, accretion disc) components to the total spectra of black hole X-ray binaries
- High precision optical polarimetry can shed new light on the radiative mechanisms that occur in BHXRBS.

Intrinsic polarization estimates of observed sample

Source	State	P_B per cent	P_V per cent	P_R per cent
XTE J1118+480	Q	1.2 ± 0.8	1.4 ± 0.8	1.3 ± 0.5
Swift J1357.2–0933	RH ^a	≤ 0.5	≤ 0.4	≤ 0.4
4U 1957+115	S	≤ 0.2	≤ 0.3	≤ 0.3
V4641 Sgr	RH ^a	≤ 0.1	≤ 0.1	≤ 0.1
	Q	≤ 0.1	≤ 0.1	≤ 0.1
V404 Cyg	RH	0.8 ± 0.3	1.1 ± 0.1	0.5 ± 0.1
	Q	≤ 0.5	≤ 0.5	≤ 0.5
MAXI J1820+070	RH1	0.28 ± 0.01	0.36 ± 0.01	0.30 ± 0.01
	RH2	0.34 ± 0.02	0.51 ± 0.02	0.53 ± 0.02
	S	0.16 ± 0.01	0.15 ± 0.01	0.02 ± 0.01
	DH	0.06 ± 0.04	0.13 ± 0.06	0.09 ± 0.04
	RH ^a	≤ 0.3	≤ 0.4	≤ 0.3
	Q	3.2 ± 0.2	1.9 ± 0.2	0.9 ± 0.1
MAXI J0637–430	S	≤ 0.2	–	–

Notes. ^aFailed outburst. States: Q – quiescence, RH – rising hard, S – soft, DH – decaying hard

Thank you for your attention!

