

ALGEBRA

SYNOPSIS - 1

ALGEBRAIC EXPRESSIONS

FUNDAMENTAL CONCEPTS

There are two types of symbols in algebra:

(i) Constants: A symbol having a fixed numerical value is called a constant.

Thus, each of the symbols $3, -2, \frac{5}{9}, 0.6$ is a constant.

In fact, every number is a constant.

(ii) Variables: Consider the following example.

Example: We know that the perimeter of a square is given by the formula,

$$p = 4 \times s$$

Here 4 is a constant.

When $s = 3$, then $p = 4 \times s$

When $s = 5$, then $p = 4 \times 5 = 20$

Thus, the values of p and s are not fixed, they vary.

A symbol which takes on various numerical values is called a variable or a literal.

In the above example, s and p are literals or variables.

OPERATIONS ON NUMBERS AND LITERALS

We shall denote the literals by a, b, c, x, y, z, m, n etc.

1. Addition:

(i) The sum of x and 8 is $x + 8$ (ii) The sum of x and y is $x + y$

(iii) For any literals, a, b, c we have

$$\text{I. } a + b = b + a$$

$$\text{II. } a + 0 = 0 + a$$

$$\text{III. } (a + b) + c = a + (b + c)$$

2. Subtraction:

(i) 6 less than x is $x - 6$

(ii) y less than x is $x - y$.

3. Multiplication:

(i) 7 times x is $7 \times x$, written as $7x$ (ii) The product of x and y is xy

(iii) For any literals, a, b, c we have

$$\text{I. } ab = ba$$

$$\text{II. } a \times 0 = 0 \times a = 0$$

$$\text{III. } (ab)c = a(bc)$$

$$\text{IV. } a(b + c) = ab + ac$$

4. Division:

(i) 5 divided by x is written as $\frac{5}{x}$ (ii) x divided by 6 is written as $\frac{x}{6}$

(iii) x divided by y is written as $\frac{x}{y}$

Algebraic Expression:

A combination of constants and variables, connected by the symbols $+$, $-$, $\times$ and $\div$ is called an algebraic expression.

Example: $5x + 6y + 2xy$ is an algebraic expression.

Terms: The several parts of an expression separated by the sign $+$ or $-$ are called the terms of the expression.

Examples:

(i) The expression $2x - 5y + 3xyz$ has three terms, namely $+2x$, $-5y$ and $+3xyz$

(ii) The expression $a^2b - 2ab^3 + 5b^2a - 8$ has four terms, namely $+a^2b - 2ab^3$, $+5b^2a$ and -8

(iii) The expression abc has only one term.

Various Types of Algebraic Expressions

- i. Monomials: An algebraic expression which contains only one term is called a monomial.

Examples: Each one of the expressions $5x, 9xy, 8a^2bc, -2z^3, 6, \frac{5}{x}$ is a monomial.

- ii. Binomials: An algebraic expression which contains two terms is called a binomial.

Examples: Each one of the expressions $5 + 2x, 1 - 3xyz, x^2 + a^2, a + \frac{1}{a}$ is a binomial.

- iii. Trinomials: An algebraic expression which contains three terms is called a trinomial.

Examples: Each one of the expressions $2x + 3y - 4z, 5 - a - bc, a^2 + b^2 - 2ab$ is a trinomial.

- iv. Multinomials: An algebraic expression containing two or more terms is called a multinomial.

Factors of a Term:

When numbers and literals are multiplied to form a product, then each quantity multiplied is called a factor of the product.

A constant factor is called a numerical factor while a variable factor is called a literal factor.

Examples: (i) In $7ab$, the numerical factor is 7 and the literal factors are a, b and ab .

(ii) In $-9x^2y$, the numerical factor is -9 and the literal factors are x, x^2, y, xy and x^2y .

Constant Term:

A term of the expression having no literal factor is called the constant term.

Examples: (i) In the expression $3x - 4y + 2$, the constant term is 2.

(ii) In the expression $a^2 + b^2 - 3ab - 5$ the constant term is -5

Coefficients:

Any of the factors of a term is called the coefficient of the product of other factors. In particular, the constant part is called the numerical coefficient of the term and the remaining part is called the literal coefficient of the term.

Examples:

In the term $-2xyz^2$:

Numerical coefficient = -2 ;

Literal coefficient = xyz^2 ;

Coefficient of $x = -2yz^2$;

Coefficient of $y = -2xz^2$.

Coefficient of $z^2 = -2xy$ etc.

Like Terms:

Two terms having same literal factors are known as like terms. Otherwise, they are known as unlike terms.

Examples: (i) $3xy, -5xy$ are like terms.

(ii) $2a^2b, 3ab^2$ are unlike terms.

(iii) $6ab^2, 9b^2a$ are like terms [Since $ab^2 = b^2a$].

Polynomials:

An algebraic expression in which the powers of the variables involved are non-negative integers, is called a polynomial.

The highest power of the variable in a polynomial is called its degree.

Examples:

(i) $3x + 7$ is a polynomial in x of degree 1.

(ii) $2y^2 - 5y + 1$ is a polynomial in y of degree 2.

(iii) $z^3 + 4z^2 + 3z - 6$ is a polynomial in z of degree 3.

(iv) $x + \frac{1}{x}$ is not a polynomial, since $\frac{1}{x} = x^{-1}$, i.e., power of x is negative integer.

Note that $x + \frac{1}{x}$ is a binomial expression but it is not a polynomial.

Polynomials in Two or More Variables:

An algebraic expression involving two or more variables with non-negative integral power is called a polynomial in these variables.

The degree of any term of a polynomial is the sum of the powers of all the variables in that term.

The degree of the highest degree term in a polynomial is called the degree of the polynomial.

Examples:

- i. $x + y + xy$ is a polynomial in x and y whose terms are of degree 1, 1 and 2 respectively. So, it is a polynomial of degree 2.
- ii. $a^2b + ab^2 + 3ab + 5$ is a polynomial in a and b whose terms are of degree 3, 3, 2 and 0 respectively. So, it is a polynomial of degree 3.
- iii. $y + z$ is a polynomial in y and z whose terms are of degree 1 and 1 respectively. So, it is a polynomial of degree 1.

Zero Value: The values which are satisfying the given polynomial.

Example: $f(x) = x + 3$. -3 is zero value of $f(x)$.

WORK SHEET - 1

SINGLE ANSWER TYPE

1. The value of the expression $\frac{n^2}{2} + \frac{n}{2}$ when $n = 12$ is
 - 1) 76
 - 2) 74
 - 3) 78
 - 4) 72
2. If $\frac{7x}{3} - \frac{7}{6}$ is a polynomial, then the zero of the polynomial is
 - 1) $\frac{1}{2}$
 - 2) $-\frac{1}{2}$
 - 3) 0
 - 4) -2
3. If the zero of the polynomial in 'x' is $-\frac{5}{4}$, then the polynomial is
 - 1) $4x - 5$
 - 2) $5x - 4$
 - 3) $5x + 4$
 - 4) $4x + 5$
4. If $A = -8x^2 - 6x + 10$, then its value when 'x' = $\frac{1}{2}$ is
 - 1) 6
 - 2) 4
 - 3) 5
 - 4) 7
5. The third degree polynomial among the following is
 - 1) $2x^{3-1} + 3x^{2-1} + 5$
 - 2) $3x^{4-1} + 2x^{3-1} + 6x^{2-1} + 8$
 - 3) $3x^{-2-1} + 4x^{-2} + 5$
 - 4) $2x^{5-3} + 3x^{4-3} + 7$
6. Among the following the expression which is not a monomial is
 - 1) $\frac{4a^3b^2c^5}{23}$
 - 2) $-147x^3y^2$
 - 3) $\frac{2}{7}x^{-2}y^5z$
 - 4) $x^3y^5z^{12}$
7. If $x = \frac{a}{2}$, then the value of $4x^2 + 8x + 18$ is
 - 1) $a^2 + 2a + 8$
 - 2) $a^2 + 3a + 18$
 - 3) $a^2 + 4a + 18$
 - 4) $a^2 + 5a + 18$

8. The value of the expression $\frac{-26}{3} - \frac{13x}{27}$ when $x = \frac{9}{13}$ is
 1) - 8 2) - 10 3) - 9 4) - 11
9. Degree of the polynomial $p + q x^m + r x^{m+2} + 5x^{m+3} + x^{m+4}$ is
 1) m 2) m + 2 3) m + 3 4) m + 4
10. If $\frac{n(n+1)(2n+1)}{6}$ represents sum of the squares of first 'n' natural numbers, then its value when n = 10 is
 1) 365 2) 375 3) 395 4) 385
11. Degree of the polynomial $\frac{1}{2}x^5 + 3x^4 + 2x^3 + 3x^2 + 6$ is
 1) 4 2) 3 3) 5 4) 2
12. Degree of the monomial $\frac{3}{5}x^2y^6z^7$ is
 1) 15 2) 9 3) 8 4) 13
13. In a polynomial $3x + 5$ where $x = a + 2$, then its value when $a = 8$ is
 1) 25 2) 45 3) 35 4) 40

MULTI ANSWER TYPE

14. Which of the following is/are true?
 1) The coefficient of 'x' in $9xy$ is $-9y$ 2) The coefficient of 'a' in $-7abc$ is $-7bc$
 3) The coefficient of 'xyz' in $-xyz$ is -1 4) The coefficient of 'b' in $-abc$ is $-ac$
15. In $6x^2y + 5xy^2 - 8xy^2 - 7yx^2$, like terms are
 1) $6x^2y, -7x^2y$ 2) $5xy^2, -8xy^2$ 3) $6x^2y, -5xy^2$ 4) $-7yx^2, 8xy^2$
16. Which of the following is/are false?
 a) 3 more than a number 'x' is $x + 3$
 b) One third of the sum of x and y is $(x + y)/3$
 c) The quotient of x by y added to the product of x and y is $(x + y) + x/y$
 d) 5 less than the quotient of x by y is $(x/y) + 5$
17. a) The degree of $x^2 + xy^2 + y^3$ is 3 b) The degree of $m^2n^3 + mn^2 + 4$ is 5
 c) The degree of $p^2q^2 + pq^2 + 1$ is 4
 which of the above is/are true?
 1) a 2) b 3) c 4) all of these
18. Which of the following statement is/are false?
 a) $x^3 - \frac{1}{x^3} + 3x^2 - \frac{3}{x^2} + 8$ is a polynomial.
 b) $p^3 + p^2q - \frac{4p}{q} + 8$ is a polynomial.
 c) $a^3 + b^3 + c^3 - abc$ is a polynomial.
 1) a 2) b 3) c 4) both 1 & 2

19. The degree of the polynomial $2x^{3^2 \times 2^3} + 3x^{4^2 \times 3^2} + 5x^{4^2 \times 3^2} + 5x^{4^2 \times 2^3}$ is
 1) 12^2 2) 146 3) 144 4) 148
20. If $A = \frac{1}{2}(m+n)p$ where $m = 12.25$, $n = 108.35$, $p = 3.5$, then the value of A is less than
 1) 211.05 2) 211.50 3) 211.00 4) 211.105
21. If $a = 1$ and $b = \frac{1}{2}$, then the value of $(25a^4b) \times (-2a^2b^2) \times (-2.1a^3b^3)$ is
 1) $131/32$ 2) $-105/32$ 3) $128/17$ 4) $-105/64$
22. If $a = 7$, $b = 5$, then the value of $(a + b)(a + b) - (a - b)(a - b) + (a^2 - b^2)$
 1) 41×2^3 2) 41×2 3) 41×2^2 4) 41×2^4

REASONING ANSWER TYPE

23. *Statement I* : The value of $(a + b)^2 + (a - b)^2 + (a^2 - b^2)$ when $a = 3$, $b = 2$ is 30
Statement II : $(a + b)^2 = a^2 + b^2 + 2ab$, $(a - b)^2 = a^2 + b^2 - 2ab$, $a^2 - b^2 = (a + b)(a - b)$
 1. Both Statements are true, Statement II is the correct explanation of Statement I.
 2. Both Statements are true, Statement II is not correct explanation of Statement I.
 3. Statement I is true, Statement II is false.
 4. Statement I is false, Statement II is true.
24. *Statement I* : If 10 oranges are taken out from a basket containing 'x' oranges and added to another basket containing y + 20 oranges, then the total number of oranges in the two baskets is $x + y + 30$.
Statement II : The sum of 'p' and 10 is subtracted from sum of 5 more than 4 times 'q' and 'r', then the resultant is $(4q + r) - (p + 5)$
 1. Both Statements are true, Statement II is the correct explanation of Statement I.
 2. Both Statements are true, Statement II is not correct explanation of Statement I.
 3. Statement I is true, Statement II is false.
 4. Statement I is false, Statement II is true.
25. *Statement I* : If $A = 4x^{2m}y^{4n}z^{3p}$ where $2m = 4n = 3p = 24$, then the degree of polynomial A is 74.
Statement II : In case of polynomials in more than one variable the sum of the power of the variable in each term is taken and the highest sum is the degree of the polynomial.
 1. Both Statements are true, Statement II is the correct explanation of Statement I.
 2. Both Statements are true, Statement II is not correct explanation of Statement I.
 3. Statement I is true, Statement II is false.
 4. Statement I is false, Statement II is true.

26. *Statement I* : $2m^2n, -4nm^2, \frac{-8}{3}m^2n$ are like terms.

Statement II : Monomials having different literal factors are like terms.

1. Both Statements are true, Statement II is the correct explanation of Statement I.
 2. Both Statements are true, Statement II is not correct explanation of Statement I.
 3. Statement I is true, Statement II is false.
 4. Statement I is false, Statement II is true.
27. *Statement I* : The value of $(5a^6) \times (-10ab^2) \times (-2.1a^2b^3)$ when $a = 1, b = 1/2$ is $105/32$.
Statement II : The value of an expression depends on the given values of the variable.
1. Both Statements are true, Statement II is the correct explanation of Statement I.
 2. Both Statements are true, Statement II is not correct explanation of Statement I.
 3. Statement I is true, Statement II is false.
 4. Statement I is false, Statement II is true.
28. *Statement I* : If $x = 2, y = -2$ and $z = 3$, then $x^2 + y^2 + z^2 - 2xy - 2yz - 2zx = 25$

Statement II : If $p = 3, q = -5$ and $r = 4$, the $p^2 + q^2 + r^2 - pq - qr - pr = 73$

1. Both Statements are true, Statement II is the correct explanation of Statement I.
2. Both Statements are true, Statement II is not correct explanation of Statement I.
3. Statement I is true, Statement II is false.
4. Statement I is false, Statement II is true.

COMPREHENSION TYPE

Writeup-1:

If $A = 18x^3y^2z^4, B = 12x^2y^3z^3, C = x^4y^3$

29. a) $A \times C$ in simplified form is $18x^7y^7z^7$

b) $A \times C$ in simplified form is $18x^6y^7z^3$

Which of the following is/are true?

- | | | | |
|------|------|-------------------|---------|
| 1) a | 2) b | 3) both (1) & (2) | 4) none |
|------|------|-------------------|---------|
30. $B \times C$ in product form is
- | | | | |
|------------------|-------------------|------------------|------------------|
| 1) $12x^6y^6z^3$ | 2) $-12x^6y^6z^3$ | 3) $12x^5y^5z^3$ | 4) $12x^6y^5z^3$ |
|------------------|-------------------|------------------|------------------|
31. $A \times B \times C$ in product form is
- | | | | |
|-------------------|-------------------|-------------------|-------------------|
| 1) $216x^9y^8z^7$ | 2) $216x^8y^9z^7$ | 3) $216x^7y^8z^7$ | 4) $216x^9y^8z^6$ |
|-------------------|-------------------|-------------------|-------------------|

Writeup-2:

$A = 6a^{p+2} + 5a^{2p+3} + 7a^{3p+4}$ and $B = 4b^{q+3} + 5b^{3q+2} + 8b^{4q+3}$ where $p = 2, q = 3$

32. The degree of polynomial A is
- | | | | |
|-------|------|------|-------|
| 1) 10 | 2) 7 | 3) 6 | 4) 11 |
|-------|------|------|-------|
33. The degree of polynomial B is
- | | | | |
|-------|-------|-------|-------|
| 1) 10 | 2) 15 | 3) 11 | 4) 12 |
|-------|-------|-------|-------|
34. Which polynomial has higher degree?
- | | | | |
|------|------|---------------|---------|
| 1) A | 2) B | 3) both 1 & 2 | 4) none |
|------|------|---------------|---------|

Writeup-3:

(i) An algebraic expression of the form $a + bx + cx^2 + dx^3 + \dots$ where a, b, c, d etc are constants is a polynomial in one variable. Here the highest power of x is the degree of the polynomial.

(ii) An algebraic expression in two or more variables is a polynomial if every variable is a polynomial if every variable in it has only positive integral powers. Here the sum of the variables in each term is taken and the highest sum is the degree of the polynomial.

35. Which of the following is a polynomial in one variable ?

- 1) $x^3 - \frac{1}{x^3}$ 2) $x^{-3} + 2$ 3) $\frac{y^3}{x}$ 4) $\frac{4}{7}t - t$

36. The degree of the Polynomial $\frac{3}{5}x^2 - \frac{5}{4}x + \frac{2}{3}$ is _____.

- 1) $3/5$ 2) 2 3) 3 4) Not a Polynomial

37. The degree of the polynomial $2p^2q + pq^2 + 3p^3 - p^2q^2$ is _____.

- 1) 2 2) 3 3) 4 4) Not a Polynomial

Writeup-4:

If $A = \frac{V}{W(R^2 - r^2)}$ where $V = 115.5$, $W = 22/7$, $R = 2.6$, $r = 2.3$, then

38. The value of $R^2 - r^2$ is

- 1) 1.48 2) 1.44 3) 1.46 4) 1.47

39. The value of V/W is

- 1) $\frac{143}{4}$ 2) $\frac{149}{4}$ 3) $\frac{141}{4}$ 4) $\frac{147}{4}$

40. The value of A is

- 1) 2.5 2) 25 3) 0.25 4) None

Writeup-5:

The value of the expression $ax^2 + bx + c$ at $x = k$ is $(ak^2 + bk + c)$

41. The value of the expression $2x^2 + (7/2)xy + 5y^2$ at $x=1$, $y = 2$ is

- 1) 92 2) 29 3) -29 4) 19

42. The value of the expression $a^2 - bc + c^2 - b^2$ at $a = 0$, $b = 1$ $c = 2$ is ____

- 1) 1 2) 2 3) 3 4) 4

43. The value of the expression $x^3 + \frac{xy}{3} + \frac{7}{2}y^2$ at $x = 2$, $y = 3$ is ____

- 1) 75 2) $72/5$ 3) $75/2$ 4) $57/2$

MATRIX MATCHING TYPE

44. **Column-I**

- a) $a.25 = 25.a$
b) $3 \times (5 + x) = 3 \times 5 + 3 \times x$
c) $4 + (5 + x) = (4 + 5) + x$
d) $5 \times (7 \times x) = (5 \times 7) \times x$

Column-II

- 1) associative law in multiplication
2) associative law in addition
3) distributive law
4) commutative law in multiplication

45. **Column-I**

- a) 7 times x increased by 5 gives Quotient of 11 by 2
b) 7 less than the quotient of x by 11 equals 2

Column-II

- 1) $2x - 9 = 23$
2) $\frac{x}{11} - 7 = 2$

c) Twice x decreased by 9 gives 23

$$3) 2 + x - 9 = 23$$

d) 2 times x exceeds 9 by 23

$$4) 7x + 5 = \frac{11}{2}$$

$$5) \frac{x}{7} - 11 = 2$$

46. **Column-I**
Polynomials

Column-II
Degree

a) $5x$

1) 5

b) $15x^2 - x + 2$

2) 4

c) $-x^4 + 2x + 1$

3) 2

d) $x^2 - x^5$

4) 1

47. **Column-I**

Column-II

a) If $l = 5$ and $m = 3$, then $2l + 3m =$ _____

1) 1

b) If $x = 3$, then $x^2 - 4x + 4 =$ _____

2) 13^2

c) If $x = 5$, $y = 12$, then $x^2 + y^2 =$ _____

3) 19

d) If $x = 4$, $y = 3$ & $z = -2$, then $xy + yz + zx =$ _____

4) -2

5) 169

INTEGER ANSWER TYPE

48. 6 less than the quotient of x by 3 equals 2. Their $x =$ _____.

49. The degree of the polynomial $p^3q^2 + 2p^2q + pq^2 + 5p^4 + 8q^3$ is _____.

50. If $a = +1$, $b = -2$ and $c = -3$, then the value of $\frac{a^3 + b^3 + c^3 - 3abc}{ab + bc + ca - (a^2 + b^2 + c^2)}$ is _____

SYNOPSIS - 2

Addition of polynomials:

1. $5x$, $-3x$, $4x$, $\frac{7}{6}x$ are like terms.

2. $-\frac{3}{2}y$, $6x$, $-7x^2$, $8x^3$ are unlike terms.

3. Like terms can be added and their sum can be simplified.

E.g. $7x - 9x + 6x = x(7 - 9 + 6) = 4x$.

4. If no two terms are alike in a polynomial, then it is said to be in the simplified standard form.

Eg : $2x^3 - 8x^2 - 6x + 9$

5. If an expression is

a) in ascending order, then its terms are arranged in increasing order of powers in the expression.

b) in descending order, then its terms are arranged in decreasing order of powers in the expression.

6. The sum or difference of two rational polynomials is also a polynomial with rational coefficients.

WORK SHEET - 2**SINGLE ANSWER TYPE**

- The sum of $\frac{3}{4}x^3, \frac{5}{6}x^3, -\frac{2}{3}x^3$ and $\frac{7}{2}x^3$ is
 - $\frac{12}{53}x^3$
 - $-\frac{53}{12}x^3$
 - $\frac{53}{12}x^3$
 - $-\frac{12}{53}x^3$
- The simplified form of $3x^3 - 2x^2 - 8x - 6x^2 + 7x^3 + 9x + 8x^3 - 9x^2 + 6x$ is
 - $-18x^3 - 17x^2 + 7x$
 - $18x^3 - 17x^2 - 7x$
 - $18x^3 + 17x^2 - 7x$
 - $18x^3 - 17x^2 + 7x$
- The ascending order of the polynomials $-3x^3 + 7x^2 - 9x^4 + 6x - 8$ is
 - $-8 + 6x + 7x^2 - 3x^3 + 9x^4$
 - $-8 - 6x - 7x^2 - 3x^3 - 9x^4$
 - $-8 + 6x + 7x^2 - 3x^3 - 9x^4$
 - $8 + 6x + 7x^2 + 3x^3 + 9x^4$
- If $A = -7x - 3x - 5x$ and $B = 9x + 3x + 2x$, then $A + B$ is
 - $2x$
 - $-2x$
 - $-x$
 - $-3x$
- If $\frac{1}{2}x - \frac{1}{3}x = A$ and $\frac{1}{3}x - \frac{1}{4}x = B$, then $A - B$ is
 - $\frac{1}{12}x$
 - $-\frac{1}{12}x$
 - $-2x$
 - 0
- The equivalent expression of $2x^3 - 3x^2 - 8x - 3$ is
 - $3x^3 - 5x^3 + 7x^2 - 5x^2 - 8x + 10x - 4 + 1$
 - $3x^3 - x^3 - 5x^2 + 2x^2 - 9x + x - 7 + 4$
 - $4x^3 - 6x^2 - 3x^3 + 3x^2 + x^2 - 9x + 3x + 6 - 3$
 - $4x^3 - 2x^3 + 3x^2 - 5x^2 - 8x + 6x + 4 - 1$
- The descending order of $4x^2 - 9x^3 + 3x^2 - 9x^4 + 3x^3 - 9x^2 + 6x - 3x + 5 - 3$ is
 - $-9x^4 + 6x^3 - 2x^2 + 3x + 2$
 - $-9x^4 - 6x^3 + 2x^2 - 3x + 2$
 - $-9x^4 - 6x^3 - 2x^2 + 3x + 2$
 - $-9x^4 + 6x^3 - 2x^2 + 3x - 2$
- If $-\frac{7}{5}x^3 + \frac{3}{4}x^3 + \frac{7}{2}x^3 + \frac{9}{3}x^3$ is added to $\frac{9x^3}{60}$, then the result is
 - $-6x^3$
 - $6x^3$
 - $60x^3$
 - $16x^3$
- If $2x - 3x + 5x = P$, $Q = -8x + 3x + 9x$ and $R = -8x - 6x - 7x$, then $(P + Q) - R$ is
 - $27x$
 - $28x$
 - $29x$
 - $26x$
- If $A = -3x^3 - 2x^3 + 4x^2 - 2x^2$, $B = -3x^2 + 5x^2 - 8x + 3x$ and $C = 2x - 9x - 7 + 8$, then $A + B + C$ in simplified form is
 - $-5x^3 + 4x^2 - 12x + 1$
 - $5x^3 - 3x^2 - 12x + 1$
 - $-5x^3 - 4x^2 - 12x - 1$
 - $5x^3 + 3x^2 + 12x + 1$
- If $4x^3y^2 + 3x^2y^3 - 8x^2y^5$ is added $-9x^2y^3 + 6x^2y^5 - 9x^3y^4$, then the result is
 - $4x^3y^2 + 5x^2y^3 - 2x^2y^5 - 9x^3y^4$
 - $4x^3y^2 - 6x^2y^3 - 2x^2y^5 - 9x^3y^4$
 - $4x^3y^2 - 6x^2y^3 + 2x^2y^5 - 9x^3y^4$
 - $-4x^2y^2 - 6x^2y^3 - 2y^2y^5 - 9x^3y^4$

12. If $0.5x^3 + 1.85x^3 + 2.96x^3 - 4.71x^3$ is added to $(1.25x^4 - 2.5x^5 + 3.6x^4 - 4.71x^4)$, then the result is
- | | |
|-----------------------|------------------------|
| 1) $0.6x^3 + 2.36x^4$ | 2) $-0.6x^3 - 2.36x^4$ |
| 3) $0.6x^3 - 2.36x^4$ | 4) $-0.6x^3 + 2.36x^4$ |

MULTI ANSWER TYPE

13. The sum of $5x^2 - \frac{1}{3}x + \frac{5}{2}$; $-\frac{1}{2}x^2 + \frac{1}{2}x - \frac{1}{3}$ and $-2x^2 + \frac{1}{5}x - \frac{1}{6}$
- 1) $\frac{5}{4}x^2 + \frac{11}{30}x + 2$ 2) $\frac{5}{4}x^2 - \frac{11}{30}x + 2$ 3) $\frac{5}{2}x^2 + \frac{11}{30}x + 2$ 4) $\frac{5}{2}x^2 + \frac{11}{30}x - 2$
14. If the lengths of the three sides of a triangle in centimetres are $\frac{7}{2}x^3 - \frac{1}{2}x^2 + \frac{5}{3}$, $\frac{3}{2}x^3 + \frac{7}{4}x^2 - x + \frac{1}{3}$ and $\frac{3}{2}x^2 - \frac{5}{2}x - 2$, then its perimeter is
- 1) $5x^3 - \frac{11}{4}x^2 + \frac{7}{2}x$ 2) $5x^3 - \frac{11}{4}x + \frac{7}{2}x^2$
- 3) $5x^3 + \frac{11}{4}x^2 - \frac{7}{2}x$ 4) $5x^3 + \frac{7}{2}x^2 - \frac{11}{4}x$

REASONING ANSWER TYPE

15. *Statement I* : If $A = 5x^2 - \frac{1}{3}x + \frac{5}{2}$, $B = -\frac{1}{2}x^2 + \frac{1}{2}x - \frac{1}{3}$, $C = -2x^2 + \frac{1}{5}x - \frac{1}{6}$, then

$$A + B + C = \frac{5}{2}x^2 - \frac{11}{30}x - 2$$

Statement II : Two or more algebraic expression can be added by arranging their terms and combining like terms.

1. Both Statements are true, Statement II is the correct explanation of Statement I.
 2. Both Statements are true, Statement II is not correct explanation of Statement I.
 3. Statement I is true, Statement II is false.
 4. Statement I is false, Statement II is true.
16. *Statement I* : The sum of $8p^2 - 9q^2$ and $-6p^2 + 5q^2$ is $2p^2 - 4q^2$
Statement II : The sum can be found by adding the dissimilar terms in both the expressions.
1. Both Statements are true, Statement II is the correct explanation of Statement I.
 2. Both Statements are true, Statement II is not correct explanation of Statement I.
 3. Statement I is true, Statement II is false.
 4. Statement I is false, Statement II is true.

COMPREHENSION TYPE

Writeup-1:

$$A = 7x^2 - 4x + 5, B = -3x^2 + 2x - 1, C = 5x^2 - x + 9$$

17. The value of $A + B$ is
- 1) $4x^2 - 2x + 4$ 2) $4x^2 + 2x - 4$ 3) $4x^2 - 2x - 4$ 4) $4x^2 + 2x + 4$

18. The value of $B + C$ is
 1) $2x^2 - x - 8$ 2) $2x^2 + x + 8$ 3) $2x^2 - x + 8$ 4) $2x^2 + x - 8$
19. The value of $2A + B + C$ is
 1) $16x^2 + 7x + 18$ 2) $16x^2 - 7x - 18$ 3) $16x^2 + 7x - 18$ 4) $16x^2 - 7x + 18$

Writeup-2:

The sum can be found by adding the similar terms in the expressions.

20. If $a = 2a - 5b + 4c$, $B = 5a - 2b + 2c$, then $A + B =$ ____
 1) $7(a - b + c)$ 2) $7(a - b) + 6c$ 3) $a - b + 6c$ 4) $7(a - b) + c$
21. If $A = x - y + 1$, $B = -2x + 7y + 3$, then $(2/5)A + B/5 =$ ____
 1) $5y + 1$ 2) $y + 5$ 3) $y - 5$ 4) $y + 1$
22. If $A = -3x - y + 5$, $B = x + 2y + 3$, then $3A - 5B =$ ____
 1) $14x - 13y$ 2) $13x - 14y$ 3) $-13x + 14y$ 4) $-14x - 13y$

MATRIX MATCHING TYPE

Add the two expressions

23. Column-I

- a) $3a - 4b$, $7a - 2b$
 b) $3a + 5b - 4c$, $2a - 5b - bc$
 c) $2ab - 5bc + 4ca$, $ab + 2bc - 5ac$
 d) $2a + 3b - 1$, $-4a + 5b - 5$

Column-II

- 1) $3ab - 3bc - ac$
 2) $10a - 6b$
 3) $-2a + 8b - 6$
 4) $5a - 10c$
 5) $5(a - 2c)$

INTEGER ANSWER TYPE

24. If $B = 2x^3 + 6x^2 - 7x + 8$ and $A + B = B$, then A is
 1) -1 2) 1 3) B 4) 0

SYNOPSIS - 3**Subtraction of polynomials :**

- Subtraction is the inverse process of addition.
- To every positive rational number there exists a negative rational number such that their sum is zero i.e., $A + (-A) = 0$.
 Here, the letter $(-A)$ is called the additive inverse of A .
 Eg : $A = -2x^2 - 8x + 7$
 \ Its additive inverse is $2x^2 + 8x - 7$.
- If $A + B = 0$, then B is called the additive inverse of A .
 Eg: $A = -2x^4 + 3x^2 + 5x$
 $B = 2x^4 - 3x^2 - 5x$ ∴ $A + B = -2x^4 + 3x^2 + 5x + 2x^4 - 3x^2 - 5x = 0$.
 \ B is the additive inverse of A .
- Sum of a polynomial and its additive inverse is zero.
 Eg : $A = 4x^3 - 4x^2 + 3x + 7$
 ∴ Additive inverse is $B = -4x^3 + 4x^2 - 3x - 7$
 \ $A + B = 4x^3 - 4x^2 + 3x + 7 - 4x^3 + 4x^2 - 3x - 7 = 0$.

WORK SHEET - 3

SINGLE ANSWER TYPE

1. If $B = -9x^2 + 3x - 7$, then the additive inverse of B is
 1) $9x^2 - 3x - 7$ 2) $9x^2 - 3x + 7$ 3) $-9x^2 - 3x - 7$ 4) $-9x^2 + 3x + 7$
2. If $A = \frac{-3x^2}{4} + \frac{2}{3}x + 7$ and $B = \frac{1}{4}x^2 - \frac{1}{3}x + 8$, then $A - B$ is
 1) $x^2 - x + 1$ 2) $-x^2 - x - 1$ 3) $-x^2 + x - 1$ 4) $x^2 + x + 1$
3. If $P = 2x^3 - 3x^2 - 5x + 6$ and $Q = \frac{1}{3}x^3 - \frac{3}{4}x^2 - \frac{5}{2}x + \frac{7}{3}$, then $Q - P$ is
 1) $\frac{5x^3}{3} + \frac{9x^2}{4} + \frac{5x}{2} - \frac{11}{3}$ 2) $\frac{-5x^3}{3} - \frac{9x^2}{4} + \frac{5x}{2} - \frac{11}{3}$
 3) $\frac{-5x^3}{3} - \frac{9x^2}{4} - \frac{5x}{2} - \frac{11}{3}$ 4) $\frac{5x^3}{3} + \frac{9x^2}{4} + \frac{5x}{2} - \frac{11}{3}$
4. If $A = -\frac{3}{2}x^3 - \frac{9}{7}x^2 + \frac{6x}{7} + 2$ and $A + B = 0$, then polynomial B is
 1) $\frac{-3x^3}{2} - \frac{9}{2}x^2 + \frac{6x}{7} + 2$ 2) $\frac{3x^3}{2} + \frac{9}{2}x^2 + \frac{6x}{7} + 2$
 3) $\frac{-3x^3}{2} - \frac{9}{2}x^2 - 6x - 2$ 4) $\frac{3x^3}{2} + \frac{9}{7}x^2 - \frac{6x}{7} - 2$
5. If $A = 2x^3 - 9x^2 - 6x + 7$ and $A + B = 5x^3 - 6x^2 - 8x + 9$, then the polynomial (A + 2) - A is
 1) $3x^3 - 3x^2 - 2x + 2$ 2) $3x^3 + 3x^2 - 2x + 2$
 3) $3x^3 + 3x^2 + 2x + 2$ 4) $-3x^3 - 3x^2 - 2x + 2$
6. If $A = 4x^3 - 9x^2 - 9x - 8$ and $A - B = -2x^3 - 8x^2 - 6x - 2$, then the polynomial $B = A - (A - 2)$ is
 1) $6x^3 - x^2 - 3x - 6$ 2) $6x^3 + x^2 + 3x + 6$
 3) $6x^3 + x^2 + 3x - 6$ 4) $-6x^3 - x^2 - 3x - 6$
7. Given $A = 2x^3 - 3x^2 + 6x + 7$ and $B = 4x^3 - 9x^2 - 3x + 7$, If C, D are additive inverses of A and B, then $D - C$ is
 1) $-2x^3 + 6x^2 + 9x$ 2) $-2x^3 + 5x^2 + 9x$
 3) $-2x^3 - 6x^2 + 9x$ 4) $-2x^3 - 6x^2 - 9x$
8. If $A - B = 2x^3 - 3x^2 + 8x - 7$ and $B = 5x^3 - 9x^2 + 6x - 8$, where $A = (A - 2) + B$, then the polynomial A is
 1) $7x^3 - 12x^2 + 14x + 18$ 2) $7x^3 - 12x^2 + 14x - 15$
 3) $7x^3 - 12x^2 - 14x + 15$ 4) $-7x^3 + 12x^2 - 14x - 15$

9. Given $C = \frac{-5}{6}x^2 - \frac{7}{6}x + \frac{3}{2}$ and $C + A = 0$. If $B = \frac{x^2}{6} - \frac{1}{6}x + \frac{1}{2}$ is added to A,

then the result is

- 1) $x^2 - x + 1$ 2) $-x^2 - x - 1$ 3) $x^2 + x - 1$ 4) $x^2 - x + 1$
 10. If $A = 7x^3 - 2x^2 - 9x + 6$, $B = 2x^3 - 8x^2 + 3x - 5$, $C = 2x^3 - 4x^2 - 8x + 7$,
 and $D = -3x^3 - 5x^2 + 6x + 7$, then $(A - 3) - (B - 4)$ is
 1) $5x^2 - 2x - 11$ 2) $5x^2 + 2x + 11$ 3) $5x^2 - 2x + 11$ 4) $-5x^2 - 2x - 11$

MULTI ANSWER TYPE

11. On multiplication of $\left(3x - \frac{4}{5}y^2x\right)$ by $\frac{1}{2}xy$, the result is

- 1) $\frac{3}{2}x^2y + \frac{5}{2}x^2y^3$ 2) $x^2y\left(\frac{3}{2} - \frac{2}{5}y^2\right)$
 3) $\frac{3}{2}x^2y - \frac{2}{5}x^2y^3$ 4) $\frac{5}{2}x^2y + \frac{3}{2}yx^2$
 12. The product of $100x \times (0.01x^4 - 0.01x^2)$ is
 1) $0.01x^4 - x^3$ 2) $0.5x^5 - x^3$ 3) $x^5 - x^3$ 4) $x^3 - x^5$
 13. Which of the following must be subtracted from $a^2 + b^2 + 2ab$ to get $-4ab + 2b^2$
 1) $2b^2 - 4ab$ 2) $a^2 - b^2 + 6ab$
 3) $a^2 - b(b + 6a)$ 4) $a^2 + b(-b + 6a)$

REASONING ANSWER TYPE

14. *Statement I* : If $a = 1$ and $b = 0.5$, then the value of $2.3a^5b^2 \times 1.2a^2b^2$ is 0.1725
Statement II : In a given expression, the process of replacing each variable by a given value of it is called substitution.
- Both Statements are true, Statement II is the correct explanation of Statement I.
 - Both Statements are true, Statement II is not correct explanation of Statement I.
 - Statement I is true, Statement II is false.
 - Statement I is false, Statement II is true.
15. *Statement I* : The perimeter of a triangle is $14a^2 + 20a + 13$. Two of its sides are $3a^2 + 5a + 1$ and $a^2 + 10a - 6$, then its 3rd side is $10a^2 + 5a + 18$
Statement II : The perimeter of a triangle is sum of its three sides.
- Both Statements are true, Statement II is the correct explanation of Statement I.
 - Both Statements are true, Statement II is not correct explanation of Statement I.
 - Statement I is true, Statement II is false.
 - Statement I is false, Statement II is true.

COMPREHENSION TYPE

Writeup-1

$$A = \frac{6}{5}x^2 - \frac{4}{5}x^3 + \frac{5}{6} + \frac{3}{2}x, B = \frac{x^3}{3} - \frac{5x^2}{2} + \frac{3}{5}x + \frac{1}{4}$$

16. The simplified form of (B - A) is

1) $\frac{17x^3}{15} - \frac{37x^2}{10} - \frac{9x}{10} - \frac{7}{12}$

2) $\frac{17x^3}{15} + \frac{37x^2}{10} + \frac{9x}{10} - \frac{7}{12}$

3) $\frac{17x^3}{15} - \frac{37x^2}{10} + \frac{9x}{10} + \frac{7}{12}$

4) $\frac{17x^3}{15} + \frac{37x^2}{10} + \frac{9x}{10} + \frac{7}{12}$

17. If $C = B - A + \frac{17x^3}{15} + \frac{37x^2}{10} + \frac{9x}{10} - \frac{7}{12}$, then the C =

1) $\frac{34x^3}{15} - \frac{7}{6}$

2) $\frac{34x^3}{15} - \frac{7}{12}$

3) $\frac{34x^3}{15} - \frac{6}{12}$

4) none

18. The simplified form of (A + C) is

1) $\frac{22x^3}{15} - \frac{1}{3} - \frac{6}{5}x^2 + \frac{3}{2}x$

2) $\frac{22x^3}{15} + \frac{1}{3} + \frac{6}{5}x^2 + \frac{3}{2}x$

3) $-\frac{22x^3}{15} + \frac{1}{3} + \frac{6}{5}x^2 - \frac{3}{2}x$

4) $\frac{22x^3}{15} + \frac{1}{4} + \frac{6}{5}x^2 + \frac{3}{2}x$

Writeup-2

$$\text{If } A = (x^2y - 1), B = -6x^2 + 15x^2y^3, C = 6x^2 - 15y^2, D = -2x^4y + x^2y^3$$

19. $A \times C =$ _____

1) $6x^4y + 15x^2y^3 - 6x^2 + 15y^2$

2) $6x^4y - 15x^2y^3 - 6x^2 + 15y^2$

3) $-6x^4y - 15x^2y^3 - 6x^2 - 15y^2$

4) $6x^4y + 15x^2y^3 + 6x^2 + 15y^2$

20. $(A \times C) - B =$ _____

1) $-2x^2y(4x^2 + 7y^2)$

2) $3y(2x^4 + 5y)$

3) $2y(3x^4 + 5y)$

4) $-3x^2y(2x^2 - 5y^2)$

21. $D - (A \times C) + B =$ _____

1) $-2x^2y(4x^2 + 7y^2)$

2) $2x^2y(4x^2 + 7y^2)$

3) $-2y(4x^4 + 7y)$

4) $-3x^2y(2x^2 - 5y^2)$

MATRIX MATCHING TYPE

22. Column-I

a) $(x - y) - (2x + y)$

b) $(3x + 2y) + (-4x - 4y)$

c) $(8y - 7x) - (6y - 8x)$

d) $2(x + y)$

Column-II

1) $-(-2y - x)$

2) $-(-2y + x)$

3) $-x - 2y$

4) $4y + 2x$

5) $2y + x$

INTEGER ANSWER TYPE

23. If $x = 2a^2 - 5a + 3$, $y = -3a^2 + a + 8$ and $z = 5a^2 - 6a - 5$, then the value of $x - (y - z)$ at

$a = -1$ is _____

KEY & HINTS

WORK SHEET – 1 (KEY)				
1) 3	2) 1	3) 3	4) 3	5) 2
6) 3	7) 3	8) 2	9) 3	10) 4
11) 3	12) 1	13) 3	14) 2,3,4	15) 1,2
16) 3,4	17) 1,2,3	18) 4	19) 3	20) 2,4
21) 2	22) 3	23) 4	24) 4	25) 4
26) 4	27) 1	28) 2	29) 4	30) 1
31) 1	32) 1	33) 2	34) 2	35) 4
36) 2	37) 3	38) 4	39) 4	40) 2
41) 2	42) 1	43) 3	44) A-4 B-3 C-2 D-1	45) A-4 B-2 C-1 D-1
46) A-4 B-3 C-2 D-1	47) A-3 B-1 C-2,5 D-4	48) 24	49) 5	50) 1

1. Substitute $n = 12$ in $\frac{n^2}{2} + \frac{n}{2}$

$$\Rightarrow \frac{n^2}{2} + \frac{n}{2} = \frac{12^2}{2} + \frac{12}{2} = \frac{144}{2} + \frac{12}{2} = 72 + 6 = 78$$

$\therefore$ Value of the expression is 78.

2. $\frac{7}{3}x - \frac{7}{6} = 0$

Transpose $-\frac{7}{6}$ to RHS

$$\Rightarrow \frac{7x}{3} = \frac{7}{6}$$

Transpose $\frac{7}{3}$ to RHS

$$\Rightarrow x = \frac{7}{6} \times \frac{3}{7} = \frac{1}{2}$$

$\therefore$ Zero of the given polynomial is $\frac{1}{2}$

3. $x = -\frac{5}{4} \Rightarrow 4x = -5$

$\therefore 4x + 5$ is the required polynomial

4. $A = -8x^2 - 6x + 10$

$$\text{Put } x = \frac{1}{2} \Rightarrow A = -8 \times \left(\frac{1}{2}\right)^2 - 6 \times \frac{1}{2} + 10$$

$$= -8 \times \frac{1}{4} - 6 \times \frac{1}{2} + 10 = -2 - 3 + 10 = 5$$

5. $3x^{4-1} + 2x^{3-1} + 6x^{2-1} + 8 = 3x^3 + 2x^2 + 6x + 8$ is a 3rd degree polynomial

6. $\frac{2}{7}x^{-2}y^5z$

7. Put $x = \frac{a}{2}$ in $4x^2 + 8x + 18$

$$\Rightarrow 4x^2 + 8x + 18 = 4 \times \left(\frac{a}{2}\right)^2 + 8 \times \frac{a}{2} + 18 = 4 \times \frac{a^2}{4} + 8 \times \frac{a}{2} + 18 = a^2 + 4a + 18$$

8. Put $x = \frac{9}{13}$ in $\frac{-26}{3} - \frac{13x}{27}$

$$\Rightarrow \frac{-26}{3} - \frac{13x}{27} = \frac{-26}{3} - \frac{13}{27} \times \frac{9}{13} = \frac{-26}{3} - \frac{1}{3} = \frac{-27}{3} = -9$$

9. Degree of the polynomial is $m + 4$

10. Put $n = 10$ in $\frac{n(n+1)(2n+1)}{6}$

$$\Rightarrow \frac{n(n+1)(2n+1)}{6} = \frac{10(10+1)(2 \times 10 + 1)}{6} = \frac{10 \times 11 \times 21}{6} = 385$$

11. The polynomial is $\frac{1}{2}x^5 + 3x^4 + 2x^3 + 3x^2 + 6$

$\therefore$ The degree of the polynomial is 5.

12. Degree of the monomial $\frac{3}{5}x^2y^6z^7$ is $2 + 6 + 7 = 15$
13. $x = a + 2$ and $a = 8 \Rightarrow x = 10$
 $\therefore$ The value of $3x + 5 = 3 \times 10 + 5 = 35$
14. Key: (2,3,4) ; Sol:- The coefficient of x in $9xy$ is '9y'
 The coefficient of 'a' in $(-7abc)$ is $(-7bc)$; The coefficient of 'xyz' on $(-xyz)$ is -1
 The coefficient of b in $-abc$ is $(-ac)$
15. Key: (1,2) ; Sol:- In $6x^2y + 5xy^2 - 8xy^2 - 7yx^2, 6x^2y, -7x^2y$ are like terms
 Also $5xy^2, -8xy^2$ are like terms
16. Key: (3, 4) ; Sol:- 1. 3 more than $x = x + 3$ (True)
 2. One third of sum of x and $y =$ one third of $(x + y) = \frac{1}{3}(x + y)$ (True)
 3. The quotient of x by $y = \frac{x}{y}$; product of x and $y = xy$
 now the quotient of x by y added to the product of x and $y = \frac{x}{y} + xy$
 but it is given that $x + y + \frac{x}{y}$ $\therefore$ (3) is false
 4. The quotient of x by $y = \frac{x}{y}$; Now 5 less than $\frac{x}{y} = \frac{x}{y} - 5$; but it is given that $\frac{x}{y} + 5$
 $\therefore$ (4) is false.
17. Key: (1,2,3) ; Sol:- The degree of $x^2 + xy^2 + y^3$ is 3 (True)
 The degree of $m^2n^3 + mn^2 + 4$ is 5 (True) ; The degree of $p^2q^2 + pq^2 + 1$ is 4 (True)
18. Key: 4 ; Sol:- Clearly (a) and (b) are not polynomials only (c) is a polynomials
 $\therefore$ (a) and (b) are false
19. Key: 3 ; Sol:- $2x^{3^2 \times 2^3} + 3x^{4^2 \times 3^2} + 5x^{4^2 \times 3^2} + 5x^{4^2 \times 3^3} = 2x^{54} + 3x^{144} + 5x^{144} + 5x^{128}$
 $\therefore$ degree = 144.
20. Key: 2, 4 ; Sol:- $A = (m+n)p/2 = (120.6)(3.5)/2 = 211.05$
 Clearly, $A < 211.50$ & Also, $A < 211.105$
21. Key: 2 ; Sol:- $(25a^4b) \times (-2a^2b^2) \times (-2.1a^3b^3) = -105 \times a^{4+2+3} b^{1+2+3} = -105a^9b^6$
 putting $a = 1$ and $b = \frac{1}{2}$ we get $-105a^9b^6 = -105(1)^9(1/2)^6 = -105/64$

22. Key: 3; Sol:- $(a + b)(a + b) - (a - b)(a - b) + (a^2 - b^2) = (7 + 5)(7 + 5) - (7 - 5)(7 - 5) + (7^2 - 5^2)$
 $= 12 \times 12 - 2 \times 2 + 49 - 25 = 144 - 4 + 24 = 164.$
23. Key: 4 ; Sol:- Clearly statement 2 is true (by conceptual formulae)
 Statement - 1: $(a + b)^2 + (a - b)^2 + (a^2 - b^2)$
 $= a^2 + b^2 + 2ab + a^2 - 2ab + b^2 + a^2 - b^2 = 3a^2 + b^2$
 $= 3(3)^2 + (2)^2$ (Q $a = 3, b = 2$)
 $= 37 + 4 = 41$ -----> statement - 1 is false.
24. Key: 4 ; Sol:- Statement I: $(x - 10) + (y + 20) + 10 = x + y + 20$
 $\therefore$ Statement I is False.
 Statement II: $[(4 \times q) + 5 + r] - (p + 10) = 4q + 5 + r - p - 10 = 4q + r - p - 5$
 $\therefore$ Statement 2 is True.
25. Key: 4 ; Sol:- Clearly statement - 2 is true
 St-1 : $A = 4x^{2m}y^{4n}z^{3p}$; Since $2m = 4n = 3p = 24 \Rightarrow m = 12, n = 6, p = 8$
 $\therefore$ Degree of A = $2m + 4n + 3p = 2(12) + 4(6) + 3(8) = 24 + 24 + 24 = 72$
 $\therefore$ statement - 1 is false.
26. Key: 4 ; Sol:- Conceptual.
27. Key: 1 ; Sol:- Clearly, St-2 is true.
 $5a^6 \times (-10)ab^2 \times (-2.1)a^2b^3 = 105a^9b^5 = 105 \times (1)^9(1/2)^5 = 105/32$ -----> St-1 is true.
 And St-2 is correct explanation of St-1
28. Key: 2; Sol:- $x^2 + y^2 + z^2 - 2xy - 2yz - 2zx = (2)^2 + (-2)^2 + (3)^2 - 2(2)(-2) - 2(-2)(3) - 2(3)(2) = 25$
 st-1 is true; $p^2 + q^2 + r^2 - pq - qr - pr = (3)^2 + (-5)^2 + (4)^2 - (3)(-5) - (-5)(4) - (4)(3) = 73$
 st-2 is true. But st-2 is not correct explanation of st-1
29. Key: 4 ; Sol:- Given $A = 18x^3y^2z^4, B = 12x^2y^3z^3$ and $C = x^4y^3$;
 $\therefore A \times C = (18x^3y^2z^4) \times (x^4y^3) = 18x^7y^5z^4$ -----> So, Both (a) & (b) is false
30. Key: 1 ; Sol:- $B \times C = (12x^2y^3z^3) \times (x^4y^3) = 12x^6y^6z^3$
31. Key: 1 ; Sol:- $A \times B \times C = (18x^3y^2z^4) \times (12x^6y^6z^3) = 216x^9y^8z^7$
32. Key: 1 ; Sol:- Given $A = 6a^{p+2} + 5a^{2p+3} + 7a^{3p+4}$; Since $p = 2, q = 3$
 $\therefore A = 6.a^{2+2} + 5.a^{2(2)+3} + 7.a^{3(2)+4} \Rightarrow A = 6a^4 + 5a^7 + 7a^{10}$; $\therefore$ degree of A = 10
33. Key: 2 ; Sol:- Since $B = 4b^{q+3} + 5b^{3q+2} + 8b^{4q+3}$; Given $q = 3$
 $\therefore B = 4.b^{3+3} + 5.b^{3(3)+2} + 8.b^{4(3)+3} \Rightarrow 4.b^6 + 5.b^{11} + 8.b^{15}$; $\therefore$ degree of B = 15
34. Key: 2 ; Sol:- Clearly, degree of A < degree of B. i.e., $10 < 15$
35. Key: 4 ; Sol:- Conceptual.
36. Key: 2 ; Sol:- Conceptual.
37. Key: 3 ; Sol:- Conceptual.

38. Key: 4 ; Sol:- $R^2 - r^2 = (R + r)(R - r) = (2.6 + 2.3)(2.6 - 2.3) = 4.9 \times 0.3 = 1.47$.

39. Key: 4 ; Sol:- $(V/W) = 115.5/(22/7) = 115.5 \times (7/22) = (73.5/2) = (147/4)$

40. Key: 2 ; Sol:- $A = \frac{115.5}{\frac{22}{7}[(2.6)^2 - (2.3)^2]} = \frac{115.5 \times 7}{22 \times (6.76 - 5.29)}$

$$= \frac{808.5}{22 \times 1.47} = \frac{808.5}{32.34} = 25$$

41. Key: 2; Sol:- $2x^2 + (7/2)xy + 5y^2 = 2(1)^2 + (7/2)(1)(2) + 5(2)^2 = 29$

42. Key: 1; Sol:- $a^2 - bc + c^2 - b^2 = (0)^2 - (1)(2) + (2)^2 - (1)^2 = 1$

43. Key: 3; Sol:- $x^3 - xy/3 + 7/2 y^2 = (2)^3 - 2(3)/3 + 7/2(3)^2 = 75/2$

44. Key: $a \rightarrow 4$; $b \rightarrow 3$; $c \rightarrow 2$; $d \rightarrow 1$; Sol:- Conceptual

45. $(a) \rightarrow 4$, $(b) \rightarrow 2$, $(c) \rightarrow 1$, $(d) \rightarrow 1$; Sol:- Conceptual.

46. Key: $a \rightarrow 4$; $b \rightarrow 3$; $c \rightarrow 2$; $d \rightarrow 1$;

Sol:- a) degree of $5x$ is 1

b) degree of $15x^2 - x + 2$ is 2

c) degree of $-x^4 + 2x + 11$ is 4

d) degree of $x^2 - x^5$ is 5.

47. key: $a-3$, $b-1$, $c-2,5$, $d-4$

Sol:- a) $2h+3m = 2(5)+3(3)=10+9=19$;

b) $x^2-4x+4=(x-2)^2=(3-2)^2=1$

c) $x^2+y^2 = 5^2+12^2=25+144=169$;

d) $xy+yz+zx = 4(3)+(3)(-2)+(-2)(4) = 12-6-8=-2$

48. Key: (24) ; Sol:- $\frac{x}{3} - 6 = 2 \Rightarrow \frac{x}{3} = 8 \Rightarrow x = 24$.

49. Key: 5 ; Sol:- Conceptual.

50. Key: 1;

$$\text{Sol:- } \frac{(+1)^3 + (-2)^3 + (-3)^3 - 3(1)(-2)(-3)}{(1)(-2) + (-2)(-3) + (-3)(1) - (1+4+9)} = \frac{1-8-27-18}{-2+6-3-14} = \frac{1-53}{6-19} = \frac{-52}{-13} = 4$$

WORK SHEET – 2 (KEY)

1) 3	2) 2	3) 1	4) 3	5) 1
6) 1	7) 3	8) 2	9) 3	10) 1
11) 2	12) 3	13) 3	14) 3	15) 4
16) 3	17) 1	18) 2	19) 4	20) 2
21) 4	22) 4	23) A-2 B-4,5 C-1 D-3	24) 4	

1. $\frac{3}{4}x^3 + \frac{5}{6}x^3 - \frac{2}{3}x^3 + \frac{7}{2}x^3 = \frac{9x^3 + 10x^3 - 8x^3 + 42x^3}{12} = \frac{x^3}{12}(9 + 10 - 8 + 42) = \frac{53x^3}{12}$
2. $3x^3 + 7x^3 + 8x^3 - 2x^2 - 6x^2 - 9x^2 - 8x + 9x + 6x$
 $= x^3(3 + 7 + 8) + x^2(-2 - 6 - 9) + x(-8 + 9 + 6)$
 $= 18x^3 - 17x^2 + 7x$
3. Ascending order of $-3x^3 + 7x^2 - 9x^4 + 6x - 8$ is $-8 + 6x + 7x^2 - 3x^3 - 9x^4$
4. $A = x(-7 - 3 - 5) \Rightarrow A = -15x$
 $B = x(9 + 3 + 2) \Rightarrow B = 14x$
 $\therefore A + B = -15x + 14x = -x$
5. $A = \frac{1}{2}x - \frac{1}{3}x = x\left(\frac{1}{2} - \frac{1}{3}\right) = x\left(\frac{3-2}{6}\right) = \frac{1}{6}x$
 $B = \frac{1}{3}x - \frac{1}{4}x = x\left(\frac{1}{3} - \frac{1}{4}\right) = x\left(\frac{4-3}{12}\right) = \frac{1}{12}x$
 $\therefore A - B = \frac{1}{6}x - \frac{1}{12}x = x\left(\frac{1}{6} - \frac{1}{12}\right) = x\left(\frac{2-1}{12}\right) = \frac{1}{12}x$
6. $3x^3 - x^3 - 5x^2 + 2x^2 - 9x + x - 7 + 4$
 $= 2x^3 - 3x^2 - 8x - 3$
7. The descending order of the given expression is $-9x^4 - 6x^3 - 2x^2 + 3x + 2$
8. $\frac{-7}{5}x^3 + \frac{3}{4}x^3 + \frac{7}{2}x^3 + \frac{9x^3}{3} + \frac{9x^3}{60}$
 $= \frac{-84x^3 + 45x^3 + 210x^3 + 180x^3 + 9x^3}{60} = \frac{360}{60}x^3 = 6x^3$
9. $P = 2x - 3x + 5x = 4x$, $Q = -8x + 3x + 9x = 4x$
 $R = -8x - 6x - 7x = -21x$
 $\therefore (P + Q) - R = (4x + 4x) - (-21x)$
 $= 8x + 21x = x(8 + 21) = 29x$
10. $A = -5x^3 + 2x^2$
 $B = 2x^2 - 5x$
 $C = -7x + 1$
 $\therefore A + B + C = -5x^3 + 2x^2 + 2x^2 - 5x - 7x + 1$
 $= -5x^3 + x^2(2 + 2) + x(-5 - 7) + 1$
 $= -5x^3 + 4x^2 - 12x + 1$
11. $4x^3y^2 + 3x^2y^3 - 9x^2y^3 - 8x^2y^5 + 6x^2y^5 - 9x^3y^4$
 $= 4x^3y^2 - 6x^2y^3 - 2x^2y^5 - 9x^3y^4$
12. $x^3(0.5 + 1.85 + 2.96 - 4.71) + x^4(1.25 - 2.5 + 3.6 - 4.71)$
 $0.6x^3 - 2.36x^4$

13. Key: 3; Sol:-

Required sum

$$\begin{aligned}
 &= \left(5x^2 - \frac{1}{3}x + \frac{5}{2}\right) + \left(-\frac{1}{2}x^2 + \frac{1}{2}x - \frac{1}{3}\right) + \left(-2x^2 + \frac{1}{5}x - \frac{1}{6}\right) \\
 &= 5x^2 - \frac{1}{2}x^2 - 2x^2 - \frac{1}{3}x + \frac{1}{2}x + \frac{1}{5}x + \frac{5}{2} - \frac{1}{6} - \frac{1}{3} \\
 &= \left(5 - \frac{1}{2} - 2\right)x^2 + \left(-\frac{1}{3} + \frac{1}{2} + \frac{1}{5}\right)x + \left(\frac{5}{2} - \frac{1}{6} - \frac{1}{3}\right) \\
 &= \frac{5}{2}x^2 + \frac{11}{30}x + 2
 \end{aligned}$$

14. Key: 3; Sol:- $\left(\frac{7}{2} + \frac{3}{2}\right)x^3 + \left(\frac{-1}{2} + \frac{7}{4} + \frac{3}{2}\right)x^2 + \left(\frac{-5}{2} - 1\right)x + \left(\frac{5}{3} + \frac{1}{3} - 2\right)$

$$= \left(\frac{10}{2}\right)x^3 + \left(\frac{-2+7+6}{4}\right)x^2 + \left(\frac{-7}{2}\right)x + \left(\frac{+6-6}{3}\right) = 5x^3 + \frac{11}{4}x^2 - \frac{7}{2}x$$

15. Key: 4; Sol:- Clearly, St-2 is true.

$$A + B + C = \left(5x^2 - \frac{1}{3}x + \frac{5}{2}\right) + \left(-\frac{1}{2}x^2 + \frac{1}{2}x - \frac{1}{3}\right) + \left(-2x^2 + \frac{1}{2}x - \frac{1}{6}\right) = \left(\frac{5}{2}x^2 + \frac{11}{30}x + 2\right)$$

So, St-1 is false.

16. Key: 3;; Sol:- clearly st-2 is false ; st-1 $8p^2 - 9q^2 - 6p^2 + 5q^2 = 2p^2 - 4q^2$ st-1 is true

17. Key: 1 ; Sol:- $A + B = (7x^2 - 4x + 5) + (-3x^2 + 2x - 1) = 4x^2 - 2x + 4$

18. Key: 2; Sol: $B + C = (-3x^2 + 2x - 1) + (5x^2 - x + 9) = 2x^2 + x + 8$

19. Key: 4; Sol:- $2A + B + C = 2(7x^2 - 4x + 5) + 2x^2 + x + 8 = 16x^2 - 7x + 18$.

20. key: 2; Sol:- $A+B = 2a-5b+4c+5a-2b+2c=7a-7b+6c$

21. key: 4; Sol:- $2/5A+B/5=$

$$\frac{2A+B}{5} = \frac{2(x-y+1) + (-2x+7y+3)}{5} = \frac{5(y+1)}{5} = Y+1$$

22. key: 4; Sol:- $3A-5B=3(-3x-y+5)-5(x+2y+3)=-9x-3y+15-5x-10y-15=-14x-13y$

23. Key: a-2; b-(4,5), c-1, d-3

; Sol:- a) $3a-4b+7a-2b=10a-6b$; b) $3a+5b-4c+2a-5b-6c=5(a-2c)$

c) $2ab-5bc+4ca+ab+2bc-5ac=3ab-3bc-ac$ d) $2a+3b-1-4a+5b-5=-2a+8b-6$

24. Key: 4; Sol:- $A + B = B$ $\therefore A = 0$

WORK SHEET – 3 (KEY)				
1) 2	2) 3	3) 2	4) 4	5) 2
6) 1	7) 1	8) 2	9) 3	10) 2
11) 2,3	12) 3	13) 2,4	14) 1	15) 1
16) 1	17) 1	18) 4	19) 2	20) 2
21) 3	22) A-2,3 B-2,3 C-1,5 D-4	23) 12		

1. $B = -9x^2 + 3x - 7$,
 $\setminus$ Additive inverse of B is $9x^2 - 3x + 7$.

2. $A - B = -\frac{3}{4}x^2 - \frac{1}{4}x^2 + \frac{2}{3}x + \frac{1}{3}x + 7 - 8 = -\frac{4x^2}{4} + \frac{3}{3}x + 7 - 8 = -x^2 + x - 1$

3. $Q - P = \frac{1}{3}x^3 - 2x^3 - \frac{3}{4}x^2 + 3x^2 - \frac{5}{2}x + 5x + \frac{7}{3} - 6 = \frac{-5x^3}{3} - \frac{9x^2}{4} + \frac{5x}{2} - \frac{11}{3}$

4. B is the additive inverse of A. (Q $A + B = 0$)

$\text{p } A = -\frac{3}{2}x^3 - \frac{9}{7}x^2 + \frac{6x}{7} + 2$

$\setminus B = \frac{3}{2}x^3 + \frac{9}{7}x^2 - \frac{6x}{7} - 2$

5. $A + B = +5x^3 - 6x^2 - 8x + 9$

$A = +2x^3 - 9x^2 - 6x + 7$
 $\quad \quad \quad - \quad + \quad + \quad -$

$\therefore (A + B) - A = 3x^3 + 3x^2 - 2x + 2$

6. $B = A - (A - B)$

$A = 4x^3 - 9x^2 - 9x - 8$
 $A - B = -2x^3 - 8x^2 - 6x - 2$
 $\quad \quad \quad + \quad + \quad + \quad +$

$\therefore B = A - (A - B) = 6x^3 - x^2 - 3x - 6$

7. D is the additive inverse of B and C is the additive inverse of A

$$D = -4x^3 + 9x^2 + 3x - 7$$

$$C = -2x^3 + 3x^2 - 6x - 7$$

$$\begin{array}{cccc} + & - & + & + \end{array}$$

$$\therefore D - C = -2x^3 + 6x^2 + 9x$$

8. $A = (A - B) + B$

$$\Rightarrow A - B = 2x^3 - 3x^2 + 8x - 7$$

$$B = 5x^3 - 9x^2 + 6x - 8$$

$$\therefore A = (A - B) + B = 7x^3 - 12x^2 + 14x - 15$$

9. A is the additive inverse of C. (Q $C + A = 0$)

$$A = \frac{5}{6}x^2 + \frac{7}{6}x - \frac{3}{2}, B = \frac{x^2}{6} - \frac{1}{6}x + \frac{1}{2}$$

$$\therefore A + B = \frac{5x^2}{6} + \frac{x^2}{6} + \frac{7}{6}x - \frac{1}{6}x - \frac{3}{2} + \frac{1}{2} = \frac{6x^2}{6} + \frac{6x}{6} - \frac{2}{2} = x^2 + x - 1$$

10. $A - C = 7x^3 - 2x^2 - 9x + 6 - (2x^3 - 4x^2 - 8x + 7) = 5x^3 + 2x^2 - x - 1$

$$B - D = 2x^3 - 8x^2 + 3x - 5 - (-3x^3 - 5x^2 + 6x + 7) = 5x^3 - 3x^2 - 3x - 12$$

$$A - C = 5x^3 + 2x^2 - x - 1$$

$$B - D = 5x^3 - 3x^2 - 3x - 12$$

$$\begin{array}{cccc} - & + & + & + \end{array}$$

$$\therefore (A - C) - (B - D) = 5x^2 + 2x + 11$$

11. Key: (2,3); Sol:- $\left(3x - \frac{4}{5}y^2x\right) \times \frac{1}{2}xy = \frac{3}{2}x^2y - \frac{2}{5}x^2y^3 = x^2y\left(\frac{3}{2} - \frac{2}{5}y^2\right)$

12. Key: 3 ; Sol:- $100x \times 0.01x^4 - 100x \times 0.01x^2 = (100 \times 0.01)x^5 - (100 \times 0.01)x^3$
 $= \left(100 \times \frac{1}{100}\right)x^5 - \left(100 \times \frac{1}{100}\right)x^3 = x^5 - x^3$

13. Key: (2,4) ; Sol: $a^2 + b^2 + 2ab - x = -4ab + 2b^2$ **P** $a^2 + b^2 + 2ab + 4ab - 2b^2 = x$
P $x = a^2 - b^2 + 6ab$

14. key: 1 ; Sol:- Clearly St- 2 is true. Now, $2.3a^5b^2 \times 1.2a^2b^2 = (2.3) \times (1.2)a^7b^4$
 $= (2.3) \times (1.2) \times (1)^7(0.5)^4 = 0.1725$

So, St-1 is true & St- 2 is correct explanation of St- 1 .

15. Key: 1; Sol:- Clearly st-2 is true; $(3a^2+5a+1)+(a^2+10a-6)=4a^2+15a-5$
 Now $(14a^2+20a+13)-(4a^2+15a-5) = 10a^2+5a+18$ so, st-1 is true
 and st-2 is correct explanation of st-2

16. Key: 1 ;

Sol:-

$$\left(\frac{x^3}{3} - \frac{5x^2}{2} + \frac{3}{5}x + \frac{1}{4}\right) - \left(\frac{6}{5}x^2 - \frac{4}{5}x^3 + \frac{5}{6} + \frac{3}{2}x\right) = \frac{17x^3}{15} - \frac{37x^2}{10} - \frac{9x}{10} - \frac{7}{12}.$$

17. Key: 1 ; Sol:- $C = \frac{17x^3}{15} - \frac{37x^2}{10} - \frac{9x}{10} - \frac{7}{12} + \frac{17x^3}{15} + \frac{37x^2}{10} + \frac{9x}{10} - \frac{7}{12} = \frac{34x^3}{15} - \frac{7}{6}$

18. Key: 4 ; Sol:- $A+C = \left(\frac{6}{5}x^2 - \frac{4}{5}x^3 + \frac{5}{6} + \frac{3}{2}x\right) + \frac{34x^3}{15} - \frac{7}{6} = \frac{22x^3}{15} + \frac{1}{4} + \frac{6}{5}x^2 + \frac{3}{2}x$

19. Key: 2; Sol:- $A \times C = (x^2y - 1) \times (6x^2 - 5y^2) = 6x^4y - 15x^2y^3 - 6x^2 + 15y^2$

20. Key 2; Sol:- $(A \times C) - B = 6x^4y - 15x^2y^3 - 6x^2 + 15y^2 + 6x^2 + 15x^2y^3 = 3y(2x^4 + 5y)$

21. Key: 3; Sol:- $D - (A \times C) + B = D - [(A \times C) - B] = -2x^4y - 6x^4y - 15y^2$
 $= -8x^4 - 14y^2 = -2y(4x^4 + 7y)$

22. Key: a® (2,3) ; b® (2,3) ; c® (1,5); d® 4;

Sol:- a) $(x - y) - (2x + y) = x - y - 2x - y = -x - 2y$

b) $(3x + 2y)(-4x - 4y) = 3x + 2y - 4x - 4y = -x - 2y$

c) $(8y - 7x) - (6y - 8x) = (8y - 6y) - 7x + 8x = 2y + x$

d) $2(x + 2y) = 2x + 4y$

23. Key: 12 ; Sol:- $y-z = (-3a^2 + a + 8) - (5a^2 - 6a - 5) = -8a^2 + 7a + 13$

$$x - (y - z) = (2a^2 - 5a + 3) - (-8a^2 + 7a + 13) = 10a^2 - 12a - 10$$

if $a = -1$, then $x - (y - z) = 10(-1)^2 - 12(-1) - 10 = 12.$ $\therefore$