

DIFFERENTIABILITY

2.4.1 DIFFERENTIABILITY OF A FUNCTION AT A POINT

(1) **Meaning of differentiability at a point** : Consider the function $f(x)$ defined on an open interval (b, c) let $P(a, f(a))$ be a point on the curve $y = f(x)$ and let $Q(a-h, f(a-h))$ and $R(a+h, f(a+h))$ be two neighbouring points on the left hand side and right hand side respectively of the point P.

$$\text{Then slope of chord } PQ = \frac{f(a-h) - f(a)}{(a-h) - a} = \frac{f(a-h) - f(a)}{-h}$$

$$\text{and, slope of chord } PR = \frac{f(a+h) - f(a)}{a+h-a} = \frac{f(a+h) - f(a)}{h}.$$

⊙ As $h \rightarrow 0$, point Q and R both tends to P from left hand and right hand respectively. Consequently, chords PQ and PR becomes tangent at point P.

$$\text{Thus, } \lim_{h \rightarrow 0} \frac{f(a-h) - f(a)}{-h} = \lim_{h \rightarrow 0} (\text{slope of chord } PQ) = \lim_{Q \rightarrow P} (\text{slope of chord } PQ)$$

Slope of the tangent at point P, which is limiting position of the chords drawn on the left hand side of point P and $\lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} = \lim_{h \rightarrow 0} (\text{slope of chord } PR) = \lim_{R \rightarrow P} (\text{slope of chord } PR).$

⇒ Slope of the tangent at point P, which is the limiting position of the chords drawn on the right hand side of point P.

$$\text{Now, } f(x) \text{ is differentiable at } x = a \Leftrightarrow \lim_{h \rightarrow 0} \frac{f(a-h) - f(a)}{-h} = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

⇔ There is a unique tangent at point P.



Thus, $f(x)$ is differentiable at point P, iff there exists a unique tangent at point P. In other words, $f(x)$ is differentiable at a point P iff the curve does not have P as a corner point. i.e., "the function is not differentiable at those points on which function has jumps (or holes) and sharp edges."

Let us consider the function $f(x) = |x-1|$, which can be graphically shown,

Which show $f(x)$ is not differentiable at $x=1$. Since, $f(x)$ has sharp edge at $x=1$.

Mathematically : The right hand derivative at $x=1$ is 1 and left-hand derivative at $x=1$ is -1 . Thus, $f(x)$ is not differentiable at $x=1$.

(2) **Right hand derivative** : Right hand derivative of $f(x)$ at $x = a$, denoted by $f'(a+0)$ or $f'(a+)$, is the $\lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}.$



(3) **Left hand derivative** : Left hand derivative of $f(x)$ at $x=a$ denoted by $f(a-0)$ or $f(a-)$, is the $\lim_{h \rightarrow 0} \frac{f(a-h) - f(a)}{-h}$.

(4) A function $f(x)$ is said to be differentiable (finitely) at $x=a$ if $f(a+0) = f(a-0) = \text{finite}$

i.e., $\lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} = \lim_{h \rightarrow 0} \frac{f(a-h) - f(a)}{-h} = \text{finite}$ and the common limit is called the derivative of $f(x)$ at $x=a$, denoted by $f'(a)$. Clearly, $f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$ $\{x \rightarrow a \text{ from the left as well as from the right}\}$.

Example: 1 Consider $f(x) = \begin{cases} x^2, & x \neq 0 \\ 0, & x = 0 \end{cases}$

- (a) $f(x)$ is discontinuous everywhere
- (b) $f(x)$ is continuous everywhere but not differentiable at $x=0$
- (c) $f'(x)$ exists in $(-1, 1)$
- (d) $f'(x)$ exists in $(-2, 2)$

Solution: (b) We have, $f(x) = \begin{cases} x^2, & x \neq 0 \\ 0, & x = 0 \end{cases} = \begin{cases} \frac{x^2}{x}, & x > 0 \\ 0, & x = 0 \\ \frac{x^2}{-x}, & x < 0 \end{cases}$

$$\Rightarrow \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} -x = 0, \quad \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} x = 0 \text{ and } f(0) = 0.$$

So $f(x)$ is continuous at $x=0$. Also $f(x)$ is continuous for all other values of x . Hence, $f(x)$ is everywhere continuous.

$$\text{Also, } Rf(0) = f(0+0) = \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h - 0} = \lim_{h \rightarrow 0} \frac{h - 0}{h} = 1$$

$$\text{i.e. } Rf(0) = 1 \text{ and } Lf(0) = f(0-0) = \lim_{h \rightarrow 0} \frac{f(-h) - f(0)}{-h} = \lim_{h \rightarrow 0} \frac{-h}{-h} = -1$$

i.e. $Lf(0) = -1$ So, $Lf(0) \neq Rf(0)$ i.e., $f(x)$ is not differentiable at $x=0$.

Example: 2 If the function f is defined by $f(x) = \frac{x}{1+|x|}$, then at what points f is differentiable

- (a) Everywhere
- (b) Except at $x = \pm 1$
- (c) Except at $x = 0$
- (d) Except at $x = 0$ or ± 1

Solution: (a) We have, $f(x) = \frac{x}{1+|x|} = \begin{cases} \frac{x}{1+x}, & x > 0 \\ 0, & x = 0 \\ \frac{x}{1-x}, & x < 0 \end{cases}$; $Lf(0) = \lim_{h \rightarrow 0} \frac{f(-h) - f(0)}{-h} = \lim_{h \rightarrow 0} \frac{\frac{-h}{1+h} - 0}{-h} = 1$

$$Rf(0) = \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{\frac{h}{1+h} - 0}{h} = \lim_{h \rightarrow 0} \frac{1}{1+h} = 1$$

So, $Lf(0) = Rf(0) = 1$

So, $f(x)$ is differentiable at $x=0$; Also $f(x)$ is differentiable at all other points.

Hence, $f(x)$ is everywhere differentiable.

Example: 3 The value of the derivative of $|x-1| + |x-3|$ at $x=2$ is

- (a) -2
- (b) 0
- (c) 2
- (d) Not defined

Solution: (b) Let $f(x) = |x-1| + |x-3| = \begin{cases} -(x-1) - (x-3) & , x < 1 \\ (x-1) - (x-3) & , 1 \leq x < 3 \\ (x-1) + (x-3) & , x \geq 3 \end{cases} = \begin{cases} -2x+4 & , x < 1 \\ 2 & , 1 \leq x < 3 \\ 2x-4 & , x \geq 3 \end{cases}$

Since, $f(x) = 2$ for $1 \leq x < 3$. Therefore $f(x) = 0$ for all $x \in (1, 3)$.

Hence, $f(x) = 0$ at $x = 2$.

Example: 4 The function f defined by $f(x) = \begin{cases} \frac{\sin x^2}{x} & , x \neq 0 \\ 0 & , x = 0 \end{cases}$


- (a) Continuous and derivable at $x = 0$ (b) Neither continuous nor derivable at $x = 0$
 (c) Continuous but not derivable at $x = 0$ (d) None of these

Solution: (a) We have, $\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \frac{\sin x^2}{x} = \lim_{x \rightarrow 0} \left(\frac{\sin x^2}{x^2} \right) x = 1 \times 0 = 0 = f(0)$

So, $f(x)$ is continuous at $x = 0$, $f(x)$ is also derivable at $x = 0$, because $\lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} = \lim_{x \rightarrow 0} \frac{\sin x^2}{x^2} = 1$ exists finitely.

Example: 5 If $f(x) = |\log |x||$, then

- (a) $f(x)$ is continuous and differentiable for all x in its domain
 (b) $f(x)$ is continuous for all x in its domain but not differentiable at $x = \pm 1$.
 (c) $f(x)$ is neither continuous nor differentiable at $x = \pm 1$
 (d) None of these



Solution: (b) It is evident from the graph of $f(x) = |\log |x||$ that $f(x)$ is everywhere continuous but not differentiable at $x = \pm 1$.

Example: 6 The left hand derivative of $f(x) = [x] \sin(\pi x)$ at $x = k$ (k is an integer), is

- (a) $(-1)^k(k-1)\pi$ (b) $(-1)^{k-1}(k-1)\pi$ (c) $(-1)^k k\pi$ (d) $(-1)^{k-1} k\pi$

Solution: (a) $f(x) = [x] \sin(\pi x)$

If x is just less than k , $[x] = k - 1$. $\therefore f(x) = (k-1)\sin(\pi x)$, when $x < k \quad \forall k \in I$

Now L.H.D. at $x = k$,

$$= \lim_{x \rightarrow k} \frac{(k-1)\sin(\pi x) - k\sin(\pi k)}{x - k} = \lim_{x \rightarrow k} \frac{(k-1)\sin(\pi x)}{(x - k)} \quad [\text{as } \sin(\pi k) = 0, k \in \text{integer}]$$

$$= \lim_{h \rightarrow 0} \frac{(k-1)\sin(\pi(k-h))}{-h} \quad [\text{Let } x = (k-h)]$$

$$= \lim_{h \rightarrow 0} \frac{(k-1)(-1)^{k-1} \sin h\pi}{-h} = \lim_{h \rightarrow 0} (k-1)(-1)^{k-1} \frac{\sin h\pi}{h\pi} \times (-\pi) = (k-1)(-1)^k \pi = (-1)^k(k-1)\pi.$$

Example: 7 The function $f(x) = |x| + |x-1|$ is

- (a) Continuous at $x=1$, but not differentiable
 (b) Both continuous and differentiable at $x=1$
 (c) Not continuous at $x=1$
 (d) None of these

Solution: (a) We have, $f(x) = |x| + |x-1| = \begin{cases} -2x+1, & x < 0 \\ 1, & 0 \leq x < 1 \\ 2x-1, & x \geq 1 \end{cases}$

Since, $\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} 1 = 1$, $\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (2x-1) = 1$ and $f(1) = 2 \times 1 - 1 = 1$


$\therefore \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x) = f(1)$. So, $f(x)$ is continuous at $x=1$.

Now, $\lim_{x \rightarrow 1^-} \frac{f(x) - f(1)}{x - 1} = \lim_{h \rightarrow 0} \frac{f(1-h) - f(1)}{-h} = \lim_{h \rightarrow 0} \frac{1-1}{-h} = 0$,

and $\lim_{x \rightarrow 1^+} \frac{f(x) - f(1)}{x - 1} = \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h} = \lim_{h \rightarrow 0} \frac{2(1+h) - 1 - 1}{h} = 2$.

\therefore (LHD at $x=1$) \neq (RHD at $x=1$). So, $f(x)$ is not differentiable at $x=1$.

Trick : The graph of $f(x) = |x| + |x-1|$ i.e. $f(x) = \begin{cases} -2x+1, & x < 0 \\ 1, & 0 \leq x < 1 \\ 2x-1, & x \geq 1 \end{cases}$ is



By graph, it is clear that the function is not differentiable at $x=0, 1$ as there it has sharp edges.

Example: 8 Let $f(x) = |x-1| + |x+1|$, then the function is

- (a) Continuous
 (b) Differentiable except $x = \pm 1$
 (c) Both (a) and (b)
 (d) None of these

Solution: (c) Here $f(x) = |x-1| + |x+1| \Rightarrow f(x) = \begin{cases} 2x, & \text{when } x > 1 \\ 2, & \text{when } -1 \leq x \leq 1 \\ -2x, & \text{when } x < -1 \end{cases}$

Graphical solution : The graph of the function is shown alongside,


From the graph it is clear that the function is continuous at all real x , also differentiable at all real x except at $x = \pm 1$. Since sharp edges at $x = -1$ and $x = 1$.

At $x=1$ we see that the slope from the right i.e., R.H.D. = 2, while slope from the left i.e., L.H.D. = 0

Similarly, at $x=-1$ it is clear that R.H.D. = 0 while L.H.D. = -2

Trick : In this method, first of all, we differentiate the function and on the derivative equality sign should be removed from doubtful points.

Here, $f'(x) = \begin{cases} -2, & x < -1 \\ 0, & -1 < x < 1 \text{ (No equality on } -1 \text{ and } +1) \\ 2, & x > 1 \end{cases}$



Now, at $x=1$, $f(1^+)=2$ while $f(1^-)=0$ and

at $x=-1$, $f(-1^+)=0$ while $f(-1^-)=-2$

Thus, $f(x)$ is not differentiable at $x=\pm 1$.

Note : \square This method is not applicable when function is discontinuous.

Example: 9 If the derivative of the function $f(x) = \begin{cases} ax^2 + b, & x < -1 \\ bx^2 + ax + 4, & x \geq -1 \end{cases}$ is everywhere continuous and differentiable at $x = 1$ then

- (a) $a = 2, b = 3$ (b) $a = 3, b = 2$ (c) $a = -2, b = -3$ (d) $a = -3, b = -2$

Solution: (a) $f(x) = \begin{cases} ax^2 + b, & x < -1 \\ bx^2 + ax + 4, & x \geq -1 \end{cases}$

$$\therefore f'(x) = \begin{cases} 2ax, & x < -1 \\ 2bx + a, & x \geq -1 \end{cases}$$

To find a, b we must have two equations in a, b

Since $f(x)$ is differentiable, it must be continuous at $x = -1$.

$$\therefore R = L = V \text{ at } x = -1 \text{ for } f(x) \Rightarrow b - a + 4 = a + b$$

$$\therefore 2a = 4 \text{ i.e., } a = 2$$

Again $f(x)$ is continuous, it must be continuous at $x = -1$.

$$\therefore R = L = V \text{ at } x = -1 \text{ for } f(x)$$

$$-2b + a = -2a \text{ Putting } a = 2, \text{ we get } -2b + 2 = -4$$

$$\therefore 2b = 6 \text{ or } b = 3.$$

Example: 10 Let f be twice differentiable function such that $f'(x) = -f(x)$ and $f(x) = g(x)$, $h(x) = \{f(x)\}^2 + \{g(x)\}^2$. If $h(5) = 11$, then $h(10)$ is equal to

- (a) 22 (b) 11 (c) 0 (d) None of these

Solution: (b) Differentiating the given relation $h(x) = [f(x)]^2 + [g(x)]^2$ w.r.t x , we get
 $h'(x) = 2f(x)f'(x) + 2g(x)g'(x)$ (i)

But we are given $f'(x) = -f(x)$ and $f(x) = g(x)$ so that $f'(x) = g'(x)$.

Then (1) may be re-written as $h'(x) = -2f'(x)f(x) + 2f(x)f'(x) = 0$. Thus $h'(x) = 0$

Whence by integrating, we get $h(x) = \text{constant} = c$ (say). Hence $h(x) = c$ for all x .

In particular, $h(5) = c$ But we are given $h(5) = 11$.

It follows that $c = 11$ and we have $h(x) = 11$ for all x . Therefore, $h(10) = 11$.

Example: 11 The function $f(x) = \begin{cases} |2x - 3|, & x \geq 1 \\ \sin\left(\frac{\pi x}{2}\right), & x < 1 \end{cases}$

- (a) Is continuous at $x = 2$ (b) Is differentiable at $x = 1$
 (c) Is continuous but not differentiable at $x = 1$ (d) None of these

Solution: (c) $[2 + h] = 2, [2 - h] = 1, [1 + h] = 1, [1 - h] = 0$

At $x = 2$, we will check $R = L = V$

$$R = \lim_{h \rightarrow 0} |4 + 2h - 3| [2 + h] = 2, V = 1.2 = 2$$

$$L = \lim_{h \rightarrow 0} |4 - 2h - 3| [2 - h] = 1, R \neq L, \therefore \text{not continuous}$$

At $x = 1$, $R = \lim_{h \rightarrow 0} |2 + 2h - 3| [1 + h] = 1.1 = 1$,

$$V = |-1| [1] = 1$$

$$L = \lim_{h \rightarrow 0} \sin \frac{\pi}{2} (1 - h) = 1$$

Since $R = L = V \therefore$ continuous at $x = 1$.

$$\text{R.H.D.} = \lim_{h \rightarrow 0} \frac{|2 + 2h - 3| [1 + h] - 1}{h} = \lim_{h \rightarrow 0} \frac{|-1| \cdot 1 - 1}{h} = \lim_{h \rightarrow 0} \frac{1 - 1}{h} = 0$$

$$\text{L.H.D.} = \lim_{h \rightarrow 0} \frac{|2 - 2h - 3| [1 - h] - 1}{-h} = \lim_{h \rightarrow 0} \frac{1.0 - 1}{-h} = \lim_{h \rightarrow 0} \frac{1}{h} = \infty$$

Since $\text{R.H.D.} \neq \text{L.H.D.} \therefore$ not differentiable. at $x = 1$.

2.4.2 DIFFERENTIABILITY IN AN OPEN INTERVAL

A function $f(x)$ defined in an open interval (a, b) is said to be differentiable or derivable in open interval (a, b) if it is differentiable at each point of (a, b) .

Differentiability in a closed interval : A function $f: [a, b] \rightarrow \mathbf{R}$ is said to be differentiable in $[a, b]$ if

- (1) $f(x)$ exists for every x such that $a < x < b$ i.e. f is differentiable in (a, b) .
- (2) Right hand derivative of f at $x = a$ exists.
- (3) Left hand derivative of f at $x = b$ exists.

Everywhere differentiable function : If a function is differentiable at each $x \in \mathbf{R}$, then it is said to be everywhere differentiable. e.g., A constant function, a polynomial function, $\sin x, \cos x$ etc. are everywhere differentiable.

Some standard results on differentiability

- (1) Every polynomial function is differentiable at each $x \in \mathbf{R}$.
- (2) The exponential function $a^x, a > 0$ is differentiable at each $x \in \mathbf{R}$.
- (3) Every constant function is differentiable at each $x \in \mathbf{R}$.
- (4) The logarithmic function is differentiable at each point in its domain.
- (5) Trigonometric and inverse trigonometric functions are differentiable in their domains.
- (6) The sum, difference, product and quotient of two differentiable functions is differentiable.
- (7) The composition of differentiable function is a differentiable function.

Important Tips

- ☞ If f is derivable in the open interval (a, b) and also at the end points 'a' and 'b', then f is said to be derivable in the closed interval $[a, b]$.
- ☞ A function f is said to be a differentiable function if it is differentiable at every point of its domain.
- ☞ If a function is differentiable at a point, then it is continuous also at that point.
i.e. Differentiability \Rightarrow Continuity, but the converse need not be true.
- ☞ If a function ' f ' is not differentiable but is continuous at $x = a$, it geometrically implies a sharp corner or kink at $x = a$.
- ☞ If $f(x)$ is differentiable at $x = a$ and $g(x)$ is not differentiable at $x = a$, then the product function $f(x).g(x)$ can still be differentiable at $x = a$.
- ☞ If $f(x)$ and $g(x)$ both are not differentiable at $x = a$ then the product function $f(x).g(x)$ can still be differentiable at $x = a$.
- ☞ If $f(x)$ is differentiable at $x = a$ and $g(x)$ is not differentiable at $x = a$ then the sum function $f(x) + g(x)$ is also not differentiable at $x = a$.
- ☞ If $f(x)$ and $g(x)$ both are not differentiable at $x = a$, then the sum function may be a differentiable function.

Example: 12 The set of points where the function $f(x) = \sqrt{1 - e^{-x^2}}$ is differentiable

- (a) $(-\infty, \infty)$ (b) $(-\infty, 0) \cup (0, \infty)$ (c) $(-1, \infty)$ (d) None of these

Solution: (b) Clearly, $f(x)$ is differentiable for all non-zero values of x , For $x \neq 0$, we have

$$f(x) = \frac{xe^{-x^2}}{\sqrt{1 - e^{-x^2}}}$$

Now, (L.H.D. at $x = 0$)

$$= \lim_{x \rightarrow 0^-} \frac{f(x) - f(0)}{x - 0} = \lim_{h \rightarrow 0} \frac{f(0 - h) - f(0)}{-h} = \lim_{h \rightarrow 0} \frac{\sqrt{1 - e^{-h^2}}}{-h} = \lim_{h \rightarrow 0} \frac{\sqrt{1 - e^{-h^2}}}{h} = - \lim_{h \rightarrow 0} \sqrt{\frac{e^{h^2} - 1}{h^2}} \times \frac{1}{\sqrt{e^{h^2}}} = -1$$

$$\text{and, (RHD at } x = 0) = \lim_{x \rightarrow 0^+} \frac{f(x) - f(0)}{x - 0} = \lim_{h \rightarrow 0} \frac{\sqrt{1 - e^{-h^2}} - 0}{h} = \lim_{h \rightarrow 0} \sqrt{\frac{e^{h^2} - 1}{h^2}} \times \frac{1}{\sqrt{e^{h^2}}} = 1.$$

So, $f(x)$ is not differentiable at $x = 0$, Hence, the points of differentiability of $f(x)$ are $(-\infty, 0) \cup (0, \infty)$.

Example: 13 The function $f(x) = e^{-|x|}$ is

- (a) Continuous everywhere but not differentiable at $x = 0$
 (b) Continuous and differentiable everywhere
 (c) Not continuous at $x = 0$
 (d) None of these

Solution: (a) We have, $f(x) = \begin{cases} e^{-x}, & x \geq 0 \\ e^x, & x < 0 \end{cases}$

Clearly, $f(x)$ is continuous and differentiable for all non-zero x .

$$\text{Now, } \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} e^x = 1 \text{ and } \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} e^{-x} = 1$$

Also, $f(0) = e^0 = 1$


So, $f(x)$ is continuous for all x .

$$(\text{LHD at } x=0) = \left(\frac{d}{dx}(e^x) \right)_{x=0} = [e^x]_{x=0} = e^0 = 1$$

$$(\text{RHD at } x=0) = \left(\frac{d}{dx}(e^{-x}) \right)_{x=0} = [-e^{-x}]_{x=0} = -1$$

So, $f(x)$ is not differentiable at $x=0$.

Hence, $f(x) = e^{-|x|}$ is everywhere continuous but not differentiable at $x=0$. This fact is also evident from the graph of the function.



Example: 14 If $f(x) = \sqrt{1 - \sqrt{1 - x^2}}$, then $f(x)$ is

- (a) Continuous on $[-1, 1]$ and differentiable on $(-1, 1)$
- (b) Continuous on $[-1, 1]$ and differentiable on $(-1, 0) \cup (0, 1)$
- (c) Continuous and differentiable on $[-1, 1]$
- (d) None of these

Solution: (b) We have, $f(x) = \sqrt{1 - \sqrt{1 - x^2}}$. The domain of definition of $f(x)$ is $[-1, 1]$.

$$\text{For } x \neq 0, x \neq 1, x \neq -1 \text{ we have } f(x) = \frac{1}{\sqrt{1 - \sqrt{1 - x^2}}} \times \frac{x}{\sqrt{1 - x^2}}$$

Since $f(x)$ is not defined on the right side of $x=1$ and on the left side of $x=-1$. Also, $f(x) \rightarrow \infty$ when $x \rightarrow -1^+$ or $x \rightarrow 1^-$. So, we check the differentiability at $x=0$.

$$\begin{aligned} \text{Now, } (\text{LHD at } x=0) &= \lim_{x \rightarrow 0} \frac{f(x) - f(0)}{x - 0} = \lim_{h \rightarrow 0} \frac{f(0 - h) - f(0)}{-h} \\ &= \lim_{h \rightarrow 0} \frac{\sqrt{1 - \sqrt{1 - h^2}} - 0}{-h} = - \lim_{h \rightarrow 0} \frac{\sqrt{1 - \{1 - (1/2)h^2 + (3/8)h^4 + \dots\}}}{h} = - \lim_{h \rightarrow 0} \sqrt{\frac{1}{2} - \frac{3}{8}h^2 + \dots} = - \frac{1}{\sqrt{2}} \end{aligned}$$

$$\text{Similarly, } (\text{RHD at } x=0) = \frac{1}{\sqrt{2}}$$

Hence, $f(x)$ is not differentiable at $x=0$.

Example: 15 Let $f(x)$ be a function differentiable at $x=c$. Then $\lim_{x \rightarrow c} f(x)$ equals

- (a) $f(c)$
- (b) $f'(c)$
- (c) $\frac{1}{f(c)}$
- (d) None of these

Solution: (d) Since $f(x)$ is differentiable at $x=c$, therefore it is continuous at $x=c$. Hence, $\lim_{x \rightarrow c} f(x) = f(c)$.

Example: 16 The function $f(x) = (x^2 - 1)|x^2 - 3x + 2| + \cos(|x|)$ is not differentiable at

- (a) -1
- (b) 0
- (c) 1
- (d) 2

Solution: (d) $(x^2 - 3x + 2) = (x - 1)(x - 2) = +ive$

When $x < 1$ or > 2 , $-ive$ when $1 \leq x \leq 2$

Also $\cos|x| = \cos x$ (since $\cos(-x) = \cos x$)

$$\therefore f(x) = -(x^2 - 1)(x^2 - 3x + 2) + \cos x, \quad 1 \leq x \leq 2$$

$$\therefore f(x) = (x^2 - 1)(x^2 - 3x + 2) + \cos x, \quad x > 2 \quad \dots\dots\dots(i)$$

Evidently $f(x)$ is not differentiable at $x=2$ as $L' \neq R'$

Note : \square For all other values like $x < 0, 0 \leq x < 1$, $f(x)$ is same as given by (i).

Example: 17 If $f(x) = \begin{cases} xe^{-\left(\frac{1}{|x|} + \frac{1}{x}\right)}, & x \neq 0 \\ 0, & x = 0 \end{cases}$, then $f(x)$ is

- (a) Continuous as well as differentiable for all x
- (b) Continuous for all x but not differentiable at $x = 0$
- (c) Neither differentiable nor continuous at $x = 0$
- (d) Discontinuous every where

Solution: (b) $f(0) = 0$ and $f(x) = xe^{-\left(\frac{1}{|x|} + \frac{1}{x}\right)}$

$$\text{R.H.L.} = \lim_{h \rightarrow 0} (0 + h)e^{-2/h} = \lim_{h \rightarrow 0} \frac{h}{e^{2/h}} = 0$$

$$\text{L.H.L.} = \lim_{h \rightarrow 0} (0 - h)e^{-\left(\frac{1}{h} + \frac{1}{h}\right)} = 0$$

$\therefore f(x)$ is continuous.

$$Rf'(x) \text{ at } (x=0) = \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{he^{-2/h}}{h} = e^{-\infty} = 0$$

$$Lf'(x) \text{ at } (x=0) = \lim_{h \rightarrow 0} \frac{f(0-h) - f(0)}{-h} = \lim_{h \rightarrow 0} \frac{-he^{-\left(\frac{1}{h} + \frac{1}{h}\right)}}{-h} = +1 \Rightarrow Lf'(x) \neq Rf'(x)$$

$f(x)$ is not differentiable at $x = 0$.

Example: 18 The function $f(x) = x^2 \sin \frac{1}{x}$, $x \neq 0$, $f(0) = 0$ at $x = 0$

- (a) Is continuous but not differentiable
- (b) Is discontinuous
- (c) Is having continuous derivative
- (d) Is continuous and differentiable

Solution: (d) $\lim_{x \rightarrow 0} f(x) = x^2 \sin \left(\frac{1}{x}\right)$ but $-1 \leq \sin \left(\frac{1}{x}\right) \leq 1$ and $x \rightarrow 0$

$$\therefore \lim_{x \rightarrow 0^+} f(x) = 0 = \lim_{x \rightarrow 0^-} f(x) = f(0)$$

Therefore $f(x)$ is continuous at $x = 0$. Also, the function $f(x) = x^2 \sin \frac{1}{x}$ is differentiable

$$\text{because } Rf'(x) = \lim_{h \rightarrow 0} \frac{h^2 \sin \frac{1}{h} - 0}{h} = 0, Lf'(x) = \lim_{h \rightarrow 0} \frac{h^2 \sin \left(\frac{1}{-h}\right)}{-h} = 0.$$

Example: 19 Which of the following is not true

- (a) A polynomial function is always continuous
- (b) A continuous function is always differentiable
- (c) A differentiable function is always continuous
- (d) e^x is continuous for all x

Solution: (b) A continuous function may or may not be differentiable. So (b) is not true.

Example: 20 If $f(x) = \text{sgn}(x^3)$, then

- (a) f is continuous but not derivable at $x = 0$
- (b) $f(0^+) = 2$
- (c) $f(0^-) = 1$
- (d) f is not derivable at $x = 0$

$$\text{Solution: (d) Here, } f(x) = \text{sgn } x^3 = \begin{cases} \frac{x^3}{|x^3|} & \text{for } x^3 \neq 0 \\ 0 & \text{for } x^3 = 0 \end{cases} \quad \text{Thus, } f(x) = \text{sgn } x^3 = \text{sgn } x, \text{ which is neither continuous}$$

$$\begin{cases} \frac{x}{|x|} & \text{for } x \neq 0 \\ 0 & \text{for } x = 0 \end{cases}$$

$$\begin{cases} -1, & x < 0 \\ 0, & x = 0 \\ 1, & x > 0 \end{cases}$$

nor derivable at 0.

$$\text{Note : } \square \quad f(0^+) = \lim_{h \rightarrow 0^+} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0^+} \frac{1-0}{h} \rightarrow \infty \quad \text{and} \quad f(0^-) = \lim_{h \rightarrow 0^-} \frac{f(0-h) - f(0)}{h} = \lim_{h \rightarrow 0^-} \frac{-1-0}{h} \rightarrow \infty.$$

$\therefore f(0^+) \neq f(0^-)$, $\therefore f$ is not derivable at $x=0$.

Example: 21 A function $f(x) = \begin{cases} 1+x, & x \leq 2 \\ 5-x, & x > 2 \end{cases}$ is

- (a) Not continuous at $x=2$ (b) Differentiable at $x=2$
 (c) Continuous but not differentiable at $x=2$ (d) None of the above

Solution: (c) $\lim_{h \rightarrow 0^-} 1+(2-h) = 3$, $\lim_{h \rightarrow 0^+} 5-(2+h) = 3$, $f(2) = 3$

Hence, f is continuous at $x=2$

$$\text{Now } Rf(x) = \lim_{h \rightarrow 0} \frac{5-(2+h)-3}{h} = -1$$

$$Lf(x) = \lim_{h \rightarrow 0} \frac{1+(2-h)-3}{-h} = 1$$

$$\ominus Rf(x) \neq Lf(x)$$


$\therefore f$ is not differentiable at $x=2$.

Example: 22 Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a function. Define $g: \mathbb{R} \rightarrow \mathbb{R}$ by $g(x) = |f(x)|$ for all x . Then g is

- (a) Onto if f is onto (b) One-one if f is one-one
 (c) Continuous if f is continuous (d) Differentiable if f is differentiable

Solution: (c) $g(x) = |f(x)| \geq 0$. So $g(x)$ cannot be onto. If $f(x)$ is one-one and $f(x_1) = -f(x_2)$ then $g(x_1) = g(x_2)$.

So, ' $f(x)$ is one-one' does not ensure that $g(x)$ is one-one.



If $f(x)$ is continuous for $x \in \mathbb{R}$ $|f(x)|$ is also continuous for $x \in \mathbb{R}$. This is obvious from the following graphical consideration.

So the answer (c) is correct. The fourth answer (d) is not correct from the above graphs $y = f(x)$ is differentiable at P while $y = |f(x)|$ has two tangents at P , i.e. not differentiable at P .

ASSIGNMENT

1. If $f(x) = \begin{cases} 1 & , x < 0 \\ 1 + \sin x & , 0 \leq x \leq \pi/2 \end{cases}$ then at $x = 0$, the value of $f'(x)$ is equal to
 (a) 1 (b) 0 (c) ∞ (d) Derivative does not exist
2. If $f(x) = |x - 3|$, then $f'(3)$ equals
 (a) 0 (b) 1 (c) -1 (d) Does not exist
3. If $f(x) = \begin{cases} x \sin(1/x), & x \neq 0 \\ 0 & , x = 0 \end{cases}$ then at $x = 0$, the function is
 (a) Discontinuous (b) Continuous but not differentiable
 (c) Both continuous and differentiable (d) None of these
4. If $f(x) = |x - 3|$, then f is
 (a) Discontinuous at $x = 2$ (b) Not differentiable at $x = 2$
 (c) Differentiable at $x = 3$ (d) Continuous but not differentiable at $x = 3$
5. If $f(x) = \begin{cases} x+1, & \text{when } x < 2 \\ 2x-1, & \text{when } x \geq 2 \end{cases}$, then $f'(x)$ at $x = 2$ equals
 (a) 0 (b) 1 (c) 2 (d) Does not exist
6. If $f(x) = \begin{cases} x^2 \sin(1/x), & \text{when } x \neq 0 \\ 0 & , \text{when } x = 0 \end{cases}$, then at $x = 0$, value of $f'(x)$ equals
 (a) 1 (b) 0 (c) ∞ (d) Does not exist
7. If $\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}$ exists finitely, then
 (a) $\lim_{x \rightarrow c} f(x) = f(c)$ (b) $\lim_{x \rightarrow c} f(x) = f(c)$ (c) $\lim_{x \rightarrow c} f(x)$ does not exist (d) $\lim_{x \rightarrow c} f(x)$ may or may not exist
8. If $f(x) = \frac{|x-1|}{x-1}$, $x \neq 1$ and $f(1) = 1$. Then which of the following statement is true
 (a) Continuous for $x \leq 1$ (b) Discontinuous at $x = 1$
 (c) Differentiable at $x = 1$ (d) Discontinuous for $x > 1$
9. Let $f(xy) = f(x)f(y)$ for all $x, y \in \mathbb{R}$ If $f(1) = 2$ and $f(4) = 4$, then $f'(4)$ equal to
 (a) 4 (b) 1 (c) $\frac{1}{2}$ (d) 2
10. The derivative of $f(x) = |x|$ at $x = 0$ is
 (a) 1 (b) 0 (c) -1 (d) Does not exist

11. If $f(x) = \begin{cases} e^x + ax, & x < 0 \\ b(x-1)^2, & x \geq 0 \end{cases}$ is differentiable at $x=0$ then (a, b) is
 (a) $(-3, -1)$ (b) $(-3, 1)$ (c) $(3, 1)$ (d) $(3, -1)$
12. At the point $x = 1$, the function $f(x) = \begin{cases} x^3 - 1; & 1 < x < \infty \\ x - 1; & -\infty < x \leq 1 \end{cases}$
 (a) Continuous and differentiable (b) Continuous and not differentiable
 (c) Discontinuous and differentiable (d) Discontinuous and not differentiable
13. The function $|x^3|$ is
 (a) Differentiable everywhere (b) Continuous but not differentiable at $x=0$
 (c) Not a continuous function (d) A function with range $[0, \infty]$
14. For the function $f(x) = |x^2 - 5x + 6|$ the derivative from the right $f'(2+)$; and the derivative from left $f'(2-)$ are respectively
 (a) $1, -1$ (b) $-1, 1$ (c) $0, 2$ (d) None of these
15. Let $f(x)$ be an even function. Then $f(x)$
 (a) Is an even function (b) Is an odd function (c) May be even or odd (d) None of these
16. Let $f(x)$ be an odd function. Then $f(x)$
 (a) Is an even function (b) Is an odd function (c) May be even or odd (d) None of these
17. Let $g(x)$ be the inverse of the function $f(x)$ and $f(x) = \frac{1}{1+x^3}$. Then $g'(x)$ is equal to
 (a) $\frac{1}{1+(g(x))^3}$ (b) $\frac{1}{1+(f(x))^3}$ (c) $1+(g(x))^3$ (d) $1+(f(x))^3$
18. Let $g(x)$ be the inverse of an invertible function $f(x)$ which is differentiable at $x=c$, then $g'(f(c))$ equals
 (a) $f'(c)$ (b) $\frac{1}{f'(c)}$ (c) $f(c)$ (d) None of these
19. If $f(x) = \begin{cases} x+2, & -1 < x < 3 \\ 5, & x = 3 \\ 8-x, & x > 3 \end{cases}$ then at $x=3$, $f(x) =$
 (a) 1 (b) -1 (c) 0 (d) Does not exist
20. If $f(x) = (x - x_0)g(x)$, where $g(x)$ is continuous at x_0 , then $f'(x_0)$ is equal to
 (a) 0 (b) x_0 (c) $g(x_0)$ (d) None of these
21. Function $f(x) = |x| + |x-1|$ is not differentiable at
 (a) $x=1, -1$ (b) $x=0, -1$ (c) $x=0, 1$ (d) $x=1, 2$
22. If $f(x) = \begin{cases} e^x; & x \leq 0 \\ |1-x|; & x > 0 \end{cases}$, then
 (a) $f(x)$ is differentiable at $x=0$ (b) $f(x)$ is continuous at $x=0$
 (c) $f(x)$ is differentiable at $x=1$ (d) $f(x)$ is continuous at $x=1$
23. The function which is continuous for all real values of x and differentiable at $x=0$, is
 (a) $|x|$ (b) $\log x$ (c) $\sin x$ (d) $x^{1/2}$

Advance Level

24. The number of points at which the function $f(x) = |x - 0.5| + |x - 1| + \tan x$ does not have a derivative in the interval $(0, 2)$ is
- (a) 1 (b) 2 (c) 3 (d) 4
25. If $f(x) = \begin{cases} ax^2 + b, & b \neq 0, x \leq 1 \\ bx^2 + ax + c, & x > 1 \end{cases}$. Then $f(x)$ is continuous and differentiable at $x = 1$ if
- (a) $c = 0, a = 2b$ (b) $a = b, c \in R$ (c) $a = b, c = 0$ (d) $a = b, c \neq 0$
26. If $f(x) = \begin{cases} ax^2 - b, & |x| < 1 \\ \frac{1}{|x|}, & |x| \geq 1 \end{cases}$ is differentiable at $x = 1$, then
- (a) $a = \frac{1}{2}, b = -\frac{1}{2}$ (b) $a = -\frac{1}{2}, b = -\frac{3}{2}$ (c) $a = b = \frac{1}{2}$ (d) $a = b = -\frac{1}{2}$
27. The set of points where the function $f(x) = |x - 1|e^x$ is differentiable is
- (a) R (b) $R - \{1\}$ (c) $R - \{-1\}$ (d) $R - \{0\}$
28. Let $f(x)$ be defined on R such that $f(1) = 2, f(2) = 8$ and $f(u + v) = f(u) + kuv - 2v^2$ for all $u, v \in R$ and $u \neq v$ (k is a fixed constant). Then
- (a) $f(x) = 8x$ (b) $f(x) = 8x$ (c) $f(x) = x$ (d) None of these
29. Let $f: R \rightarrow R$ be a function defined by $f(x) = \max\{x, x^3\}$. The set of all points where $f(x)$ is not differentiable is
- (a) $\{-1, 1\}$ (b) $\{-1, 0\}$ (c) $\{0, 1\}$ (d) $\{-1, 0, 1\}$
30. Let $f(x) = \begin{cases} 0, & x < 0 \\ x^2, & x \geq 0 \end{cases}$ then for all values of x
- (a) f is continuous but not differentiable (b) f is differentiable but not continuous
(c) f' is continuous but not differentiable (d) f' is continuous and differentiable
31. If $u(x) = \sin x$ $\begin{cases} 1 & \text{for } x > 0 \\ 0 & \text{for } x = 0 \\ -1 & \text{for } x < 0 \end{cases}$ then $u(x) \cdot v(x)$ has a derivative at $x = 1$ is
- $v(x) = \text{sgn}(x) = \begin{cases} 1 & \text{for } x > 0 \\ 0 & \text{for } x = 0 \\ -1 & \text{for } x < 0 \end{cases}$
- (a) $\cos 1$ (b) $\sin 1$ (c) Not continuous at $x = 1$ (d) None of these
32. The coefficient a and b that make the function $f(x) = \begin{cases} \frac{1}{|x|} & \text{for } |x| \geq 1 \\ ax^2 + b & \text{for } |x| < 1 \end{cases}$ continuous and differentiable at any point are given by
- (a) $a = -1/2, b = 3/2$ (b) $a = 1/2, b = -3/2$ (c) $a = 1, b = -1$ (d) None of these
33. If $f(x) = \int_{-1}^x |t| dt, x \geq -1$, then
- (a) f and f' are continuous for $x + 1 > 0$ (b) f is continuous but f' is not for $x + 1 > 0$
(c) f and f' are continuous at $x = 0$ (d) f is continuous at $x = 0$ but f' is not so
34. Let $f(x) = \begin{cases} x^n \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$, then $f(x)$ is continuous but not differentiable at $x = 0$ if
- (a) $n \in (0, 1]$ (b) $n \in [1, \infty)$ (c) $n \in (-\infty, \infty)$ (d) $n = 0$

35. Which of the following is differentiable at $x=0$
 (a) $\cos(|x|)+|x|$ (b) $\cos(|x|)-|x|$ (c) $\sin(|x|)+|x|$ (d) $\sin(|x|)-|x|$
36. If $x+4|y|=6y$, then y as a function of x is
 (a) Continuous at $x=0$ (b) Derivable at $x=0$ (c) $\frac{dy}{dx}=\frac{1}{2}$ for all x (d) None of these
37. The set of point where the function $f(x)=x|x|$ is differentiable is
 (a) $(-\infty, \infty)$ (b) $(-\infty, 0) \cup (0, \infty)$ (c) $(0, \infty)$ (d) $[0, \infty)$
38. If $f(x)=\sin^{-1}\left(\frac{2x}{1+x^2}\right)$ then $f(x)$ is differentiable on
 (a) $[-1, 1]$ (b) $\mathbb{R}-\{-1, 1\}$ (c) $\mathbb{R} - (-1, 1)$ (d) None of these
39. Let $f(x)=|x|$ and $g(x)=|x^3|$, then
 (a) $f(x)$ and $g(x)$ both are continuous at $x=0$ (b) $f(x)$ and $g(x)$ both are differentiable at $x=0$
 (c) $f(x)$ is differentiable but $g(x)$ is not differentiable at $x=0$ (d) $f(x)$ and $g(x)$ both are not differentiable at $x=0$
40. The function $f(x)=\sin^{-1}(\cos x)$ is
 (a) Discontinuous at $x=0$ (b) Continuous at $x=0$ (c) Differentiable at $x=0$ (d) None of these
41. Let $f(x)=(x+|x|)|x|$. Then for all x
 (a) f is continuous (b) f is differentiable for some x
 (c) f' is continuous (d) f is continuous
42. The set of all those points, where the function $f(x)=\frac{x}{1+|x|}$ is differentiable, is
 (a) $(-\infty, \infty)$ (b) $[0, \infty)$ (c) $(-\infty, 0) \cup (0, \infty)$ (d) $(0, \infty)$
43. $f(x)$ and $g(x)$ are two differentiable function on $[0, 2]$ such that $f'(x)-g'(x)=0$, $f(1)=2$, $g(1)=4$, $f(2)=3$, $g(2)=9$, then $f(x)-g(x)$ at $x=\frac{3}{2}$ is
 (a) 0 (b) 2 (c) 10 (d) -5
44. The set of points of differentiability of the $f(x)=\begin{cases} \frac{\sqrt{x+1}-1}{\sqrt{x}}, & \text{for } x \neq 0 \\ 0, & \text{for } x=0 \end{cases}$ is
 (a) \mathbb{R} (b) $[0, \infty)$ (c) $(0, \infty)$ (d) $\mathbb{R}-\{0\}$
45. If $f(x)=a|\sin x|+be^{|x|}+c|x|^3$ and if $f(x)$ is differentiable at $x=0$, then
 (a) $a=b=c=0$ (b) $a=0, b=0, c \in \mathbb{R}$ (c) $b=c=0, a \in \mathbb{R}$ (d) $c=0, a=0, b \in \mathbb{R}$
46. If $f(x)=\begin{cases} \frac{x-1}{2x^2-7x+5}, & x \neq 1 \\ -\frac{1}{3}, & x=1 \end{cases}$, then $f(1)$ equals
 (a) $\frac{2}{9}$ (b) $-\frac{2}{9}$ (c) 0 (d) Does not exist
47. Function $f(x)=1+|\sin x|$ is
 (a) Continuous no where (b) Differentiable no where
 (c) Every where continuous (d) Not differentiable at $x=0$

48. Function $f(x) = \begin{cases} x^2, & x \leq 0 \\ 1, & 0 < x \leq 1 \\ 1/x, & x > 1 \end{cases}$ is

- (a) Differentiable at $x=0, 1$ (b) Differentiable only at $x=0$
(c) Differentiable at only $x=1$ (d) Not differentiable at $x=0, 1$

49. Function $f(x) = \begin{cases} -x, & \text{if } x < 0 \\ x^2, & \text{if } 0 \leq x \leq 1 \\ x^3 - x + 1, & \text{if } x > 1 \end{cases}$, is differentiable at

- (a) $x=0$ but not at $x=1$ (b) $x=1$ but not at $x=0$
(c) $x=0$ and $x=1$ (d) Neither $x=0$ nor $x=1$

50. If $g(x) = x f(x)$ where $f(x) = \begin{cases} x \sin\left(\frac{1}{x}\right), & x \neq 0 \\ 0, & x = 0 \end{cases}$ then at $x=0$

- (a) g is differentiable but g' is discontinuous function
(b) Both f and g are differentiable
(c) g is differentiable and g' is continuous function (d) None of these

51. The set of points where $f(x) = x|x|$ is differentiable two times is

- (a) R_0 (b) R_+ (c) R (d) None of these

52. If $f(x) = \begin{cases} \frac{x(3e^{1/x} + 4)}{2 - e^{1/x}}, & x \neq 0 \\ 0, & x = 0 \end{cases}$, then

- (a) $\lim_{x \rightarrow 0} f(x) = 1$ (b) $f(x)$ is continuous at $x=0$
(c) $f(x)$ is differentiable at $x=0$ (d) $f(0+0) = 3$

ANSWER

ASSIGNMENT (BASIC & ADVANCE LEVEL)

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
d	d	b	d	d	b	a	b	d	d	b	b	a,d	a	b	a	c	b	d	c
21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
c	b,d	c	c	a	b	b	d	d	c	a	a	a	a	d	a	a	b	a	b
41	42	43	44	45	46	47	48	49	50	51	52								