

ELLIPSE

5.2.1 DEFINITION

An ellipse is the locus of a point which moves in such a way that its distance from a fixed point is in constant ratio (<1) to its distance from a fixed line. The fixed point is called the **focus** and fixed line is called the **directrix** and the constant ratio is called the **eccentricity** of the ellipse, denoted by (e).

In other words, we can say an ellipse is the locus of a point which moves in a plane so that the sum of its distances from two fixed points is constant and is more than the distance between the two fixed points.


Let $S(\alpha, \beta)$ is the focus, ZZ is the directrix and P is any point on the ellipse. Then by definition,

$$\frac{SP}{PM} = e \Rightarrow SP = e \cdot PM$$

$$\sqrt{(x-\alpha)^2 + (y-\beta)^2} = e \frac{Ax + By + C}{\sqrt{A^2 + B^2}}$$

Squaring both sides, $(A^2 + B^2)[(x-\alpha)^2 + (y-\beta)^2] = e^2(Ax + By + C)^2$

Note : \square The condition for second degree equation in x and y to represent an ellipse is that $h^2 - ab < 0$ and $\Delta = abc + 2fgh - af^2 - bg^2 - cf^2 \neq 0$



Example: 1 The equation of an ellipse whose focus is $(-1, 1)$, whose directrix is $x - y + 3 = 0$ and whose eccentricity is $\frac{1}{2}$, is given by

(a) $7x^2 + 2xy + 7y^2 + 10x - 10y + 7 = 0$

(b) $7x^2 - 2xy + 7y^2 - 10x + 10y + 7 = 0$

(c) $7x^2 - 2xy + 7y^2 - 10x - 10y - 7 = 0$

(d) $7x^2 - 2xy + 7y^2 + 10x + 10y - 7 = 0$

Solution: (a) Let any point on it be (x, y) then by definition,

$$\sqrt{(x+1)^2 + (y-1)^2} = \frac{1}{2} \left| \frac{x-y+3}{\sqrt{1^2+1^2}} \right|$$

Squaring and simplifying, we get

$$7x^2 + 2xy + 7y^2 + 10x - 10y + 7 = 0, \text{ which is the required ellipse.}$$

5.2.2 STANDARD EQUATION OF THE ELLIPSE


Let S be the focus, ZM be the directrix of the ellipse and $P(x, y)$ is any point on the ellipse, then by

definition $\frac{SP}{PM} = e$

$$\Rightarrow (SP)^2 = e^2 (PM)^2$$


$$(x - ae)^2 + (y - 0)^2 = e^2 \left(\frac{a}{e} - x \right)^2 \Rightarrow \frac{x^2}{a^2} + \frac{y^2}{a^2(1-e^2)} = 1$$

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, \text{ where } b^2 = a^2(1-e^2)$$



Since $e < 1$, therefore $a^2(1 - e^2) < a^2 \Rightarrow b^2 < a^2$. Some terms related to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, a > b$:

- (1) **Centre:** The point which bisects each chord of the ellipse passing through it, is called centre $(0, 0)$ denoted by C .



- (2) **Major and minor axes:** The diameter through the foci, is called the major axis and the diameter bisecting it at right angles is called the minor axis. The major and minor axes are together called principal axes.

Length of the major axis $AA' = 2a$, Length of the minor axis $BB' = 2b$

The ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, is symmetrical about both the axes.

- (3) **Vertices:** The extremities of the major axis of an ellipse are called vertices. The coordinates of vertices A and A' are $(a, 0)$ and $(-a, 0)$ respectively.

- (4) **Foci:** S and S' are two foci of the ellipse and their coordinates are $(ae, 0)$ and $(-ae, 0)$ respectively. Distance between foci $SS' = 2ae$.

- (5) **Directrices:** ZM and $Z'M$ are two directrices of the ellipse and their equations are $x = \frac{a}{e}$ and $x = -\frac{a}{e}$ respectively. Distance between directrices $ZZ' = \frac{2a}{e}$.

- (6) **Eccentricity of the ellipse:** For the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$,

$$\text{we have } b^2 = a^2(1 - e^2) \Rightarrow e^2 = 1 - \frac{b^2}{a^2} = 1 - \frac{4b^2}{4a^2} = 1 - \left(\frac{2b}{2a}\right)^2; \quad e = \sqrt{1 - \left(\frac{\text{Minoraxis}}{\text{Majoraxis}}\right)^2}$$

This formula gives the eccentricity of the ellipse.

- (7) **Ordinate and double ordinate:** Let P be a point on the ellipse and let PN be perpendicular to the major axis AA' such that PN produced meets the ellipse at P' . Then PN is called the ordinate of P and PNP' the double ordinate of P .

If abscissa of P is h , then ordinate of P , $\frac{y^2}{b^2} = 1 - \frac{h^2}{a^2} \Rightarrow y = \frac{b}{a}\sqrt{(a^2 - h^2)}$ (For first quadrant)

And ordinate of P' is $y = -\frac{b}{a}\sqrt{(a^2 - h^2)}$ (For fourth quadrant)

Hence coordinates of P and P' are $\left(h, \frac{b}{a}\sqrt{(a^2 - h^2)}\right)$ and $\left(h, -\frac{b}{a}\sqrt{(a^2 - h^2)}\right)$ respectively.

- (8) **Latus-rectum:** Chord through the focus and perpendicular to the major axis is called its latus rectum.

The double ordinates LL' and L_1L_1' are latus rectum of the ellipse.

Length of latus rectum $LL' = L_1L'_1 = \frac{2b^2}{a}$ and end points of latus-rectum are $L = \left(ae, \frac{b^2}{a} \right), L' = \left(ae, -\frac{b^2}{a} \right)$ and $L_1 = \left(-ae, \frac{b^2}{a} \right); L'_1 = \left(-ae, -\frac{b^2}{a} \right)$


(9) **Focal chord:** A chord of the ellipse passing through its focus is called a focal chord.

(10) **Focal distances of a point:** The distance of a point from the focus is its focal distance. The sum of the focal distances of any point on an ellipse is constant and equal to the length of the major axis of the ellipse.

Let $P(x_1, y_1)$ be any point on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

$$SP = ePM = e \left(\frac{a}{e} - x_1 \right) = a - ex_1 \text{ and } S'P = ePM = e \left(\frac{a}{e} + x_1 \right) = a + ex_1$$

$$\therefore SP + S'P = (a - ex_1) + (a + ex_1) = 2a = AA' = \text{major axis.}$$



Example: 2 The length of the latus-rectum of the ellipse $5x^2 + 9y^2 = 45$ is

- (a) $\sqrt{5}/4$ (b) $\sqrt{5}/2$ (c) $5/3$ (d) $10/3$

Solution: (d) Here the ellipse is $\frac{x^2}{9} + \frac{y^2}{5} = 1$

$$\text{Here } a^2 = 9 \text{ and } b^2 = 5. \text{ So, latus-rectum} = \frac{2b^2}{a} = \frac{2(5)}{3} = \frac{10}{3}.$$

Example: 3 In an ellipse the distance between its foci is 6 and its minor axis is 8. Then its eccentricity is

- (a) $\frac{4}{5}$ (b) $\frac{1}{\sqrt{52}}$ (c) $\frac{3}{5}$ (d) $\frac{1}{2}$

Solution: (c) Distance between foci = 6 $\Rightarrow 2ae = 6 \Rightarrow ae = 3$, Minor axis = 8 $\Rightarrow 2b = 8 \Rightarrow b = 4 \Rightarrow b^2 = 16$

$$\text{From } b^2 = a^2(1 - e^2), \Rightarrow 16 = a^2 - a^2e^2 \Rightarrow a^2 - 9 = 16 \Rightarrow a = 5$$

$$\text{Hence, } ae = 3 \Rightarrow e = \frac{3}{5}$$

Example: 4 What is the equation of the ellipse with foci $(\pm 2, 0)$ and eccentricity $\frac{1}{2}$

- (a) $3x^2 + 4y^2 = 48$ (b) $4x^2 + 3y^2 = 48$ (c) $3x^2 + 4y^2 = 0$ (d) $4x^2 + 3y^2 = 0$

Solution: (a) Here $ae = \pm 2$ $\ominus e = \frac{1}{2}$, $\therefore a = \pm 4$

$$\text{Form } b^2 = a^2(1 - e^2) \Rightarrow b^2 = 16 \left(1 - \frac{1}{4} \right) \Rightarrow b^2 = 12$$

$$\text{Hence, the equation of ellipse is } \frac{x^2}{16} + \frac{y^2}{12} = 1 \text{ or } 3x^2 + 4y^2 = 48$$

Example: 5 If $P(x, y), F_1 = (3, 0), F_2 = (-3, 0)$ and $16x^2 + 25y^2 = 400$, then $PF_1 + PF_2$ equals

- (a) 8 (b) 6 (c) 10 (d) 12

Solution: (c) We have $16x^2 + 25y^2 = 400 \Rightarrow \frac{x^2}{25} + \frac{y^2}{16} = 1$ or $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, where $a^2 = 25$ and $b^2 = 16$

$$\text{This equation represents an ellipse with eccentricity given by } e^2 = 1 - \frac{b^2}{a^2} = 1 - \frac{16}{25} = \frac{9}{25}$$

$$\Rightarrow e = 3/5$$

So, the coordinates of the foci are $(\pm ae, 0)$ i.e. $(3, 0)$ and $(-3, 0)$, Thus, F_1 and F_2 are the foci of the ellipse.

Since, the sum of the focal distance of a point on an ellipse is equal to its major axis,

$$\therefore PF_1 + PF_2 = 2a = 10$$

Example: 6 An ellipse has OB as semi minor axis. F and F are its foci and the angle FBF is a right angle. Then the eccentricity of the ellipse is

(a) $\frac{1}{2}$

(b) $\frac{1}{\sqrt{2}}$

(c) $\frac{2}{3}$

(d) $\frac{1}{3}$


Solution: (b) Since $\angle FBF = \frac{\pi}{2}$

$$\therefore \angle FBC = \angle FCB = \frac{\pi}{4}$$

$$\therefore CB = CF \Rightarrow b = ae$$

$$\Rightarrow b^2 = a^2 e^2 \Rightarrow a^2(1 - e^2) = a^2 e^2$$

$$\Rightarrow 1 - e^2 = e^2 \Rightarrow 2e^2 = 1 \Rightarrow e = \frac{1}{\sqrt{2}}$$



Example: 7 Let P be a variable point on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ with foci F_1 and F_2 . If A is the area of the triangle PF_1F_2 , then the maximum value of A is

(a) $2abe$

(b) abe

(c) $\frac{1}{2}abe$

(d) None of these

Solution: (b) Let $P(a \cos \theta, b \sin \theta)$ and $F_1(-ae, 0), F_2(ae, 0)$

$$A = \text{Area of } \triangle PF_1F_2 = \frac{1}{2} \begin{vmatrix} a \cos \theta & b \sin \theta & 1 \\ ae & 0 & 1 \\ -ae & 0 & 1 \end{vmatrix} = \frac{1}{2} |2ae b \sin \theta| = aeb |\sin \theta|$$

$$\therefore A \text{ is maximum, when } |\sin \theta| = 1.$$

$$\text{Hence, maximum value of } A = aeb$$

Example: 8 The eccentricity of an ellipse, with its centre at the origin is $\frac{1}{2}$. If one of the directrices is $x = 4$, then the equation of the ellipse is
[AIEEE 2004]

(a) $4x^2 + 3y^2 = 1$

(b) $3x^2 + 4y^2 = 12$

(c) $4x^2 + 3y^2 = 12$

(d) $3x^2 + 4y^2 = 1$

Solution: (b) Given $e = \frac{1}{2}, \frac{a}{e} = 4$. So, $a = 2 \Rightarrow a^2 = 4$

$$\text{From } b^2 = a^2(1 - e^2) \Rightarrow b^2 = 4\left(1 - \frac{1}{4}\right) = 4 \times \frac{3}{4} = 3$$

Hence the equation of ellipse is


$$\frac{x^2}{4} + \frac{y^2}{3} = 1, \text{ i.e. } 3x^2 + 4y^2 = 12$$

5.2.3 EQUATION OF ELLIPSE IN OTHER FORM

In the equation of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, if $a > b$ or $a^2 > b^2$ (denominator of x^2 is greater than that of y^2), then the major and minor axis lie along x -axis and y -axis respectively. But if $a < b$ or $a^2 < b^2$ (denominator of x^2 is less than that of y^2), then the major axis of the ellipse lies along the y -axis and is of length $2b$ and the minor axis along the x -axis and is of length $2a$.

The coordinates of foci S and S' are $(0, be)$ and $(0, -be)$ respectively.

The equation of the directrices ZK and $Z'K'$ are $y = \pm b/e$ and eccentricity e is given by the formula $a^2 = b^2(1 - e^2)$ or $e = \sqrt{1 - \frac{a^2}{b^2}}$



Difference between both ellipse will be clear from the following table.

| Ellipse | $\left\{ \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \right\}$ | |
|------------------------------------|--|---|
| Basic fundamentals | | |
| | For $a > b$ | For $b > a$ |
| Centre | $(0, 0)$ | $(0, 0)$ |
| Vertices | $(\pm a, 0)$ | $(0, \pm b)$ |
| Length of major axis | $2a$ | $2b$ |
| Length of minor axis | $2b$ | $2a$ |
| Foci | $(\pm ae, 0)$ | $(0, \pm be)$ |
| Equation of directrices | $x = \pm a/e$ | $y = \pm b/e$ |
| Relation in a, b and e | $b^2 = a^2(1 - e^2)$ | $a^2 = b^2(1 - e^2)$ |
| Length of latus rectum | $\frac{2b^2}{a}$ | $\frac{2a^2}{b}$ |
| Ends of latus-rectum | $\left(\pm ae, \pm \frac{b^2}{a} \right)$ | $\left(\pm \frac{a^2}{b}, \pm be \right)$ |
| Parametric equations | $(a \cos \phi, b \sin \phi)$ | $(a \cos \phi, b \sin \phi) \quad (0 \leq \phi < 2\pi)$ |
| Focal radii | $SP = a - ex_1$ and $S'P = a + ex_1$ | $SP = b - ey_1$ and $S'P = b + ey_1$ |
| Sum of focal radii $SP + S'P =$ | $2a$ | $2b$ |
| Distance between foci | $2ae$ | $2be$ |
| Distance between directrices | $2a/e$ | $2b/e$ |
| Tangents at the vertices | $x = -a, x = a$ | $y = b, y = -b$ |

Example: 9 The equation of a directrix of the ellipse $\frac{x^2}{16} + \frac{y^2}{25} = 1$ is

- (a) $y = \frac{25}{3}$ (b) $x = 3$ (c) $x = -3$ (d) $x = \frac{3}{25}$

Solution: (a) From the given equation of ellipse $a^2 = 16, b^2 = 25$ (since $b > a$)

$$\text{So, } a^2 = b^2(1 - e^2), \therefore 16 = 25(1 - e^2) \Rightarrow 1 - e^2 = \frac{16}{25} \Rightarrow e^2 = \frac{9}{25} \Rightarrow e = \frac{3}{5}$$

$$\therefore \text{One directrix is } y = \frac{b}{e} = \frac{5}{3/5} = \frac{25}{3}$$

Example: 10 The distances from the foci of $P(x_1, y_1)$ on the ellipse $\frac{x^2}{9} + \frac{y^2}{25} = 1$ are

- (a) $4 \pm \frac{5}{4} y_1$ (b) $5 \pm \frac{4}{5} x_1$ (c) $5 \pm \frac{4}{5} y_1$ (d) None of these

Solution: (c) For the given ellipse $b > a$ so the two foci lie on y-axis and their coordinates are $(0, \pm be)$,

$$\text{Where } b = 5, a = 3. \text{ So } e = \sqrt{1 - \frac{a^2}{b^2}} = \sqrt{1 - \frac{9}{25}} = \frac{4}{5}$$

The focal distances of a point (x_1, y_1) on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, Where $b^2 > a^2$ are given by

$$b \pm ey_1. \text{ So, Required distances are } b \pm ey_1 = 5 \pm \frac{4}{5} y_1.$$

5.2.4 PARAMETRIC FORM OF THE ELLIPSE

Let the equation of ellipse in standard form will be given by $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

Then the equation of ellipse in the parametric form will be given by $x = a \cos \phi, y = b \sin \phi$, where ϕ is the eccentric angle whose value vary from $0 \leq \phi < 2\pi$. Therefore coordinate of any point P on the ellipse will be given by $(a \cos \phi, b \sin \phi)$

Example: 11 The curve represented by $x = 3(\cos t + \sin t), y = 4(\cos t - \sin t)$ is

- (a) Ellipse (b) Parabola (c) Hyperbola (d) Circle

Solution: (a) Given, $x = 3(\cos t + \sin t), y = 4(\cos t - \sin t) \Rightarrow \frac{x}{3} = (\cos t + \sin t), \frac{y}{4} = (\cos t - \sin t)$

Squaring and adding, we get $\frac{x^2}{9} + \frac{y^2}{16} = (1 + \sin 2t) + (1 - \sin 2t) \Rightarrow \frac{x^2}{9} + \frac{y^2}{16} = 2$, which represents ellipse.

Example: 12 The distance of the point 'P' on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ from a focus is

- (a) $a(e + \cos \theta)$ (b) $a(e - \cos \theta)$ (c) $a(1 + e \cos \theta)$ (d) $a(1 + 2e \cos \theta)$

Solution: (c) Focal distance of any point $P(x, y)$ on the ellipse is equal to $SP = a + ex$. Here $x = a \cos \theta$.

$$\text{Hence, } SP = a + ae \cos \theta = a(1 + e \cos \theta)$$

5.2.5 SPECIAL FORMS OF AN ELLIPSE

(1) If the centre of the ellipse is at point (h, k) and the directions of the axes are parallel to the coordinate axes, then its equation is $\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$

If we shift the origin at (h, k) without rotating the coordinate axes, then $x = X + h$ and $y = Y + k$

(2) If the equation of the curve is $\frac{(lx + my + n)^2}{a^2} + \frac{(mx - ly + p)^2}{b^2} = 1$ where $lx + my + n = 0$ and $mx - ly + p = 0$ are perpendicular lines, then we substitute $\frac{lx + my + n}{\sqrt{l^2 + m^2}} = X$, $\frac{mx - ly + p}{\sqrt{l^2 + m^2}} = Y$, to put the equation in the standard form.

Example: 13 The foci of the ellipse $25(x+1)^2 + 9(y+2)^2 = 225$ are

- (a) $(-1, 2), (6, 1)$ (b) $(-1, -2), (1, 6)$ (c) $(1, -2), (1, -6)$ (d) $(-1, 2), (-1, -6)$

Solution: (d) Given ellipse is $\frac{(x+1)^2}{9} + \frac{(y+2)^2}{25} = 1$ i.e. $\frac{X^2}{9} + \frac{Y^2}{25} = 1$, where $X = x+1$ and $Y = y+2$

Here $a^2 = 25, b^2 = 9$ [Type : $\frac{X^2}{b^2} + \frac{Y^2}{a^2} = 1$]

Eccentricity is given by $e^2 = \frac{a^2 - b^2}{a^2} = \frac{25 - 9}{25} = \frac{16}{25}$, $\therefore e = \frac{4}{5}$


Foci are given by $Y = \pm ae = \pm 5\left(\frac{4}{5}\right) = \pm 4$

$X = 0 \Rightarrow y + 2 = \pm 4 \Rightarrow y = -2 \pm 4 = -6$ or 2

$x + 1 = 0 \Rightarrow x = -1$. Hence foci are $(-1, -6)$ or $(-1, 2)$.

5.2.6 POSITION OF A POINT WITH RESPECT TO AN ELLIPSE

Let $P(x_1, y_1)$ be any point and let $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is the equation of an ellipse. The point lies outside, on or inside the ellipse as if $S_1 = \frac{x_1^2}{a^2} + \frac{y_1^2}{b^2} - 1 > 0, = 0, < 0$



Example: 14 Let E be the ellipse $\frac{x^2}{9} + \frac{y^2}{4} = 1$ and C be the circle $x^2 + y^2 = 9$. Let P and Q be the points $(1, 2)$ and $(2, 1)$ respectively. Then

- (a) Q lies inside C but outside E (b) Q lies outside both C and E
 (c) P lies inside both C and E (d) P lies inside C but outside E

Solution: (d) The given ellipse is $\frac{x^2}{9} + \frac{y^2}{4} = 1$. The value of the expression $\frac{x^2}{9} + \frac{y^2}{4} - 1$ is positive for $x=1, y=2$ and negative for $x=2, y=1$. Therefore P lies outside E and Q lies inside E . The value of the expression $x^2 + y^2 - 9$ is negative for both the points P and Q . Therefore P and Q both lie inside C . Hence P lies inside C but outside E .

5.2.7 INTERSECTION OF A LINE AND AN ELLIPSE

Let the ellipse be $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ (i) and the given line be $y = mx + c$ (ii)

Eliminating y from equation (i) and (ii), then $\frac{x^2}{a^2} + \frac{(mx+c)^2}{b^2} = 1$

i.e., $(a^2m^2 + b^2)x^2 + 2mcax + a^2(c^2 - b^2) = 0$

The above equation being a quadratic in x , its discriminant $= 4m^2c^2a^4 - 4a^2(a^2m^2 + b^2)(c^2 - b^2)$
 $= b^2\{(a^2m^2 + b^2) - c^2\}$

Hence the line intersects the ellipse in two distinct points if $a^2m^2 + b^2 > c^2$ in one point if $c^2 = a^2m^2 + b^2$ and does not intersect if $a^2m^2 + b^2 < c^2$.

5.2.8 EQUATIONS OF TANGENT IN DIFFERENT FORMS

(1) **Point form:** The equation of the tangent to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ at the point (x_1, y_1) is $\frac{xx_1}{a^2} + \frac{yy_1}{b^2} = 1$

(2) **Slope form:** If the line $y = mx + c$ touches the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, then $c^2 = a^2m^2 + b^2$. Hence, the straight line $y = mx \pm \sqrt{a^2m^2 + b^2}$ always represents the tangents to the ellipse.

Points of contact: Line $y = mx \pm \sqrt{a^2m^2 + b^2}$ touches the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ at

$$\left(\frac{\pm a^2m}{\sqrt{a^2m^2 + b^2}}, \frac{\pm b^2}{\sqrt{a^2m^2 + b^2}} \right)$$

(3) **Parametric form:** The equation of tangent at any point $\phi(a\cos\phi, b\sin\phi)$ is $\frac{x}{a}\cos\phi + \frac{y}{b}\sin\phi = 1$

Note : \square The straight line $lx + my + n = 0$ touches the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, if $a^2l^2 + b^2m^2 = n^2$.

\square The line $x\cos\alpha + y\sin\alpha = p$ touches the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, if $a^2\cos^2\alpha + b^2\sin^2\alpha = p^2$ and

that point of contact is $\left(\frac{a^2\cos\alpha}{p}, \frac{b^2\sin\alpha}{p} \right)$.

\square Two tangents can be drawn from a point to an ellipse. The two tangents are real and distinct or coincident or imaginary according as the given point lies outside, on or inside the ellipse.

- The tangents at the extremities of latus-rectum of an ellipse intersect on the corresponding directrix.

Important Tips

- ☞ A circle of radius r is concentric with the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, then the common tangent is inclined to the major axis at an angle $\tan^{-1} \sqrt{\frac{r^2 - b^2}{a^2 - r^2}}$.
- ☞ The locus of the foot of the perpendicular drawn from centre upon any tangent to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is $(x^2 + y^2)^2 = a^2 x^2 + b^2 y^2$ or $r^2 = a^2 \cos^2 \theta + b^2 \sin^2 \theta$ (in polar coordinates)
- ☞ The locus of the mid points of the portion of the tangents to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ intercepted between the axes is $a^2 y^2 + b^2 x^2 = 4x^2 y^2$.
- ☞ The product of the perpendiculars from the foci to any tangent of an ellipse is equal to the square of the semi minor axis, and the feet of these perpendiculars lie on the auxiliary circle.

Example: 15 The number of values of 'c' such that the straight line $y = 4x + c$ touches the curve $\frac{x^2}{4} + y^2 = 1$ is

- (a) 0 (b) 1 (c) 2 (d) Infinite

Solution: (c) We know that the line $y = mx + c$ touches the curve $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ iff $c^2 = a^2 m^2 + b^2$

$$\text{Here, } a^2 = 4, b^2 = 1, m = 4 \quad \therefore c^2 = 64 + 1 \Rightarrow c = \pm \sqrt{65}$$

Example: 16 On the ellipse $4x^2 + 9y^2 = 1$, the points at which the tangents are parallel to the line $8x = 9y$ are

- (a) $\left(\frac{2}{5}, \frac{1}{5}\right)$ (b) $\left(\frac{-2}{5}, \frac{1}{5}\right)$ (c) $\left(\frac{-2}{5}, \frac{-1}{5}\right)$ (d) $\left(\frac{2}{5}, \frac{-1}{5}\right)$

Solution: (b,d) Ellipse is $\frac{x^2}{1/4} + \frac{y^2}{1/9} = 1 \Rightarrow a^2 = \frac{1}{4}, b^2 = \frac{1}{9}$. The equation of its tangent is $4xx' + 9yy' = 1$

$$\therefore m = -\frac{4x'}{9y'} = \frac{8}{9} \Rightarrow x' = -2y' \text{ and } 4x'^2 + 9y'^2 = 1 \Rightarrow 4x'^2 + 9\frac{x'^2}{4} = 1 \Rightarrow x' = \pm \frac{2}{5}$$

$$\text{When } x' = \frac{2}{5}, \text{ then } y' = \frac{-1}{5} \text{ and when } x' = \frac{-2}{5}, \text{ then } y' = \frac{1}{5}.$$

$$\text{Hence points are } \left(\frac{2}{5}, \frac{-1}{5}\right), \left(\frac{-2}{5}, \frac{1}{5}\right)$$

Example: 17 If any tangent to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ intercepts equal lengths l on the axes, then $l =$

- (a) $a^2 + b^2$ (b) $\sqrt{a^2 + b^2}$ (c) $(a^2 + b^2)^2$ (d) None of these

Solution: (b) The equation of any tangent to the given ellipse is $\frac{x}{a} \cos \theta + \frac{y}{b} \sin \theta = 1$

$$\text{This line meets the coordinate axes at } P\left(\frac{a}{\cos \theta}, 0\right) \text{ and } Q\left(0, \frac{b}{\sin \theta}\right)$$

$$\therefore \frac{a}{\cos \theta} = l = \frac{b}{\sin \theta} \Rightarrow \cos \theta = \frac{a}{l} \text{ and } \sin \theta = \frac{b}{l} \Rightarrow \cos^2 \theta + \sin^2 \theta = \frac{a^2}{l^2} + \frac{b^2}{l^2} \Rightarrow l^2 = a^2 + b^2 \Rightarrow l = \sqrt{a^2 + b^2}.$$

Example: 18 The area of the quadrilateral formed by the tangents at the end points of latus-rectum to the ellipse $\frac{x^2}{9} + \frac{y^2}{5} = 1$, is

- (a) $27/4$ sq. units (b) 9 sq. units (c) $27/2$ sq. units (d) 27 sq. units

Solution: (d) By symmetry the quadrilateral is a rhombus. So area is four times the area of the right angled triangle formed by the tangents and axes in the 1st quadrant.

Now $ae = \sqrt{a^2 - b^2} \Rightarrow ae = 2 \Rightarrow$ Tangent (in the first quadrant) at one end of latus rectum $\left(2, \frac{5}{3}\right)$

$$\text{is } \frac{2}{9}x + \frac{5}{3} \cdot \frac{y}{5} = 1$$

$$\text{i.e. } \frac{x}{9/2} + \frac{y}{3} = 1. \text{ Therefore area} = 4 \cdot \frac{1}{2} \cdot \frac{9}{2} \cdot 3 = 27 \text{ sq units.}$$

5.2.9 EQUATION OF PAIR OF TANGENTS $SS_1 = T^2$


Pair of tangents: Let $P(x_1, y_1)$ be any point lying outside the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and let a pair of tangents PA, PB can be drawn to it from P .

Then the equation of pair of tangents PA and PB is $SS_1 = T^2$

$$\text{where } S \equiv \frac{x^2}{a^2} + \frac{y^2}{b^2} - 1 = 0$$

$$S_1 \equiv \frac{x_1^2}{a^2} + \frac{y_1^2}{b^2} - 1 = 0$$

$$T \equiv \frac{xx_1}{a^2} + \frac{yy_1}{b^2} - 1 = 0$$




Director circle: The director circle is the locus of points from which perpendicular tangents are drawn to the ellipse.

Let $P(x_1, y_1)$ be any point on the locus. Equation of tangents through $P(x_1, y_1)$ is given by $SS_1 = T^2$

$$\text{i.e., } \left(\frac{x^2}{a^2} + \frac{y^2}{b^2} - 1 \right) \left(\frac{x_1^2}{a^2} + \frac{y_1^2}{b^2} - 1 \right) = \left[\frac{xx_1}{a^2} + \frac{yy_1}{b^2} - 1 \right]^2$$

They are perpendicular, So coeff. of x^2 + coeff. of $y^2 = 0$

$$\therefore \left(\frac{1}{a^2} + \frac{1}{b^2} \right) \left(\frac{x_1^2}{a^2} + \frac{y_1^2}{b^2} - 1 \right) - \left(\frac{x_1^2}{a^4} + \frac{y_1^2}{b^4} \right) = 0 \text{ or } x_1^2 + y_1^2 = a^2 + b^2$$



Hence locus of $P(x_1, y_1)$ i.e., equation of director circle is $x^2 + y^2 = a^2 + b^2$

Example: 19 The angle between the pair of tangents drawn from the point $(1, 2)$ to the ellipse $3x^2 + 2y^2 = 5$ is

- (a) $\tan^{-1}(12/5)$ (b) $\tan^{-1}(6/\sqrt{5})$ (c) $\tan^{-1}(12/\sqrt{5})$ (d) $\tan^{-1}(6/5)$

Solution: (c) The combined equation of the pair of tangents drawn from $(1, 2)$ to the ellipse $3x^2 + 2y^2 = 5$ is $(3x^2 + 2y^2 - 5)(3 + 8 - 5) = (3x + 4y - 5)^2$ [using $SS_1 = T^2$]

$$\Rightarrow 9x^2 - 24xy - 4y^2 + \dots = 0$$

The angle between the lines given by this equation is $\tan\theta = \frac{2\sqrt{h^2 - ab}}{a+b}$

Where $a=9$, $h=-12$, $b=-4 \Rightarrow \tan\theta = 12/\sqrt{5} \Rightarrow \theta = \tan^{-1}(12/\sqrt{5})$

Example: 20 The locus of the point of intersection of the perpendicular tangents to the ellipse $\frac{x^2}{9} + \frac{y^2}{4} = 1$ is

- (a) $x^2 + y^2 = 9$ (b) $x^2 + y^2 = 4$ (c) $x^2 + y^2 = 13$ (d) $x^2 + y^2 = 5$

Solution: (c) The locus of point of intersection of two perpendicular tangents drawn on the ellipse is $x^2 + y^2 = a^2 + b^2$, which is called “director circle”.

Given ellipse is $\frac{x^2}{9} + \frac{y^2}{4} = 1$. \therefore Locus is $x^2 + y^2 = 9 + 4$, i.e. $x^2 + y^2 = 13$.

Example: 21 The locus of the middle point of the intercept of the tangents drawn from an external point to the ellipse $x^2 + 2y^2 = 2$ between the coordinate axes, is

- (a) $\frac{1}{x^2} + \frac{1}{2y^2} = 1$ (b) $\frac{1}{4x^2} + \frac{1}{2y^2} = 1$ (c) $\frac{1}{2x^2} + \frac{1}{4y^2} = 1$ (d) $\frac{1}{2x^2} + \frac{1}{y^2} = 1$

Solution: (c) Let the point of contact be $R = (\sqrt{2}\cos\theta, \sin\theta)$


Equation of tangent AB is $\frac{x}{\sqrt{2}}\cos\theta + y\sin\theta = 1$

$\Rightarrow A = (\sqrt{2}\sec\theta, 0); B = (0, \csc\theta)$

Let the middle point Q of AB be (h, k) .

$\Rightarrow h = \frac{\sec\theta}{\sqrt{2}}, k = \frac{\csc\theta}{2} \Rightarrow \cos\theta = \frac{1}{h\sqrt{2}}, \sin\theta = \frac{1}{2k} \Rightarrow \frac{1}{2h^2} + \frac{1}{4k^2} = 1$


Thus required locus is $\frac{1}{2x^2} + \frac{1}{4y^2} = 1$



5.2.10 EQUATIONS OF NORMAL IN DIFFERENT FORMS

(1) **Point form:** The equation of the normal at (x_1, y_1) to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is

$$\frac{a^2 x}{x_1} - \frac{b^2 y}{y_1} = a^2 - b^2.$$



(2) **Parametric form:** The equation of the normal to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ at $(a\cos\phi, b\sin\phi)$ is $ax\sec\phi - by\csc\phi = a^2 - b^2$.

(3) **Slope form:** If m is the slope of the normal to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, then the equation of normal is

$$y = mx \pm \frac{m(a^2 - b^2)}{\sqrt{a^2 + b^2 m^2}}$$

The coordinates of the point of contact are $\left(\frac{\pm a^2}{\sqrt{a^2 + b^2 m^2}}, \frac{\pm mb^2}{\sqrt{a^2 + b^2 m^2}} \right)$


Note : \square If $y = mx + c$ is the normal of $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, then condition of normality is $c^2 = \frac{m^2(a^2 - b^2)^2}{(a^2 + b^2 m^2)}$.

\square The straight line $lx + my + n = 0$ is a normal to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, if $\frac{a^2}{l^2} + \frac{b^2}{m^2} = \left(\frac{a^2 - b^2}{n^2} \right)^2$.

\square Four normals can be drawn from a point to an ellipse.

Important Tips

\Rightarrow If S be the focus and G be the point where the normal at P meets the axis of an ellipse, then $SG = eSP$, and the tangent and normal at P bisect the external and internal angles between the focal distances of P.



- \Rightarrow Any point P of an ellipse is joined to the extremities of the major axis then the portion of a directrix intercepted by them subtends a right angle at the corresponding focus.
- \Rightarrow With a given point and line as focus and directrix, a series of ellipse can be described. The locus of the extremities of their minor axis is a parabola.
- \Rightarrow The equations to the normals at the end of the latera recta and that each passes through an end of the minor axis, if $e^4 + e^2 + 1 = 0$
- \Rightarrow If two concentric ellipse be such that the foci of one be on the other and if e and e' be their eccentricities. Then the angle between their axes is $\cos^{-1} \sqrt{\frac{e^2 + e'^2 - 1}{ee'}}$.

Example: 22 The equation of normal at the point (0, 3) of the ellipse $9x^2 + 5y^2 = 45$ is

(a) $y - 3 = 0$

(b) $y + 3 = 0$

(c) x-axis

(d) y-axis

Solution: (d) For $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, equation of normal at point (x_1, y_1) , is $\frac{(x - x_1)a^2}{x_1} = \frac{(y - y_1)b^2}{y_1}$

Here, $(x_1, y_1) = (0, 3)$ and $a^2 = 5$, $b^2 = 9$, Therefore $\frac{(x-0)}{0} \cdot 5 = \frac{(y-3)}{3} \cdot 9$ or $x = 0$ i.e., y-axis.

Example: 23 If the normal at any point P on the ellipse cuts the major and minor axes in G and g respectively and C be the centre of the ellipse, then

- (a) $a^2(CG)^2 + b^2(Cg)^2 = (a^2 - b^2)^2$ (b) $a^2(CG)^2 - b^2(Cg)^2 = (a^2 - b^2)^2$
 (c) $a^2(CG)^2 - b^2(Cg)^2 = (a^2 + b^2)^2$ (d) None of these

Solution: (a) Let at point (x_1, y_1) normal will be $\frac{(x-x_1)}{x_1} a^2 = \frac{(y-y_1)}{y_1} b^2$

At G, $y = 0 \Rightarrow x = CG = \frac{x_1(a^2 - b^2)}{a^2}$ and at g, $x = 0 \Rightarrow y = Cg = \frac{y_1(b^2 - a^2)}{b^2}$

$$\frac{x_1^2}{a^2} + \frac{y_1^2}{b^2} = 1 \Rightarrow a^2(CG)^2 + b^2(Cg)^2 = (a^2 - b^2)^2.$$

Example: 24 The equation of the normal to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ at the positive end of the latus-rectum is

- (a) $x + ey + e^3a = 0$ (b) $x - ey - e^3a = 0$ (c) $x - ey - e^2a = 0$ (d) None of these

Solution: (b) The equation of the normal at (x_1, y_1) to the given ellipse is $\frac{a^2x}{x_1} - \frac{b^2y}{y_1} = a^2 - b^2$. Here, $x_1 = ae$ and $y_1 = \frac{b^2}{a}$

So, the equation of the normal at positive end of the latus-rectum is

$$\frac{a^2x}{ae} - \frac{b^2y}{b^2/a} = a^2 - b^2 \quad [\because b^2 = a^2(1 - e^2)] \Rightarrow \frac{ax}{e} - ay = a^2 - b^2 \Rightarrow x - ey - e^3a = 0$$


5.2.11 AUXILIARY CIRCLE

The circle described on the major axis of an ellipse as diameter is called an auxiliary circle of the ellipse.

If $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is an ellipse, then its auxiliary circle is $x^2 + y^2 = a^2$

Eccentric angle of a point: Let P be any point on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$. Draw PM perpendicular from P on the major axis of the ellipse and produce MP to meet the auxiliary circle in Q. Join CQ. The angle $\angle XCQ = \phi$ is called the eccentric angle of the point P on the ellipse.

Note that the angle $\angle XCP$ is not the eccentric angle of point P.




5.2.12 PROPERTIES OF ECCENTRIC ANGLES OF THE CO-NORMAL POINTS

(1) The sum of the eccentric angles of the co-normal points on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is equal to odd multiple of π .

(2) If α, β, γ are the eccentric angles of three points on the ellipse, the normals at which are concurrent, then $\sin(\alpha + \beta) + \sin(\beta + \gamma) + \sin(\gamma + \alpha) = 0$.

(3) **Co-normal points lie on a fixed curve:** Let $P(x_1, y_1), Q(x_2, y_2), R(x_3, y_3)$ and $S(x_4, y_4)$ be co-normal points, then $PQRS$ lie on the curve $(a^2 - b^2)xy + b^2kx - a^2hy = 0$.

This curve is called Apollonian rectangular hyperbola.



Note : \square The feet of the normals from any fixed point to the ellipse lie at the intersections of the apollonian rectangular hyperbola with the ellipse.

Important Tips

The area of the triangle formed by the three points, on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, whose eccentric angles are θ, ϕ and ψ is $2ab \sin\left(\frac{\phi - \psi}{2}\right) \sin\left(\frac{\psi - \theta}{2}\right) \sin\left(\frac{\theta - \phi}{2}\right)$.

The eccentricity of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is given by $2 \cot w = \frac{e^2 \sin 2\theta}{\sqrt{1 - e^2}}$, where w is one of the angles between the normals at the points whose eccentric angles are θ and $\frac{\pi}{2} + \theta$.

Example: 25 The eccentric angle of a point on the ellipse $\frac{x^2}{6} + \frac{y^2}{2} = 1$, whose distance from the centre of the ellipse is 2, is

(a) $\pi/4$

(b) $3\pi/2$

(c) $5\pi/3$

(d) $7\pi/6$

Solution: (a) Let θ be the eccentric angle of the point P . Then the coordinates of P are $(\sqrt{6} \cos \theta, \sqrt{2} \sin \theta)$

The centre of the ellipse is at the origin, It is given that $OP = 2$

$$\Rightarrow \sqrt{6 \cos^2 \theta + 2 \sin^2 \theta} = 2 \Rightarrow 6 \cos^2 \theta + 2 \sin^2 \theta = 4 \Rightarrow 3 \cos^2 \theta + \sin^2 \theta = 2 \Rightarrow 2 \sin^2 \theta = 1$$

$$\Rightarrow \sin^2 \theta = \frac{1}{2} \Rightarrow \sin \theta = \pm \frac{1}{\sqrt{2}} \Rightarrow \theta = \pm \pi/4$$

Example: 26 The area of the rectangle formed by the perpendiculars from the centre of the ellipse to the tangent and normal at the point-whose eccentric angle is $\pi/4$, is

(a) $\left(\frac{a^2 - b^2}{a^2 + b^2}\right)ab$ (b) $\left(\frac{a^2 + b^2}{a^2 - b^2}\right)ab$ (c) $\frac{1}{ab} \left(\frac{a^2 - b^2}{a^2 + b^2}\right)ab$ (d) $\frac{1}{ab} \left(\frac{a^2 + b^2}{a^2 - b^2}\right)ab$

Solution: (a) The given point is $(a \cos \pi/4, b \sin \pi/4)$ i.e. $\left(\frac{a}{\sqrt{2}}, \frac{b}{\sqrt{2}}\right)$.

So, the equation of the tangent at this point is $\frac{x}{a} + \frac{y}{b} = \sqrt{2}$ (i)

$$\therefore p_1 = \text{length of the perpendicular from } (0, 0) \text{ on (i)} = \left| \frac{\frac{0}{a} + \frac{0}{b} - \sqrt{2}}{\sqrt{1/a^2 + 1/b^2}} \right| = \frac{\sqrt{2}ab}{\sqrt{a^2 + b^2}}$$


$$\text{Equation of the normal at } \left(\frac{a}{\sqrt{2}}, \frac{b}{\sqrt{2}} \right) \text{ is } \frac{a^2 x}{a/\sqrt{2}} - \frac{b^2 y}{b/\sqrt{2}} = a^2 - b^2 \Rightarrow \sqrt{2}ax - \sqrt{2}by = a^2 - b^2 \quad \dots (ii)$$

$$\text{Therefore, } p_2 = \text{length of the perpendicular from } (0, 0) \text{ on (ii)} = \frac{a^2 b^2}{\sqrt{(\sqrt{2}a)^2 + (-\sqrt{2}b)^2}} = \frac{a^2 - b^2}{\sqrt{2(a^2 + b^2)}}$$

$$\text{So, area of the rectangle} = p_1 p_2 = \frac{\sqrt{2}ab}{\sqrt{a^2 + b^2}} \times \frac{a^2 - b^2}{\sqrt{2(a^2 + b^2)}} = \left(\frac{a^2 - b^2}{a^2 + b^2} \right) ab$$


5.2.13 CHORD OF CONTACT

If PQ and PR be the tangents through point $P(x_1, y_1)$ to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, then the equation of the chord of contact QR is $\frac{xx_1}{a^2} + \frac{yy_1}{b^2} = 1$ or $T = 0$ at (x_1, y_1)



5.2.14 EQUATION OF CHORD WITH MID POINT (x_1, y_1)

The equation of the chord of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, whose mid point be (x_1, y_1) is $T = S_1$, where $T = \frac{xx_1}{a^2} + \frac{yy_1}{b^2} - 1 = 0$, $S_1 = \frac{x_1^2}{a^2} + \frac{y_1^2}{b^2} - 1 = 0$



5.2.15 EQUATION OF THE CHORD JOINING TWO POINTS ON AN ELLIPSE

Let $P(a \cos \theta, b \sin \theta)$; $Q(a \cos \phi, b \sin \phi)$ be any two points of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$. Then, the equation of the chord joining these two points is $y - b \sin \theta = \frac{b \sin \phi - b \sin \theta}{a \cos \phi - a \cos \theta} (x - a \cos \theta)$

Thus, the equation of the chord joining two points having eccentric angles θ and ϕ on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is $\frac{x}{a} \cos \left(\frac{\theta + \phi}{2} \right) + \frac{y}{b} \sin \left(\frac{\theta + \phi}{2} \right) = \cos \left(\frac{\theta - \phi}{2} \right)$

Note : ☐ If the chord joining two points whose eccentric angles are α and β cut the major axis of an ellipse at a distance 'c' from the centre, then $\tan \frac{\alpha}{2} \tan \frac{\beta}{2} = \frac{c - a}{c + a}$.

☐ If α and β be the eccentric angles of the extremities of a focal chord of an ellipse of eccentricity e , then $\tan \frac{\alpha}{2} \tan \frac{\beta}{2} + \frac{1 \pm e}{1 \mp e} = 0$.

Example: 27 What will be the equation of that chord of ellipse $\frac{x^2}{36} + \frac{y^2}{9} = 1$ which passes from the point (2,1) and bisected on the point

- (a) $x + y = 2$ (b) $x + y = 3$ (c) $x + 2y = 1$ (d) $x + 2y = 4$

Solution: (d) Let required chord meets to ellipse on the points P and Q whose coordinates are (x_1, y_1) and (x_2, y_2) respectively

⊖ Point (2,1) is mid point of chord PQ

$$\therefore 2 = \frac{1}{2}(x_1 + x_2) \text{ or } x_1 + x_2 = 4 \text{ and } 1 = \frac{1}{2}(y_1 + y_2) \text{ or } y_1 + y_2 = 2$$

Again points (x_1, y_1) and (x_2, y_2) are situated on ellipse; $\therefore \frac{x_1^2}{36} + \frac{y_1^2}{9} = 1$ and $\frac{x_2^2}{36} + \frac{y_2^2}{9} = 1$

$$\text{On subtracting } \frac{x_2^2 - x_1^2}{36} + \frac{y_2^2 - y_1^2}{9} = 0 \text{ or } \frac{y_2 - y_1}{x_2 - x_1} = -\frac{(x_2 + x_1)}{4(y_2 + y_1)} = \frac{-4}{4 \times 2} = \frac{-1}{2}$$

$$\therefore \text{Gradient of chord } PQ = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-1}{2}$$

Therefore, required equation of chord PQ is as follows, $y - 1 = -\frac{1}{2}(x - 2)$ or $x + 2y = 4$

Alternative: $S_1 = T$ (If mid point of chord is known)

$$\therefore \frac{2^2}{36} + \frac{1^2}{9} - 1 = \frac{2x}{36} + \frac{1y}{9} - 1 \Rightarrow x + 2y = 4$$

Example: 28 What will be the equation of the chord of contact of tangents drawn from (3, 2) to the ellipse $x^2 + 4y^2 = 9$

- (a) $3x + 8y = 9$ (b) $3x + 8y = 25$ (c) $3x + 4y = 9$ (d) $3x + 8y + 9 = 0$

Solution: (a) The required equation is $T = 0$ i.e., $3x + 4(2y) - 9 = 0$ or $3x + 8y = 9$.

Example: 29 A tangent to the ellipse $x^2 + 4y^2 = 4$ meets the ellipse $x^2 + 2y^2 = 6$ at P and Q . The angle between the tangents at P and Q of the ellipse $x^2 + 2y^2 = 6$ is

- (a) $\frac{\pi}{2}$ (b) $\frac{\pi}{3}$ (c) $\frac{\pi}{4}$ (d) $\frac{\pi}{6}$

Solution: (a) The given ellipse $x^2 + 4y^2 = 4$ can be written as $\frac{x^2}{4} + \frac{y^2}{1} = 1$ (i)

Any tangent to ellipse (i) is $\frac{x}{2} \cos \theta + y \sin \theta = 1$

Second ellipse is $x^2 + 2y^2 = 6$, i.e., $\frac{x^2}{6} + \frac{y^2}{3} = 1$


Let the tangents at P, Q meet at (h, k) .

$$\therefore \text{Equation of } PQ, \text{ i.e. chord of contact is } \frac{hx}{6} + \frac{ky}{3} = 1 \quad \text{.....(iv)}$$

Since (ii) and (iv) represent the same line, $\therefore \frac{h/6}{(\cos \theta)/2} = \frac{k/3}{\sin \theta} = \frac{1}{1} \Rightarrow h = 3 \cos \theta$ and $k = 3 \sin \theta$

So, $h^2 + k^2 = 9$ or $x^2 + y^2 = 9$ is the locus of (h, k) which is the director circle of the ellipse

$$\frac{x^2}{6} + \frac{y^2}{3} = 1$$



\therefore The angle between the tangents at P and Q will be $\pi/2$.

Example: 30 The locus of mid-points of a focal chord of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is

- (a) $\frac{x^2}{a^2} + \frac{y^2}{b^2} = \frac{ex}{a}$ (b) $\frac{x^2}{a^2} - \frac{y^2}{b^2} = \frac{ex}{a}$ (c) $x^2 + y^2 = a^2 + b^2$ (d) None of these

Solution: (a) Let (h, k) be the mid point of a focal chord. Then its equation is $S_1 = T$ or $\frac{hx}{a^2} + \frac{ky}{b^2} = \frac{h^2}{a^2} + \frac{k^2}{b^2}$.

This passes through $(ae, 0)$, $\therefore \frac{hae}{a^2} = \frac{h^2}{a^2} + \frac{k^2}{b^2}$. So, locus of (h, k) is $\frac{xe}{a} = \frac{x^2}{a^2} + \frac{y^2}{b^2}$

Example: 31 If α and β are the eccentric angles of the extremities of a focal chord of an ellipse, then the eccentricity of the ellipse is

- (a) $\frac{\cos\alpha + \cos\beta}{\cos(\alpha - \beta)}$ (b) $\frac{\sin\alpha - \sin\beta}{\sin(\alpha - \beta)}$ (c) $\frac{\cos\alpha - \cos\beta}{\cos(\alpha - \beta)}$ (d) $\frac{\sin\alpha + \sin\beta}{\sin(\alpha + \beta)}$

Solution: (d) The equation of a chord joining points having eccentric angles α and β is given by



$$\frac{x}{a} \cos\left(\frac{\alpha + \beta}{2}\right) + \frac{y}{b} \sin\left(\frac{\alpha + \beta}{2}\right) = \cos\left(\frac{\alpha - \beta}{2}\right)$$

If it passes through $(ae, 0)$ then $e \cos\left(\frac{\alpha + \beta}{2}\right) = \cos\left(\frac{\alpha - \beta}{2}\right)$

$$\Rightarrow e = \frac{\cos\left(\frac{\alpha - \beta}{2}\right)}{\cos\left(\frac{\alpha + \beta}{2}\right)} \Rightarrow e = \frac{2 \sin\left(\frac{\alpha + \beta}{2}\right) \cos\left(\frac{\alpha - \beta}{2}\right)}{2 \sin\left(\frac{\alpha + \beta}{2}\right) \cos\left(\frac{\alpha + \beta}{2}\right)} \Rightarrow e = \frac{\sin\alpha + \sin\beta}{\sin(\alpha + \beta)}$$

5.2.16 POLE AND POLAR

Let $P(x_1, y_1)$ be any point inside or outside the ellipse. A chord through P intersects the ellipse at A and B respectively. If tangents to the ellipse at A and B meet at $Q(h, k)$ then locus of Q is called polar of P with respect to ellipse and point P is called pole.




Equation of polar: Equation of polar of the point (x_1, y_1) with respect to ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is given by

$$\frac{xx_1}{a^2} + \frac{yy_1}{b^2} = 1 \quad (i.e. \quad T = 0)$$

Coordinates of pole: The pole of the line $lx + my + n = 0$ with respect to ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is

$$P\left(\frac{-a^2l}{n}, \frac{-b^2m}{n}\right)$$



Note : ☐ The polar of any point on the directrix, passes through the focus.

☐ Any tangent is the polar of its own point of contact.

Properties of pole and polar

(1) If the polar of $P(x_1, y_1)$ passes through $Q(x_2, y_2)$, then the polar of $Q(x_2, y_2)$ goes through $P(x_1, y_1)$ and such points are said to be conjugate points.

(2) If the pole of a line $l_1x + m_1y + n_1 = 0$ lies on the another line $l_2x + m_2y + n_2 = 0$, then the pole of the second line will lie on the first and such lines are said to be conjugate lines.

(3) Pole of a given line is same as point of intersection of tangents at its extremities.

Example: 32 The pole of the straight line $x + 4y = 4$ with respect to ellipse $x^2 + 4y^2 = 4$ is

(a) (1, 4)

(b) (1, 1)

(c) (4, 1)

(d) (4, 4)

Solution: (b) Equation of polar of (x_1, y_1) w.r.t the ellipse is $xx_1 + 4yy_1 = 4$ (i)

Comparing with $x + 4y = 4$ (ii)

$$\frac{x_1}{1} = \frac{4y_1}{4} = 1 \Rightarrow x_1 = 1, y_1 = 1 \therefore \text{Coordinates of pole } (x_1, y_1) = (1, 1)$$

Example: 33 If the polar with respect to $y^2 = 4ax$ touches the ellipse $\frac{x^2}{\alpha^2} + \frac{y^2}{\beta^2} = 1$, the locus of its pole is

(a) $\frac{x^2}{\alpha^2} - \frac{y^2}{(4a^2\alpha^2/\beta^2)} = 1$

(b) $\frac{x^2}{\alpha^2} + \frac{\beta^2 y^2}{4a^2} = 1$

(c) $\alpha^2 x^2 + \beta^2 y^2 = 1$

(d) None of these

Solution: (a) Let $P(h, k)$ be the pole. Then the equation of the polar is $ky = 2a(x + h)$ or $y = \frac{2a}{k}x + \frac{2ah}{k}$.

This touches $\frac{x^2}{\alpha^2} + \frac{y^2}{\beta^2} = 1$, So $\left(\frac{2ah}{k}\right)^2 = \alpha^2 \left(\frac{2a}{k}\right)^2 + \beta^2$, (using $c^2 = a^2 m^2 + b^2$)


$$\Rightarrow 4a^2 h^2 = 4a^2 \alpha^2 + k^2 \beta^2. \text{ So, locus of } (h, k) \text{ is } 4a^2 x^2 = 4a^2 \alpha^2 + \beta^2 y^2 \text{ or } \frac{x^2}{\alpha^2} - \frac{y^2}{\left(\frac{4a^2 \alpha^2}{\beta^2}\right)} = 1$$

5.2.17 DIAMETER OF THE ELLIPSE

Definition : The locus of the mid- point of a system of parallel chords of an ellipse is called a diameter and the chords are called its double ordinates *i.e.* A line through the centre of an ellipse is called a diameter of the ellipse.

The point where the diameter intersects the ellipse is called the vertex of the diameter.


Equation of a diameter to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$: Let $y = mx + c$ be a system of parallel chords of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, where m is a constant and c is a variable.



The equation of the diameter bisecting the chords of slope m of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is $y = -\frac{b^2}{a^2 m} x$, which is passing through (0, 0).

Conjugate diameter: Two diameters of an ellipse are said to be conjugate diameter if each bisects all chords parallel to the other.

Conjugate diameter of circle *i.e.* AA' and BB' are perpendicular to each other. Hence, conjugate diameter of ellipse are PP' and QQ' . Hence, angle between conjugate diameters of ellipse $> 90^\circ$.



Now the coordinates of the four extremities of two conjugate diameters are

$$P(a \cos \phi, b \sin \phi); P'(-a \cos \phi, -b \sin \phi); Q(-a \sin \phi, b \cos \phi); Q'(a \sin \phi, -b \cos \phi)$$

If $y = m_1 x$ and $y = m_2 x$ be two conjugate diameters of an ellipse, then $m_1 m_2 = -\frac{b^2}{a^2}$

(1) Properties of diameters

(i) The tangent at the extremity of any diameter is parallel to the chords it bisects or parallel to the conjugate diameter.


(ii) The tangent at the ends of any chord meet on the diameter which bisects the chord.

(2) Properties of conjugate diameters

(i) The eccentric angles of the ends of a pair of conjugate diameters of an ellipse differ by a right angle,


$$\text{i.e. } \phi - \phi' = \frac{\pi}{2}$$

(ii) The sum of the squares of any two conjugate semi-diameters of an ellipse is constant and equal to the sum of the squares of the semi axes of the ellipse, *i.e.* $CP^2 + CD^2 = a^2 + b^2$




(iii) The product of the focal distances of a point on an ellipse is equal to the square of the semi-diameter which is conjugate to the diameter through the point,

$$\text{i.e., } SP \cdot S'P = CD^2$$



(iv) The tangents at the extremities of a pair of conjugate diameters form a parallelogram whose area is constant and equal to product of the axes, *i.e.*

Area of parallelogram = $(2a)(2b)$ = Area of rectangle contained under major and minor axes.



(v) The polar of any point with respect to ellipse is parallel to the diameter to the one on which the point lies. Hence obtain the equation of the chord whose mid point is (x_1, y_1) , *i.e.* chord is $T = S_1$.

(3) **Equi-conjugate diameters:** Two conjugate diameters are called equi-conjugate, if their lengths are equal *i.e.* $(CP)^2 = (CD)^2$

$$\therefore a^2 \cos^2 \phi + b^2 \sin^2 \phi = a^2 \sin^2 \phi + b^2 \cos^2 \phi$$

$$\Rightarrow a^2(\cos^2 \phi - \sin^2 \phi) - b^2(\cos^2 \phi - \sin^2 \phi) = 0 \Rightarrow (a^2 - b^2)(\cos^2 \phi - \sin^2 \phi) = 0$$

$$\ominus (a^2 - b^2) \neq 0, \therefore \cos 2\phi = 0. \text{ So, } \phi = \frac{\pi}{4} \text{ or } \frac{3\pi}{4}$$

$$\therefore (CP) = (CD) = \sqrt{\frac{(a^2 + b^2)}{2}} \text{ for equi-conjugate diameters.}$$

Important Tips

- ☞ If the point of intersection of the ellipses $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and $\frac{x^2}{\alpha^2} + \frac{y^2}{\beta^2} = 1$ be at the extremities of the conjugate diameters of the former, then $\frac{a^2}{\alpha^2} + \frac{b^2}{\beta^2} = 2$
- ☞ The sum of the squares of the reciprocal of two perpendicular diameters of an ellipse is constant.
- ☞ In an ellipse, the major axis bisects all chords parallel to the minor axis and vice-versa, therefore major and minor axes of an ellipse are conjugate diameters of the ellipse but they do not satisfy the condition $m_1 m_2 = -b^2/a^2$ and are the only perpendicular conjugate diameters.

Example: 34 If one end of a diameter of the ellipse $4x^2 + y^2 = 16$ is $(\sqrt{3}, 2)$, then the other end is

- (a) $(-\sqrt{3}, 2)$ (b) $(\sqrt{3}, -2)$ (c) $(-\sqrt{3}, -2)$ (d) $(0, 0)$

Solution: (c) Since every diameter of an ellipse passes through the centre and is bisected by it, therefore the coordinates of the other end are $(-\sqrt{3}, -2)$.

Example: 35 If θ and ϕ are eccentric angles of the ends of a pair of conjugate diameters of the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, \text{ then } \theta - \phi \text{ is equal to}$$

- (a) $\pm \frac{\pi}{2}$ (b) $\pm \pi$ (c) 0 (d) None of these

Solution: (a) Let $y = m_1 x$ and $y = m_2 x$ be a pair of conjugate diameter of an ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and let

$$P(a \cos \theta, b \sin \theta) \text{ and } Q(a \cos \phi, b \sin \phi) \text{ be ends of these two diameters. Then } m_1 m_2 = \frac{-b^2}{a^2}$$

$$\Rightarrow \frac{b \sin \theta - 0}{a \cos \theta - 0} \times \frac{b \sin \phi - 0}{a \cos \phi - 0} = \frac{-b^2}{a^2} \Rightarrow \sin \theta \sin \phi = -\cos \theta \cos \phi \Rightarrow \cos(\theta - \phi) = 0 \Rightarrow \theta - \phi = \pm \pi/2.$$


5.2.18 SUBTANGENT AND SUBNORMAL

Let the tangent and normal at $P(x_1, y_1)$ meet the x -axis at A and B respectively.

Length of subtangent at $P(x_1, y_1)$ to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is $DA = CA - CD = \frac{a^2}{x_1} - x_1$

Length of sub-normal at $P(x_1, y_1)$ to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is

$$BD = CD - CB = x_1 - \left(x_1 - \frac{b^2}{a^2} x_1 \right) = \frac{b^2}{a^2} x_1 = (1 - e^2) x_1.$$



Note : \square The tangent and normal to any point of an ellipse bisects respectively the internal and external angles between the focal radii of that point.

Example: 36 Length of subtangent and subnormal at the point $\left(\frac{-5\sqrt{3}}{2}, 2 \right)$ of the ellipse $\frac{x^2}{25} + \frac{y^2}{16} = 1$ are


- (a) $\left(\frac{5\sqrt{3}}{2} - \frac{10}{\sqrt{3}} \right), \frac{8\sqrt{3}}{5}$ (b) $\left(\frac{5\sqrt{3}}{2} + \frac{10}{\sqrt{3}} \right), \frac{8\sqrt{3}}{10}$ (c) $\left(\frac{5\sqrt{3}}{2} + \frac{12}{\sqrt{3}} \right), \frac{16\sqrt{3}}{5}$ (d) None of these

Solution: (a) Here $a^2 = 25, b^2 = 16, x_1 = \frac{-5\sqrt{3}}{2}$. Length of subtangent $= \left| \frac{a^2}{x_1} - x_1 \right| = \left| \frac{25}{-5\sqrt{3}/2} + \frac{5\sqrt{3}}{2} \right| = \left| \frac{5\sqrt{3}}{2} - \frac{10}{\sqrt{3}} \right|$.

$$\text{Length of subnormal} = \left| \frac{b^2}{a^2} x_1 \right| = \left| \frac{16}{25} \left(\frac{-5\sqrt{3}}{2} \right) \right| = \left| \frac{8\sqrt{3}}{5} \right|$$

5.2.19 CONCYCLIC POINTS

Any circle intersects an ellipse in two or four points. They are called concyclic points and the sum of their eccentric angles is an even multiple of π .



If $\alpha, \beta, \gamma, \delta$ be the eccentric angles of the four concyclic points on an ellipse, then $\alpha + \beta + \gamma + \delta = 2n\pi$, where n is any integer.


Note : \square The common chords of a circle and an ellipse are equally inclined to the axes of the ellipse.

Important Tips

- ☞ The centre of a circle $x^2 + y^2 + 2gx + 2fy + c = 0$ passing through the three points, on an ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ (whose eccentric angles are α, β, γ) is
- $$-g = \left(\frac{a^2 - b^2}{4a} \right) \{ \cos \alpha + \cos \beta + \cos \gamma + \cos(\alpha + \beta + \gamma) \} \text{ and } -f = \left(\frac{b^2 - a^2}{4a} \right) \{ \sin \alpha + \sin \beta + \sin \gamma - \sin(\alpha + \beta + \gamma) \}$$
- ☞ PCP and DCD are conjugate diameters of an ellipse and α is the eccentric angles of P . Then the eccentric angles of the point where the circle through P, P, D again cuts the ellipse is $\pi/2 - 3\alpha$.

5.2.20 REFLECTION PROPERTY OF AN ELLIPSE

Let S and S' be the foci and PN the normal at the point P of the ellipse, then $\angle SPS' = \angle SQS'$. Hence if an incoming light ray aimed towards one focus strike the concave side of the mirror in the shape of an ellipse then it will be reflected towards the other focus.



Example: 37 A ray emanating from the point $(-3, 0)$ is incident on the ellipse $16x^2 + 25y^2 = 400$ at the point P with ordinate 4. Then the equation of the reflected ray after first reflection is

- (a) $4x + 3y = 12$ (b) $3x + 4y = 12$ (c) $4x - 3y = 12$ (d) $3x - 4y = 12$

Solution: (a) For point P y-coordinate = 4

Given ellipse is $16x^2 + 25y^2 = 400$

$$16x^2 + 25(4)^2 = 400, \therefore x = 0$$


\therefore co-ordinate of P is $(0, 4)$

$$e^2 = 1 - \frac{16}{25} = \frac{9}{25}$$

$$\therefore e = \frac{3}{5}$$

\therefore Foci $(\pm ae, 0)$, i.e. $(\pm 3, 0)$

\therefore Equation of reflected ray (i.e. PS) is $\frac{x}{3} + \frac{y}{4} = 1$ or $4x + 3y = 12$.



ASSIGNMENT

DEFINITION OF THE ELLIPSE

Basic Level

1. If a bar of given length moves with its extremities on two fixed straight lines at right angles, then the locus of any point on bar marked on the bar describes a/an
 (a) Circle (b) Parabola (c) Ellipse (d) Hyperbola
2. If the eccentricity of an ellipse becomes zero, then it takes the form of
 (a) A circle (b) A parabola (c) A straight line (d) None of these
3. The locus of a variable point whose distance from $(-2, 0)$ is $\frac{2}{3}$ times its distance from the line $x = -\frac{9}{2}$, is
 (a) Ellipse (b) Parabola (c) Hyperbola (d) None of these
4. If A and B are two fixed points and P is a variable point such that $PA + PB = 4$, where $AB < 4$, then the locus of P is
 (a) A parabola (b) An ellipse (c) A hyperbola (d) None of these
5. Equation of the ellipse whose focus is $(6, 7)$ directrix is $x + y + 2 = 0$ and $e = 1/\sqrt{3}$ is
 (a) $5x^2 + 2xy + 5y^2 - 76x - 88y + 506 = 0$ (b) $5x^2 - 2xy + 5y^2 - 76x - 88y + 506 = 0$
 (c) $5x^2 - 2xy + 5y^2 + 76x + 88y - 506 = 0$ (d) None of these
6. The locus of the centre of the circle $x^2 + y^2 + 4x \cos \theta - 2y \sin \theta - 10 = 0$ is
 (a) An ellipse (b) A circle (c) A hyperbola (d) A parabola

STANDARD & OTHER FORMS OF AN ELLIPSE , TERMS RELATED TO AN ELLIPSE

Basic Level

7. The equation $2x^2 + 3y^2 = 30$ represents
 (a) A circle (b) An ellipse (c) A hyperbola (d) A parabola
8. The equation $\frac{x^2}{2-r} + \frac{y^2}{r-5} + 1 = 0$ represents an ellipse, if
 (a) $r > 2$ (b) $2 < r < 5$ (c) $r > 5$ (d) None of these
9. Equation of the ellipse with eccentricity $\frac{1}{2}$ and foci at $(\pm 1, 0)$ is
 (a) $\frac{x^2}{3} + \frac{y^2}{4} = 1$ (b) $\frac{x^2}{4} + \frac{y^2}{3} = 1$ (c) $\frac{x^2}{4} + \frac{y^2}{3} = \frac{4}{3}$ (d) None of these
10. The equation of the ellipse whose foci are $(\pm 5, 0)$ and one of its directrix is $5x = 36$, is
 (a) $\frac{x^2}{36} + \frac{y^2}{11} = 1$ (b) $\frac{x^2}{6} + \frac{y^2}{\sqrt{11}} = 1$ (c) $\frac{x^2}{6} + \frac{y^2}{11} = 1$ (d) None of these

11. The equation of ellipse whose distance between the foci is equal to 8 and distance between the directrix is 18, is
 (a) $5x^2 - 9y^2 = 180$ (b) $9x^2 + 5y^2 = 180$ (c) $x^2 + 9y^2 = 180$ (d) $5x^2 + 9y^2 = 180$
12. The equation of the ellipse whose one of the vertices is (0,7) and the corresponding directrix is $y = 12$, is
 (a) $95x^2 + 144y^2 = 4655$ (b) $144x^2 + 95y^2 = 4655$ (c) $95x^2 + 144y^2 = 13680$ (d) None of these
13. The equation of the ellipse whose centre is at origin and which passes through the points (-3, 1) and (2,-2) is
 (a) $5x^2 + 3y^2 = 32$ (b) $3x^2 + 5y^2 = 32$ (c) $5x^2 - 3y^2 = 32$ (d) $3x^2 + 5y^2 + 32 = 0$
14. An ellipse passes through the point (-3,1) and its eccentricity is $\sqrt{\frac{2}{5}}$. The equation of the ellipse is
 (a) $3x^2 + 5y^2 = 32$ (b) $3x^2 + 5y^2 = 25$ (c) $3x^2 + y^2 = 4$ (d) $3x^2 + y^2 = 9$
15. If the centre, one of the foci and semi- major axis of an ellipse be (0, 0), (0, 3) and 5 then its equation is
 (a) $\frac{x^2}{16} + \frac{y^2}{25} = 1$ (b) $\frac{x^2}{25} + \frac{y^2}{16} = 1$ (c) $\frac{x^2}{9} + \frac{y^2}{25} = 1$ (d) None of these
16. The equation of the ellipse whose latus rectum is 8 and whose eccentricity is $\frac{1}{\sqrt{2}}$, referred to the principal axes of coordinates, is
 (a) $\frac{x^2}{18} + \frac{y^2}{32} = 1$ (b) $\frac{x^2}{8} + \frac{y^2}{9} = 1$ (c) $\frac{x^2}{64} + \frac{y^2}{32} = 1$ (d) $\frac{x^2}{16} + \frac{y^2}{24} = 1$
17. The lengths of major and minor axes of an ellipse are 10 and 8 respectively and its major axis is along the y-axis. The equation of the ellipse referred to its centre as origin is
 (a) $\frac{x^2}{25} + \frac{y^2}{16} = 1$ (b) $\frac{x^2}{16} + \frac{y^2}{25} = 1$ (c) $\frac{x^2}{100} + \frac{y^2}{64} = 1$ (d) $\frac{x^2}{64} + \frac{y^2}{100} = 1$
18. The equation of the ellipse whose vertices are ($\pm 5, 0$) and foci are ($\pm 4, 0$) is
 (a) $9x^2 + 25y^2 = 225$ (b) $25x^2 + 9y^2 = 225$ (c) $3x^2 + 4y^2 = 192$ (d) None of these
19. The latus rectum of an ellipse is 10 and the minor axis is equal to the distance between the foci. The equation of the ellipse is
 (a) $x^2 + 2y^2 = 100$ (b) $x^2 + \sqrt{2}y^2 = 10$ (c) $x^2 - 2y^2 = 100$ (d) None of these
20. The eccentricity of the ellipse $4x^2 + 9y^2 = 36$, is
 (a) $\frac{1}{2\sqrt{3}}$ (b) $\frac{1}{\sqrt{3}}$ (c) $\frac{\sqrt{5}}{3}$ (d) $\frac{\sqrt{5}}{6}$
21. Eccentricity of the conic $16x^2 + 7y^2 = 112$ is
 (a) $\frac{3}{\sqrt{7}}$ (b) $\frac{7}{16}$ (c) $\frac{3}{4}$ (d) $\frac{4}{3}$
22. Eccentricity of the ellipse $9x^2 + 25y^2 = 225$ is
 (a) $\frac{3}{5}$ (b) $\frac{4}{5}$ (c) $\frac{9}{25}$ (d) $\frac{\sqrt{34}}{5}$

23. The eccentricity of the ellipse $25x^2 + 16y^2 = 400$ is
 (a) $\frac{3}{5}$ (b) $\frac{1}{3}$ (c) $\frac{2}{5}$ (d) $\frac{1}{5}$
24. For the ellipse $\frac{x^2}{64} + \frac{y^2}{28} = 1$, the eccentricity is
 (a) $\frac{3}{4}$ (b) $\frac{4}{3}$ (c) $\frac{2}{\sqrt{7}}$ (d) $\frac{1}{3}$
25. If the latus rectum of an ellipse be equal to half of its minor axis, then its eccentricity is
 (a) $\frac{3}{2}$ (b) $\frac{\sqrt{3}}{2}$ (c) $\frac{2}{3}$ (d) $\frac{\sqrt{2}}{3}$
26. If the length of the major axis of an ellipse is three times the length of its minor axis, then its eccentricity is
 (a) $\frac{1}{3}$ (b) $\frac{1}{\sqrt{3}}$ (c) $\frac{1}{\sqrt{2}}$ (d) $\frac{2\sqrt{2}}{3}$
27. The length of the latus rectum of an ellipse is $\frac{1}{3}$ of the major axis. Its eccentricity is
 (a) $\frac{2}{3}$ (b) $\sqrt{\frac{2}{3}}$ (c) $\frac{5 \times 4 \times 3}{7^3}$ (d) $\left(\frac{3}{4}\right)^4$
28. Eccentricity of the ellipse whose latus rectum is equal to the distance between two focus points, is
 (a) $\frac{\sqrt{5}+1}{2}$ (b) $\frac{\sqrt{5}-1}{2}$ (c) $\frac{\sqrt{5}}{2}$ (d) $\frac{\sqrt{3}}{2}$
29. If the distance between the foci of an ellipse be equal to its minor axis, then its eccentricity is
 (a) $\frac{1}{2}$ (b) $\frac{1}{\sqrt{2}}$ (c) $\frac{1}{3}$ (d) $\frac{1}{\sqrt{3}}$
30. The length of the latus rectum of the ellipse $\frac{x^2}{36} + \frac{y^2}{49} = 1$ is
 (a) $\frac{98}{6}$ (b) $\frac{72}{7}$ (c) $\frac{72}{14}$ (d) $\frac{98}{12}$
31. For the ellipse $3x^2 + 4y^2 = 12$, the length of latus rectum is
 (a) $\frac{3}{2}$ (b) 3 (c) $\frac{8}{3}$ (d) $\sqrt{\frac{3}{2}}$
32. The length of the latus rectum of the ellipse $9x^2 + 4y^2 = 1$, is
 (a) $\frac{3}{2}$ (b) $\frac{8}{3}$ (c) $\frac{4}{9}$ (d) $\frac{8}{9}$
33. In an ellipse, minor axis is 8 and eccentricity is $\frac{\sqrt{5}}{3}$. Then major axis is
 (a) 6 (b) 12 (c) 10 (d) 16
34. The distance between the foci of an ellipse is 16 and eccentricity is $\frac{1}{2}$. Length of the major axis of the ellipse is
 (a) 8 (b) 64 (c) 16 (d) 32

35. If the eccentricity of an ellipse be $1/\sqrt{2}$, then its latus rectum is equal to its
 (a) Minor axis (b) Semi-minor axis (c) Major axis (d) Semi-major axis
36. If the distance between a focus and corresponding directrix of an ellipse be 8 and the eccentricity be $1/2$, then the length of the minor axis is
 (a) 3 (b) $4\sqrt{2}$ (c) 6 (d) None of these
37. The sum of focal distances of any point on the ellipse with major and minor axes as $2a$ and $2b$ respectively, is equal to
 (a) $2a$ (b) $2\frac{a}{b}$ (c) $2\frac{b}{a}$ (d) $\frac{b^2}{a}$
38. P is any point on the ellipse $9x^2 + 36y^2 = 324$ whose foci are S and S' . Then $SP + S'P$ equals
 (a) 3 (b) 12 (c) 36 (d) 324
39. The foci of $16x^2 + 25y^2 = 400$ are
 (a) $(\pm 3, 0)$ (b) $(0, \pm 3)$ (c) $(3, -3)$ (d) $(-3, 3)$
40. In an ellipse $9x^2 + 5y^2 = 45$, the distance between the foci is
 (a) $4\sqrt{5}$ (b) $3\sqrt{5}$ (c) 3 (d) 4
41. The distance between the directrices of the ellipse $\frac{x^2}{36} + \frac{y^2}{20} = 1$ is
 (a) 8 (b) 12 (c) 18 (d) 24
42. If the eccentricity of the two ellipse $\frac{x^2}{169} + \frac{y^2}{25} = 1$, and $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ are equal, then the value of a/b is
 (a) $\frac{5}{13}$ (b) $\frac{6}{13}$ (c) $\frac{13}{5}$ (d) $\frac{13}{6}$
43. The equation of the ellipse whose one focus is at $(4, 0)$ and whose eccentricity is $4/5$, is
 (a) $\frac{x^2}{3^2} + \frac{y^2}{5^2} = 1$ (b) $\frac{x^2}{5^2} + \frac{y^2}{3^2} = 1$ (c) $\frac{x^2}{5^2} + \frac{y^2}{4^2} = 1$ (d) $\frac{x^2}{4^2} + \frac{y^2}{5^2} = 1$
44. S and T are the foci of an ellipse and B is an end of the minor axis. If STB is an equilateral triangle, the eccentricity of the ellipse is
 (a) $\frac{1}{4}$ (b) $\frac{1}{3}$ (c) $\frac{1}{2}$ (d) $\frac{2}{3}$
45. If C is the centre of the ellipse $9x^2 + 16y^2 = 144$ and S is one focus, the ratio of CS to semi-major axis, is
 (a) $\sqrt{7} : 16$ (b) $\sqrt{7} : 4$ (c) $\sqrt{5} : \sqrt{7}$ (d) None of these
46. If $LR = 10$, distance between foci = length of minor axis, then equation of ellipse is
 (a) $\frac{x^2}{50} + \frac{y^2}{100} = 1$ (b) $\frac{x^2}{100} + \frac{y^2}{50} = 1$ (c) $\frac{x^2}{50} + \frac{y^2}{20} = 1$ (d) None of these
47. Line joining foci subtends an angle of 90° at an extremity of minor axis, then eccentricity is
 (a) $\frac{1}{\sqrt{6}}$ (b) $\frac{1}{\sqrt{3}}$ (c) $\frac{1}{\sqrt{2}}$ (d) None of these

48. If foci are points $(0,1), (0,-1)$ and minor axis is of length 1, then equation of ellipse is
- (a) $\frac{x^2}{1/4} + \frac{y^2}{5/4} = 1$ (b) $\frac{x^2}{5/4} + \frac{y^2}{1/4} = 1$ (c) $\frac{x^2}{3/4} + \frac{y^2}{1/4} = 1$ (d) $\frac{x^2}{1/4} + \frac{y^2}{3/4} = 1$
49. The eccentricity of the ellipse $5x^2 + 9y^2 = 1$ is
- (a) $\frac{2}{3}$ (b) $\frac{3}{4}$ (c) $\frac{4}{5}$ (d) $\frac{1}{2}$
50. For the ellipse $x^2 + 4y^2 = 9$
- (a) The eccentricity is $\frac{1}{2}$ (b) The latus rectum is $\frac{2}{3}$ (c) A focus is $(3\sqrt{3}, 0)$ (d) A directrix is $x = 2\sqrt{3}$
51. The sum of the distances of any point on the ellipse $3x^2 + 4y^2 = 24$ from its foci is
- (a) $8\sqrt{2}$ (b) $4\sqrt{2}$ (c) $16\sqrt{2}$ (d) None of these
52. The sum of the focal distances from any point on the ellipse $9x^2 + 16y^2 = 144$ is
- (a) 32 (b) 18 (c) 16 (d) 8
53. The distance of a focus of the ellipse $9x^2 + 16y^2 = 144$ from an end of the minor axis is
- (a) $\frac{3}{2}$ (b) 3 (c) 4 (d) None of these
54. The equation of ellipse in the form $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, given the eccentricity to be $\frac{2}{3}$ and latus rectum $\frac{2}{3}$ is
- (a) $25x^2 + 45y^2 = 9$ (b) $25x^2 - 4y^2 = 9$ (c) $25x^2 - 45y^2 = 9$ (d) $25x^2 + 4y^2 = 1$
55. The equation of the ellipse with axes along the x -axis and the y -axis, which passes through the points $P(4, 3)$ and $Q(6, 2)$ is
- (a) $\frac{x^2}{50} + \frac{y^2}{13} = 1$ (b) $\frac{x^2}{52} + \frac{y^2}{13} = 1$ (c) $\frac{x^2}{13} + \frac{y^2}{52} = 1$ (d) $\frac{x^2}{52} + \frac{y^2}{17} = 1$
56. P is a variable point on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ with AA' as the major axis. Then the maximum value of the area of the triangle APA' is
- (a) ab (b) $2ab$ (c) $\frac{ab}{2}$ (d) None of these
57. The latus rectum of the ellipse $x^2 \tan^2 \alpha + y^2 \sec^2 \alpha = 1$ is $1/2$ then $\alpha (0 < \alpha < \pi)$ is equal to
- (a) $\pi/12$ (b) $\pi/6$ (c) $5\pi/12$ (d) None of these

Advance Level

58. An ellipse is described by using an endless string which is passed over two pins. If the axes are 6 cm and 4 cm, the necessary length of the string and the distance between the pins respectively in cm, are
- (a) $6, 2\sqrt{5}$ (b) $6, \sqrt{5}$ (c) $4, 2\sqrt{5}$ (d) None of these
59. A man running round a race-course notes that the sum of the distances of two flag-posts from him is always 10 meters and the distance between the flag-posts is 8 meters. The area of the path he encloses in square metres is
- (a) 15π (b) 12π (c) 18π (d) 8π

60. The equation $\frac{x^2}{1-r} - \frac{y^2}{1+r} = 1$, $r > 1$ represents
 (a) An ellipse (b) A hyperbola (c) A circle (d) An imaginary ellipse
61. The radius of the circle having its centre at (0,3) and passing through the foci of the ellipse $\frac{x^2}{16} + \frac{y^2}{9} = 1$, is
 (a) 3 (b) 3.5 (c) 4 (d) $\sqrt{12}$
62. The centre of an ellipse is C and PN is any ordinate and A, A' are the end points of major axis, then the value of $\frac{PN^2}{AN \cdot A'N}$ is
 (a) $\frac{b^2}{a^2}$ (b) $\frac{a^2}{b^2}$ (c) $a^2 + b^2$ (d) 1
63. Let P be a variable point on the ellipse $\frac{x^2}{25} + \frac{y^2}{16} = 1$ with foci at S and S' . If A be the area of triangle PSS' , then the maximum value of A is
 (a) 24 sq. units (b) 12 sq. units (c) 36 sq. units (d) None of these
64. The eccentricity of the ellipse which meets the straight line $\frac{x}{7} + \frac{y}{2} = 1$ on the axis of x and the straight line $\frac{x}{3} - \frac{y}{5} = 1$ on the axis of y and whose axes lie along the axes of coordinates, is
 (a) $\frac{3\sqrt{2}}{7}$ (b) $\frac{2\sqrt{6}}{7}$ (c) $\frac{\sqrt{3}}{7}$ (d) None of these
65. If the focal distance of an end of the minor axis of an ellipse (referred to its axes as the axes of x and y respectively) is k and the distance between its foci is $2h$, then its equation is
 (a) $\frac{x^2}{k^2} + \frac{y^2}{h^2} = 1$ (b) $\frac{x^2}{k^2} + \frac{y^2}{k^2 - h^2} = 1$ (c) $\frac{x^2}{k^2} + \frac{y^2}{h^2 - k^2} = 1$ (d) $\frac{x^2}{k^2} + \frac{y^2}{k^2 + h^2} = 1$
66. If (5, 12) and (24, 7) are the foci of a conic passing through the origin, then the eccentricity of conic is
 (a) $\frac{\sqrt{386}}{38}$ (b) $\frac{\sqrt{386}}{12}$ (c) $\frac{\sqrt{386}}{13}$ (d) $\frac{\sqrt{386}}{25}$
67. The maximum area of an isosceles triangle inscribed in the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ with the vertex at one end of the major axis is
 (a) $\sqrt{3}ab$ (b) $\frac{3\sqrt{3}}{4}ab$ (c) $\frac{5\sqrt{3}}{4}ab$ (d) None of these
68. The radius of the circle passing through the foci of the ellipse $\frac{x^2}{16} + \frac{y^2}{9} = 1$ and having its centre (0, 3) is
 (a) 4 (b) 3 (c) $\sqrt{12}$ (d) $\frac{7}{2}$

69. The locus of extremities of the latus rectum of the family of ellipse $b^2x^2 + y^2 = a^2b^2$ is
 (a) $x^2 - ay = a^2$ (b) $x^2 - ay = b^2$ (c) $x^2 + ay = a^2$ (d) $x^2 + ay = b^2$

SPECIAL FORM OF AN ELLIPSE , PARAMETRIC EQUATION OF AN ELLIPSE

Basic Level

70. The equation of the ellipse whose centre is (2, -3), one of the foci is (3, -3) and the corresponding vertex is
 (a) $\frac{(x-2)^2}{3} + \frac{(y+3)^2}{4} = 1$ (b) $\frac{(x-2)^2}{4} + \frac{(y+3)^2}{3} = 1$ (c) $\frac{x^2}{3} + \frac{y^2}{4} = 1$ (d) None of these
71. The equation of an ellipse, whose vertices are (2, -2), (2, 4) and eccentricity $\frac{1}{3}$, is
 (a) $\frac{(x-2)^2}{9} + \frac{(y-1)^2}{8} = 1$ (b) $\frac{(x-2)^2}{8} + \frac{(y-1)^2}{9} = 1$ (c) $\frac{(x+2)^2}{8} + \frac{(y+1)^2}{9} = 1$ (d) $\frac{(x-2)^2}{9} + \frac{(y+1)^2}{8} = 1$
72. The equation of an ellipse whose eccentricity $1/2$ is and the vertices are (4, 0) and (10, 0) is
 (a) $3x^2 + 4y^2 - 42x + 120 = 0$ (b) $3x^2 + 4y^2 + 42x + 120 = 0$
 (c) $3x^2 + 4y^2 + 42x - 120 = 0$ (d) $3x^2 + 4y^2 - 42x - 120 = 0$
73. For the ellipse $3x^2 + 4y^2 - 6x + 8y - 5 = 0$
 (a) Centre is (2, -1) (b) Eccentricity is $\frac{1}{3}$
 (c) Foci are (3, 1) and (-1, 1) (d) Centre is (1, -1), $e = \frac{1}{2}$, foci are (3, -1) and (-1, -1)
74. The eccentricity of the ellipse $9x^2 + 5y^2 - 18x - 2y - 16 = 0$
 (a) $1/2$ (b) $2/3$ (c) $1/3$ (d) $3/4$
75. The eccentricity of the ellipse $\frac{(x-1)^2}{9} + \frac{(y+1)^2}{25} = 1$ is
 (a) $4/5$ (b) $3/5$ (c) $5/4$ (d) Imaginary
76. The eccentricity of the ellipse $9x^2 + 5y^2 - 30y = 0$, is
 (a) $\frac{1}{3}$ (b) $\frac{2}{3}$ (c) $\frac{3}{4}$ (d) None of these
77. The eccentricity of the ellipse $4x^2 + 9y^2 + 8x + 36y + 4 = 0$ is
 (a) $\frac{5}{6}$ (b) $\frac{3}{5}$ (c) $\frac{\sqrt{2}}{3}$ (d) $\frac{\sqrt{5}}{3}$
78. The eccentricity of the curve represented by the equation $x^2 + 2y^2 - 2x + 3y + 2 = 0$ is
 (a) 0 (b) $1/2$ (c) $1/\sqrt{2}$ (d) $\sqrt{2}$
79. The centre of the ellipse $\frac{(x+y-2)^2}{9} + \frac{(x-y)^2}{16} = 1$, is
 (a) (0, 0) (b) (1, 1) (c) (1, 0) (d) (0, 1)
80. The centre of the ellipse $4x^2 + 9y^2 - 16x - 54y + 61 = 0$ is
 (a) (1, 3) (b) (2, 3) (c) (3, 2) (d) (3, 1)
81. Latus rectum of ellipse $4x^2 + 9y^2 - 8x - 36y + 4 = 0$ is
 (a) $8/3$ (b) $4/3$ (c) $\frac{\sqrt{5}}{3}$ (d) $16/3$

82. The length of the axes of the conic $9x^2 + 4y^2 - 6x + 4y + 1 = 0$, are
 (a) $\frac{1}{2}, 9$ (b) $3, \frac{2}{5}$ (c) $1, \frac{2}{3}$ (d) $3, 2$
83. Equations $x = a \cos \theta, y = b \sin \theta (a > b)$ represent a conic section whose eccentricity e is given by
 (a) $e^2 = \frac{a^2 + b^2}{a^2}$ (b) $e^2 = \frac{a^2 + b^2}{b^2}$ (c) $e^2 = \frac{a^2 - b^2}{a^2}$ (d) $e^2 = \frac{a^2 - b^2}{b^2}$
84. The curve with parametric equations $x = 1 + 4 \cos \theta, y = 2 + 3 \sin \theta$ is
 (a) An ellipse (b) A parabola (c) A hyperbola (d) A circle
85. The equations $x = a \cos \theta, y = b \sin \theta, 0 \leq \theta < 2\pi, a \neq b$, represent
 (a) An ellipse (b) A parabola (c) A circle (d) A hyperbola
86. The curve represented by $x = 2(\cos t + \sin t), y = 5(\cos t - \sin t)$ is
 (a) A circle (b) A parabola (c) An ellipse (d) A hyperbola
87. The equations $x = a \left(\frac{1 - t^2}{1 + t^2} \right), y = \frac{2bt}{1 + t^2}; t \in R$ represent
 (a) A circle (b) An ellipse (c) A parabola (d) A hyperbola
88. The eccentricity of the ellipse represented by $25x^2 + 16y^2 - 150x - 175 = 0$ is
 (a) $\frac{2}{5}$ (b) $\frac{3}{5}$ (c) $\frac{4}{5}$ (d) None of these
89. The set of values of a for which $(13x - 1)^2 + (13y - 2)^2 = a(5x + 12y - 1)^2$ represents an ellipse is
 (a) $1 < a < 2$ (b) $0 < a < 1$ (c) $2 < a < 3$ (d) None of these

Advance Level

90. The parametric representation of a point on the ellipse whose foci are $(-1, 0)$ and $(7, 0)$ and eccentricity $1/2$ is
 (a) $(3 + 8 \cos \theta, 4\sqrt{3} \sin \theta)$ (b) $(8 \cos \theta, 4\sqrt{3} \sin \theta)$ (c) $(3 + 4\sqrt{3} \cos \theta, 8 \sin \theta)$ (d) None of these
91. If $P(\theta)$ and $Q\left(\frac{\pi}{2} + \theta\right)$ are two points on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, then locus of the mid-point of PQ is
 (a) $\frac{x^2}{a^2} + \frac{y^2}{b^2} = \frac{1}{2}$ (b) $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 4$ (c) $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 2$ (d) None of these

POSITION OF A POINT, TANGENTS, PAIR OF TANGENTS & DIRECTOR CIRCLE OF AN ELLIPSE

Basic Level

92. The line $lx + my - n = 0$ will be a tangent to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, if
 (a) $a^2 l^2 + b^2 m^2 = n^2$ (b) $al^2 + bm^2 = n^2$ (c) $a^2 l + b^2 m = n$ (d) None of these
93. The line $x \cos \alpha + y \sin \alpha = p$ will be a tangent to the conic $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, if
 (a) $p^2 = a^2 \sin^2 \alpha + b^2 \cos^2 \alpha$ (b) $p^2 = a^2 + b^2$
 (c) $p^2 = b^2 \sin^2 \alpha + a^2 \cos^2 \alpha$ (d) None of these

94. The equations of the tangents of the ellipse $9x^2 + 16y^2 = 144$, which passes through the point (2, 3) is
 (a) $y = 3, x + y = 5$ (b) $y = -3, x - y = 5$ (c) $y = 4, x + y = 3$ (d) $y = -4, x - y = 3$
95. The equation of the tangent to the conic $x^2 - y^2 - 8x + 2y + 11 = 0$ at (2, 1) is
 (a) $x + 2 = 0$ (b) $2x + 1 = 0$ (c) $x - 2 = 0$ (d) $x + y + 1 = 0$
96. The position of the point (1, 3) with respect to the ellipse $4x^2 + 9y^2 - 16x - 54y + 61 = 0$ is
 (a) Outside the ellipse (b) On the ellipse (c) On the major axis (d) On the minor axis
97. The ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and the straight line $y = mx + c$ intersect in real points only if
 (a) $a^2 m^2 < c^2 - b^2$ (b) $a^2 m^2 > c^2 - b^2$ (c) $a^2 m^2 \geq c^2 - b^2$ (d) $c \geq b$
98. If the line $y = mx + c$ touches the ellipse $\frac{x^2}{b^2} + \frac{y^2}{a^2} = 1$, then $c =$
 (a) $\pm \sqrt{b^2 m^2 + a^2}$ (b) $\pm \sqrt{a^2 m^2 + b^2}$ (c) $\pm \sqrt{b^2 m^2 - a^2}$ (d) $\pm \sqrt{a^2 m^2 - b^2}$
99. If the line $y = 2x + c$ be a tangent to the ellipse $\frac{x^2}{8} + \frac{y^2}{4} = 1$, then $c =$
 (a) ± 4 (b) ± 6 (c) ± 1 (d) ± 8
100. The equation of the tangent to the ellipse $x^2 + 16y^2 = 16$ making an angle of 60° with x -axis
 (a) $\sqrt{3}x - y + 7 = 0$ (b) $\sqrt{3}x - y - 7 = 0$ (c) $\sqrt{3}x - y \pm 7 = 0$ (d) None of these
101. The position of the point (4, -3) with respect to the ellipse $2x^2 + 5y^2 = 20$ is
 (a) Outside the ellipse (b) On the ellipse (c) On the major axis (d) None of these
102. The angle between the pair of tangents drawn to the ellipse $3x^2 + 2y^2 = 5$ from the point (1, 2) is
 (a) $\tan^{-1}\left(\frac{12}{5}\right)$ (b) $\tan^{-1}(6\sqrt{5})$ (c) $\tan^{-1}\left(\frac{12}{\sqrt{5}}\right)$ (d) $\tan^{-1}(12\sqrt{5})$
103. If any tangent to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ cuts off intercepts of length h and k on the axes, then
 $\frac{a^2}{h^2} + \frac{b^2}{k^2} =$
 (a) 0 (b) 1 (c) -1 (d) None of these
104. The equation of the tangents drawn at the ends of the major axis of the ellipse $9x^2 + 5y^2 - 30y = 0$, are
 (a) $y = \pm 3$ (b) $x = \pm \sqrt{5}$ (c) $y = 0, y = 6$ (d) None of these
105. The locus of the point of intersection of mutually perpendicular tangent to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, is
 (a) A straight line (b) A parabola (c) A circle (d) None of these
106. Two perpendicular tangents drawn to the ellipse $\frac{x^2}{25} + \frac{y^2}{16} = 1$ intersect on the curve
 (a) $x = \frac{a}{e}$ (b) $x^2 + y^2 = 41$ (c) $x^2 + y^2 = 9$ (d) $x^2 - y^2 = 41$

107. The product of the perpendiculars drawn from the two foci of an ellipse to the tangent at any point of the ellipse is
- (a) a^2 (b) b^2 (c) $4a^2$ (d) $4b^2$
108. The equations of the tangents to the ellipse $4x^2 + 3y^2 = 5$, which are inclined at 60° to the axis of x are
- (a) $y = \sqrt{3}x \pm \sqrt{\frac{65}{12}}$ (b) $y = \sqrt{3}x \pm \sqrt{\frac{12}{65}}$ (c) $y = \frac{x}{\sqrt{3}} \pm \sqrt{\frac{65}{12}}$ (d) None of these
109. If the straight line $y = 4x + c$ is a tangent to the ellipse $\frac{x^2}{8} + \frac{y^2}{4} = 1$, then c will be equal to
- (a) ± 4 (b) ± 6 (c) ± 1 (d) $\pm \sqrt{132}$
110. Tangents are drawn to the ellipse $3x^2 + 5y^2 = 32$ and $25x^2 + 9y^2 = 450$ passing through the point $(3, 5)$. The number of such tangents are
- (a) 2 (b) 3 (c) 4 (d) 0
111. If $\frac{x}{a} + \frac{y}{b} = \sqrt{2}$ touches the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, then its eccentric angle θ is equal to
- (a) 0° (b) 90° (c) 45° (d) 60°
112. Locus of point of intersection of tangents at $(a\cos\alpha, b\sin\alpha)$ and $(a\cos\beta, b\sin\beta)$ for the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is
- (a) A circle (b) A straight line (c) An ellipse (d) A parabola
113. The equation of the tangent at the point $(1/4, 1/4)$ of the ellipse $\frac{x^2}{4} + \frac{y^2}{12} = 1$ is
- (a) $3x + y = 48$ (b) $3x + y = 3$ (c) $3x + y = 16$ (d) None of these
114. If F_1 and F_2 be the feet of the perpendiculars from the foci S_1 and S_2 of an ellipse $\frac{x^2}{5} + \frac{y^2}{3} = 1$ on the tangent at any point P on the ellipse, then $(S_1F_1)(S_2F_2)$ is equal to
- (a) 2 (b) 3 (c) 4 (d) 5
115. Equations of tangents to the ellipse $\frac{x^2}{9} + \frac{y^2}{4} = 1$, which cut off equal intercepts on the axes is
- (a) $y = x + \sqrt{13}$ (b) $y = -x + \sqrt{13}$ (c) $y = x - \sqrt{13}$ (d) $y = -x - \sqrt{13}$
116. The line $x = at^2$ meets the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ in the real points, if
- (a) $|t| < 2$ (b) $|t| \leq 1$ (c) $|t| > 1$ (d) None of these

Advance Level

117. The locus of mid points of parts in between axes and tangents of ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ will be
- (a) $\frac{a^2}{x^2} + \frac{b^2}{y^2} = 1$ (b) $\frac{a^2}{x^2} + \frac{b^2}{y^2} = 2$ (c) $\frac{a^2}{x^2} + \frac{b^2}{y^2} = 3$ (d) $\frac{a^2}{x^2} + \frac{b^2}{y^2} = 4$
118. The angle of intersection of ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and circle $x^2 + y^2 = ab$ is
- (a) $\tan^{-1}\left(\frac{a-b}{ab}\right)$ (b) $\tan^{-1}\left(\frac{a+b}{ab}\right)$ (c) $\tan^{-1}\left(\frac{a+b}{\sqrt{ab}}\right)$ (d) $\tan^{-1}\left(\frac{a-b}{\sqrt{ab}}\right)$
119. Locus of the foot of the perpendicular drawn from the centre upon any tangent to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is
- (a) $(x^2 + y^2)^2 = b^2x^2 + a^2y^2$ (b) $(x^2 + y^2)^2 = b^2x^2 - a^2y^2$
(c) $(x^2 + y^2)^2 = a^2x^2 - b^2y^2$ (d) $(x^2 + y^2)^2 = a^2x^2 + b^2y^2$
120. If a tangent having slope of $-\frac{4}{3}$ to the ellipse $\frac{x^2}{18} + \frac{y^2}{32} = 1$ intersects the major and minor axes in points A and B respectively, then the area of $\triangle OAB$ is equal to (O is centre of the ellipse)
- (a) 12 sq. units (b) 48 sq. units (c) 64 sq. units (d) 24 sq. units
121. Tangent is drawn to ellipse $\frac{x^2}{27} + y^2 = 1$ at $(3\sqrt{3}\cos\theta, \sin\theta)$ (where $\theta \in \left(0, \frac{\pi}{2}\right)$). Then the value of θ such that sum of intercepts on axes made by this tangent is minimum, is
- (a) $\pi/3$ (b) $\pi/6$ (c) $\pi/8$ (d) $\pi/4$
122. If the tangent at the point $\left(4\cos\phi, \frac{16}{\sqrt{11}}\sin\phi\right)$ to the ellipse $16x^2 + 11y^2 = 256$ is also a tangent to the circle $x^2 + y^2 - 2x = 15$, then the value of ϕ is
- (a) $\pm \frac{\pi}{2}$ (b) $\pm \frac{\pi}{4}$ (c) $\pm \frac{\pi}{3}$ (d) $\pm \frac{\pi}{6}$
123. An ellipse passes through the point (4, -1) and its axes are along the axes of co-ordinates. If the line $x + 4y - 10 = 0$ is a tangent to it, then its equation is
- (a) $\frac{x^2}{100} + \frac{y^2}{5} = 1$ (b) $\frac{x^2}{80} + \frac{y^2}{5/4} = 1$ (c) $\frac{x^2}{20} + \frac{y^2}{5} = 1$ (d) None of these
124. The sum of the squares of the perpendiculars on any tangent to the ellipse $x^2/a^2 + y^2/b^2 = 1$ from two points on the minor axis each distance $\sqrt{a^2 - b^2}$ from the centre is
- (a) a^2 (b) b^2 (c) $2a^2$ (d) $2b^2$
125. The tangent at a point $P(a\cos\theta, b\sin\theta)$ of an ellipse $x^2/a^2 + y^2/b^2 = 1$, meets its auxiliary circle in two points, the chord joining which subtends a right angle at the centre, then the eccentricity of the ellipse is
- (a) $(1 + \sin^2\theta)^{-1}$ (b) $(1 + \sin^2\theta)^{-1/2}$ (c) $(1 + \sin^2\theta)^{-3/2}$ (d) $(1 + \sin^2\theta)^{-2}$

126. The locus of the point of intersection of tangents to an ellipse at two points, sum of whose eccentric angles is constant is
 (a) A parabola (b) A circle (c) An ellipse (d) A straight line
127. The sum of the squares of the perpendiculars on any tangents to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ from two points on the minor axis each at a distance ae from the centre is
 (a) $2a^2$ (b) $2b^2$ (c) $a^2 + b^2$ (d) $a^2 - b^2$
128. The equation of the circle passing through the points of intersection of ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and $\frac{x^2}{b^2} + \frac{y^2}{a^2} = 1$ is
 (a) $x^2 + y^2 = a^2$ (b) $x^2 + y^2 = b^2$ (c) $x^2 + y^2 = \frac{a^2 b^2}{a^2 + b^2}$ (d) $x^2 + y^2 = \frac{2a^2 b^2}{a^2 + b^2}$
129. The slope of a common tangent to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and a concentric circle of radius r is
 (a) $\tan^{-1} \sqrt{\frac{r^2 - b^2}{a^2 - r^2}}$ (b) $\sqrt{\frac{r^2 - b^2}{a^2 - r^2}}$ (c) $\left(\frac{r^2 - b^2}{a^2 - r^2} \right)$ (d) $\sqrt{\frac{a^2 - r^2}{r^2 - b^2}}$
130. The tangents from which of the following points to the ellipse $5x^2 + 4y^2 = 20$ are perpendicular
 (a) $(1, 2\sqrt{2})$ (b) $(2\sqrt{2}, 1)$ (c) $(2, \sqrt{5})$ (d) $(\sqrt{5}, 2)$

NORMAL , ECCENTRIC ANGLES OF CO – ORDINATE POINTS

Basic Level

131. The line $y = mx + c$ is a normal to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, if $c =$
 (a) $-(2am + bn^2)$ (b) $\frac{(a^2 + b^2)m}{\sqrt{a^2 + b^2 m^2}}$ (c) $-\frac{(a^2 - b^2)m}{\sqrt{a^2 + b^2 m^2}}$ (d) $\frac{(a^2 - b^2)m}{\sqrt{a^2 + b^2}}$
132. The line $lx + my + n = 0$ is a normal to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, if
 (a) $\frac{a^2}{m^2} + \frac{b^2}{l^2} = \frac{(a^2 - b^2)}{n^2}$ (b) $\frac{a^2}{l^2} + \frac{b^2}{m^2} = \frac{(a^2 - b^2)^2}{n^2}$ (c) $\frac{a^2}{l^2} - \frac{b^2}{m^2} = \frac{(a^2 - b^2)^2}{n^2}$ (d) None of these
133. If the line $x \cos \alpha + y \sin \alpha = p$ be a normal to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, then
 (a) $p^2(a^2 \cos^2 \alpha + b^2 \sin^2 \alpha) = a^2 - b^2$ (b) $p^2(a^2 \cos^2 \alpha + b^2 \sin^2 \alpha) = (a^2 - b^2)^2$
 (c) $p^2(a^2 \sec^2 \alpha + b^2 \csc^2 \alpha) = a^2 - b^2$ (d) $p^2(a^2 \sec^2 \alpha + b^2 \csc^2 \alpha) = (a^2 - b^2)^2$
134. The equation of the normal at the point $(2, 3)$ on the ellipse $9x^2 + 16y^2 = 180$, is
 (a) $3y = 8x - 10$ (b) $3y - 8x + 7 = 0$ (c) $8y + 3x + 7 = 0$ (d) $3x + 2y + 7 = 0$

135. The eccentric angles of the extremities of latus-rectum of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ are given by
 (a) $\tan^{-1}\left(\pm \frac{ae}{b}\right)$ (b) $\tan^{-1}\left(\pm \frac{be}{a}\right)$ (c) $\tan^{-1}\left(\pm \frac{b}{ae}\right)$ (d) $\tan^{-1}\left(\pm \frac{a}{be}\right)$
136. The number of normals that can be drawn from a point to a given ellipse is
 (a) 2 (b) 3 (c) 4 (d) 1
137. The eccentric angle of a point on the ellipse $\frac{x^2}{6} + \frac{y^2}{2} = 1$, whose distances from the centre of the ellipse is 2, is
 (a) $\frac{\pi}{4}$ (b) $\frac{3\pi}{2}$ (c) $\frac{5\pi}{3}$ (d) $\frac{7\pi}{6}$

Advance Level

138. If the normal at the point $P(\theta)$ to the ellipse $\frac{x^2}{14} + \frac{y^2}{5} = 1$ intersects it again at the point $Q(2\theta)$, then $\cos\theta$ is equal to
 (a) $\frac{2}{3}$ (b) $-\frac{2}{3}$ (c) $\frac{3}{2}$ (d) $-\frac{3}{2}$
139. If the normal at any point P on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ meets the coordinates axes in G and g respectively, then $PG : Pg =$
 (a) $a : b$ (b) $a^2 : b^2$ (c) $b^2 : a^2$ (d) $b : a$
140. If α and β are eccentric angles of the ends of a focal chord of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, then $\tan\frac{\alpha}{2} \tan\frac{\beta}{2}$ is equal to
 (a) $\frac{1-e}{1+e}$ (b) $\frac{e-1}{e+1}$ (c) $\frac{e+1}{e-1}$ (d) None of these
141. If the normal at one end of the latus-rectum of an ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ passes through the one end of the minor axis, then
 (a) $e^4 - e^2 + 1 = 0$ (b) $e^2 - e + 1 = 0$ (c) $e^2 + e + 1 = 0$ (d) $e^4 + e^2 - 1 = 0$
142. The line $2x + y = 3$ cuts the ellipse $4x^2 + y^2 = 5$ at P and Q . If θ be the angle between the normals at these points, then $\tan\theta =$
 (a) $1/2$ (b) $3/4$ (c) $3/5$ (d) 5
143. The eccentric angles of extremities of a chord of an ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ are θ_1 and θ_2 . If this chord passes through the focus, then
 (a) $\tan\frac{\theta_1}{2} \tan\frac{\theta_2}{2} + \frac{1-e}{1+e} = 0$ (b) $\cos\frac{\theta_1 - \theta_2}{2} = e \cos\frac{\theta_1 + \theta_2}{2}$
 (c) $e = \frac{\sin\theta_1 + \sin\theta_2}{\sin(\theta_1 + \theta_2)}$ (d) $\cot\frac{\theta_1}{2} \cdot \cot\frac{\theta_2}{2} = \frac{e+1}{e-1}$

144. Let F_1, F_2 be two foci of the ellipse and PT and PN be the tangent and the normal respectively to the ellipse at point P then
- (a) PN bisects $\angle F_1PF_2$ (b) PT bisects $\angle F_1PF_2$
 (c) PT bisects angle $(180^\circ - \angle F_1PF_2)$ (d) None of these
145. If CF is the perpendicular from the centre C of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ on the tangent at any point P and G is the point when the normal at P meets the major axis, then $CF \cdot PG =$
- (a) a^2 (b) ab (c) b^2 (d) b^3

CHORD OF CONTACT , EQUATION OF CHORD JOINING TWO POINTS OF AN ELLIPSE

Basic Level

146. The equation of the chord of the ellipse $2x^2 + 5y^2 = 20$ which is bisected at the point $(2, 1)$ is
- (a) $4x + 5y + 13 = 0$ (b) $4x + 5y = 13$ (c) $5x + 4y + 13 = 0$ (d) None of these
147. If the chords of contact of tangents from two points (x_1, y_1) and (x_2, y_2) to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ are at right angles, then $\frac{x_1 x_2}{y_1 y_2}$ is equal to
- (a) $\frac{a^2}{b^2}$ (b) $-\frac{b^2}{a^2}$ (c) $-\frac{a^4}{b^4}$ (d) $-\frac{b^4}{a^4}$
148. Chords of an ellipse are drawn through the positive end of the minor axis. Then their mid-point lies on
- (a) A circle (b) A parabola (c) An ellipse (d) A hyperbola
149. The length of the common chord of the ellipse $\frac{(x-1)^2}{9} + \frac{(y-2)^2}{4} = 1$ and the circle $(x-1)^2 + (y-2)^2 = 1$ is
- (a) Zero (b) One (c) Three (d) Eight
150. If $\tan \theta_1 \tan \theta_2 = -\frac{a^2}{b^2}$, then the chord joining two points θ_1 and θ_2 on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ will subtend a right angle at
- (a) Focus (b) Centre (c) End of the major axis (d) End of the minor axis
151. If θ and ϕ are the eccentric angles of the ends of a focal chord of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, then
- (a) $\cos \frac{\theta - \phi}{2} = e \cos \frac{\theta + \phi}{2}$ (b) $\cos \frac{\theta - \phi}{2} + e \cos \frac{\theta + \phi}{2} = 0$ (c) $\cos \frac{\theta + \phi}{2} = e \cos \frac{\theta - \phi}{2}$ (d) None of these

DIAMETER OF ELLIPSE , POLE AND POLAR & CONJUGATE DIAMETER

Basic Level

152. With respect to the ellipse $3x^2 + 2y^2 = 1$, the pole of the line $9x + 2y = 1$ is
(a) $(-1, -3)$ (b) $(-1, 3)$ (c) $(3, -1)$ (d) $(3, 1)$
153. In the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, the equation of diameter conjugate to the diameter $y = \frac{b}{a}x$ is
(a) $y = -\frac{b}{a}x$ (b) $y = -\frac{a}{b}x$ (c) $x = -\frac{b}{a}y$ (d) None of these
154. If CP and CD are semi conjugate diameters of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, then $CP^2 + CD^2 =$
(a) $a + b$ (b) $a^2 + b^2$ (c) $a^2 - b^2$ (d) $\sqrt{a^2 + b^2}$
155. The eccentricity of an ellipse whose pair of a conjugate diameter are $y = x$ and $3y = -2x$ is
(a) $2/3$ (b) $1/3$ (c) $1/\sqrt{3}$ (d) None of these
156. If eccentric angle of one diameter is $\frac{5\pi}{6}$, then eccentric angle of conjugate diameter is
(a) $\frac{2\pi}{3}$ (b) $\frac{4\pi}{3}$ (c) $\frac{2\pi}{3}$ or $\frac{4\pi}{3}$ (d) None of these
157. For the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, the equation of the diameter conjugate to $ax - by = 0$ is
(a) $bx + ay = 0$ (b) $bx - ay = 0$ (c) $a^3y + b^3x = 0$ (d) $a^3y - b^3x = 0$
158. Equation of equi-conjugate diameter for an ellipse $\frac{x^2}{25} + \frac{y^2}{16} = 1$ is
(a) $x = \pm \frac{5}{4}y$ (b) $y = \pm \frac{5}{4}x$ (c) $x = \pm \frac{25}{16}y$ (d) None of these

Advance Level

159. The locus of the point of intersection of tangents at the ends of semi-conjugate diameter of ellipse is
(a) Parabola (b) Hyperbola (c) Circle (d) Ellipse
160. AB is a diameter of $x^2 + 9y^2 = 25$. The eccentric angle of A is $\pi/6$. Then the eccentric angle of B is
(a) $5\pi/6$ (b) $-5\pi/6$ (c) $-2\pi/3$ (d) None of these
161. If the points of intersection of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and $\frac{x^2}{p^2} + \frac{y^2}{q^2} = 1$ be the extremities of the conjugate diameter of first ellipse, then
(a) $\frac{x^2}{p^2} + \frac{y^2}{q^2} = 2$ (b) $\frac{a^2}{p^2} + \frac{b^2}{q^2} = 1$ (c) $\frac{a}{p} + \frac{b}{q} = 1$ (d) $\frac{a^2}{p^2} + \frac{b^2}{q^2} = 2$

ANSWER

BASIC & ADVANCE LEVEL

| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 |
|-----|-----|-----------------|-----|-----|-----|-----|-----|-----|-----------------|-----|-----|-----|-----|-----------------|-----|-----|-----|-----|-----|
| c | a | a | b | b | a | b | b | b | a | d | b | b | a | a | c | b | a | a | c |
| 21 | 22 | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 | 34 | 35 | 36 | 37 | 38 | 39 | 40 |
| c | b | a | a | b | d | b | b | b | b | b | c | b | d | d | d | a | b | a | d |
| 41 | 42 | 43 | 44 | 45 | 46 | 47 | 48 | 49 | 50 | 51 | 52 | 53 | 54 | 55 | 56 | 57 | 58 | 59 | 60 |
| c | c | b | c | b | b | c | a | a | d | b | d | c | a | b | a | a,c | d | a | d |
| 61 | 62 | 63 | 64 | 65 | 66 | 67 | 68 | 69 | 70 | 71 | 72 | 73 | 74 | 75 | 76 | 77 | 78 | 79 | 80 |
| c | a | b | b | b | a,b | b | a | a,c | b | b | a | d | b | a | b | d | c | b | b |
| 81 | 82 | 83 | 84 | 85 | 86 | 87 | 88 | 89 | 90 | 91 | 92 | 93 | 94 | 95 | 96 | 97 | 98 | 99 | 100 |
| a | c | c | a | a | c | b | b | b | a | a | a | c | a | c | c | c | a | b | c |
| 101 | 102 | 103 | 104 | 105 | 106 | 107 | 108 | 109 | 110 | 111 | 112 | 113 | 114 | 115 | 116 | 117 | 118 | 119 | 120 |
| a | c | b | c | c | b | b | a | d | b | c | c | d | b | a,b ,c, d | b | d | d | d | d |
| 121 | 122 | 123 | 124 | 125 | 126 | 127 | 128 | 129 | 130 | 131 | 132 | 133 | 134 | 135 | 136 | 137 | 138 | 139 | 140 |
| b | c | b,c | c | b | d | a | d | b | a,b ,c, d | c | b | d | b | c | c | a | b | c | b |
| 141 | 142 | 143 | 144 | 145 | 146 | 147 | 148 | 149 | 150 | 151 | 152 | 153 | 154 | 155 | 156 | 157 | 158 | 159 | 160 |
| d | c | a,b ,c, d | a,c | c | b | c | c | a | b | a | d | a | b | c | c | c | a | d | b |
| 161 | | | | | | | | | | | | | | | | | | | |
| d | | | | | | | | | | | | | | | | | | | |