


# HYPERBOLA

## 5.3.1 DEFINITION

A hyperbola is the locus of a point in a plane which moves in the plane in such a way that the ratio of its distance from a fixed point in the same plane to its distance from a fixed line is always constant which is always greater than unity.

Fixed point is called focus, fixed straight line is called directrix and the constant ratio is called eccentricity of the hyperbola. Eccentricity is denoted by  $e$  and  $e > 1$ .



A hyperbola is the particular case of the conic

$$ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$$

When,  $abc + 2fgh - af^2 - bg^2 - ch^2 \neq 0$  i.e.,  $\Delta \neq 0$  and  $h^2 > ab$ .

Let  $S(h, k)$  is the focus, directrix is the line  $ax + by + c = 0$  and the eccentricity is  $e$ . Let  $P(x_1, y_1)$  be a point which moves such that  $SP = e \cdot PM$

$$\Rightarrow \sqrt{(x_1 - h)^2 + (y_1 - k)^2} = e \cdot \frac{ax_1 + by_1 + c}{\sqrt{a^2 + b^2}}$$

$$\Rightarrow (a^2 + b^2)[(x_1 - h)^2 + (y_1 - k)^2] = e^2(ax_1 + by_1 + c)^2$$

Hence, locus of  $(x_1, y_1)$  is given by  $(a^2 + b^2)[(x - h)^2 + (y - k)^2] = e^2(ax + by + c)^2$

Which is a second degree equation to represent a hyperbola ( $e > 1$ ).

**Example: 1** The equation of the conic with focus at  $(1, -1)$ , directrix along  $x - y + 1 = 0$  and with eccentricity  $\sqrt{2}$  is

(a)  $x^2 - y^2 = 1$

(b)  $xy = 1$

(c)  $2xy - 4x + 4y + 1 = 0$

(d)  $2xy + 4x - 4y - 1 = 0$

**Solution:** (c) Here, focus  $(S) = (1, -1)$ , eccentricity  $(e) = \sqrt{2}$

From definition,  $SP = e \cdot PM$

$$\sqrt{(x-1)^2 + (y+1)^2} = \frac{\sqrt{2} \cdot (x - y + 1)}{\sqrt{1^2 + 1^2}}$$

$$\Rightarrow (x-1)^2 + (y+1)^2 = (x - y + 1)^2 \Rightarrow 2xy - 4x + 4y + 1 = 0, \text{ which is the required equation of conic}$$

(Rectangular hyperbola)

**Example: 2** The centre of the hyperbola  $9x^2 - 36x - 16y^2 + 96y - 252 = 0$  is

(a)  $(2, 3)$

(b)  $(-2, -3)$

(c)  $(-2, 3)$

(d)  $(2, -3)$

**Solution:** (a) Here  $a = 9$ ,  $b = -16$ ,  $h = 0$ ,  $g = -18$ ,  $f = 48$ ,  $c = -252$

$$\text{Centre of hyperbola} = \left( \frac{hf - bg}{ab - h^2}, \frac{gh - af}{ab - h^2} \right) = \left( \frac{(0)(48) - (-16)(-18)}{(9)(-16) - 0}, \frac{(-18)(0) - (9)(48)}{(9)(-16) - 0} \right) = (2, 3)$$


## 5.3.2 STANDARD EQUATION OF THE HYPERBOLA

Let  $S$  be the focus,  $ZM$  be the directrix and  $e$  be the eccentricity of the hyperbola, then by definition,

$$\Rightarrow \frac{SP}{PM} = e \Rightarrow (SP)^2 = e^2(PM)^2$$

$$\Rightarrow (x - ae)^2 + (y - 0)^2 = e^2 \left( x - \frac{a}{e} \right)^2$$

$$\Rightarrow \frac{x^2}{a^2} - \frac{y^2}{a^2(e^2 - 1)} = 1 \Rightarrow \frac{x^2}{a^2} - \frac{y^2}{b^2} = 1, \text{ where } b^2 = a^2(e^2 - 1)$$



This is the standard equation of the hyperbola.

**Some terms related to hyperbola :** Let the equation of hyperbola is  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$

(1) **Centre :** All chords passing through  $C$  are bisected at  $C$ . Here  $C(0,0)$

(2) **Vertex:** The point  $A$  and  $A'$  where the curve meets the line joining the foci  $S$  and  $S'$  are called vertices of hyperbola. The co-ordinates of  $A$  and  $A'$  are  $(a, 0)$  and  $(-a, 0)$  respectively.

(3) **Transverse and conjugate axes :** The straight line joining the vertices  $A$  and  $A'$  is called transverse axis of the hyperbola. The straight line perpendicular to the transverse axis and passing through the centre is called conjugate axis.

Here, transverse axis =  $AA' = 2a$

Conjugate axis =  $BB' = 2b$

(4) **Eccentricity :** For the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$

$$\text{We have } b^2 = a^2(e^2 - 1), \quad e = \sqrt{1 + \left( \frac{2b}{2a} \right)^2} = \sqrt{1 + \left( \frac{\text{Conjugate axis}}{\text{Transverse axis}} \right)^2}$$

(5) **Double ordinates :** If  $Q$  be a point on the hyperbola,  $QN$  perpendicular to the axis of the hyperbola and produced to meet the curve again at  $Q'$ . Then  $QQ'$  is called a double ordinate at  $Q$ .

If abscissa of  $Q$  is  $h$ , then co-ordinates of  $Q$  and  $Q'$  are  $\left( h, \frac{b}{a} \sqrt{h^2 - a^2} \right)$  and  $\left( h, -\frac{b}{a} \sqrt{h^2 - a^2} \right)$  respectively.

(6) **Latus-rectum :** The chord of the hyperbola which passes through the focus and is perpendicular to its transverse axis is called latus-rectum.

Length of latus-rectum  $LL' = L_1L'_1 = \frac{2b^2}{a} = 2a(e^2 - 1)$  and end points of latus-rectum  $L \left( ae, \frac{b^2}{a} \right); L' \left( ae, -\frac{b^2}{a} \right); L_1 \left( -ae, \frac{b^2}{a} \right); L'_1 \left( -ae, -\frac{b^2}{a} \right)$  respectively.

(7) **Foci and directrices:** The points  $S(ae, 0)$  and  $S'(-ae, 0)$  are the foci of the hyperbola and  $ZM$  and  $Z'M'$  are two directrices of the hyperbola and their equations are  $x = \frac{a}{e}$  and  $x = -\frac{a}{e}$  respectively.

Distance between foci  $SS' = 2ae$  and distance between directrices  $ZZ' = 2a/e$ .

(8) **Focal chord :** A chord of the hyperbola passing through its focus is called a focal chord.

(9) **Focal distance :** The difference of any point on the hyperbola from the focus is called the focal distance of the point.

From the figure,  $SP = ePM = e\left(x_1 - \frac{a}{e}\right) = ex_1 - a$ ,  $SP = ePM = e\left(x_1 + \frac{a}{e}\right) = ex_1 + a$

The difference of the focal distance of a point on the hyperbola is constant and is equal to the length of transverse axis.

$$|SP - SP'| = 2a = AA' = \text{Transverse axis}$$

**Example: 3** The eccentricity of the hyperbola which passes through (3, 0) and  $(3\sqrt{2}, 2)$  is

- (a)  $\sqrt{13}$  (b)  $\frac{\sqrt{13}}{3}$  (c)  $\sqrt{\frac{13}{4}}$  (d) None of these

**Solution:** (b) Let equation of hyperbola is  $x^2/a^2 - y^2/b^2 = 1$ . Point (3, 0) lies on hyperbola

So,  $\frac{(3)^2}{a^2} - \frac{0}{b^2} = 1$  or  $\frac{9}{a^2} = 1$  or  $a^2 = 9$  and point  $(3\sqrt{2}, 2)$  also lies on hyperbola. So,

$$\frac{3(\sqrt{2})^2}{a^2} - \frac{(2)^2}{b^2} = 1$$

Put  $a^2 = 9$  we get,  $\frac{18}{9} - \frac{4}{b^2} = 1$  or  $2 - \frac{4}{b^2} = 1$  or  $-\frac{4}{b^2} = 1 - 2$  or  $\frac{4}{b^2} = 1$  or  $b^2 = 4$

We know that  $b^2 = a^2(e^2 - 1)$ . Putting values of  $a^2$  and  $b^2$

$$4 = 9(e^2 - 1) \text{ or } e^2 - 1 = \frac{4}{9} \text{ or } e^2 = 1 + \frac{4}{9} \text{ or } e = \sqrt{(1 + 4/9)} \text{ or } e = \sqrt{(13/9)} = \frac{\sqrt{13}}{3}.$$

**Example: 4** The foci of the hyperbola  $9x^2 - 16y^2 = 144$  are

- (a)  $(\pm 4, 0)$  (b)  $(0, \pm 4)$  (c)  $(\pm 5, 0)$  (d)  $(0, \pm 5)$

**Solution:** (c) The equation of hyperbola is  $\frac{x^2}{16} - \frac{y^2}{9} = 1$ .

Now,  $b^2 = a^2(e^2 - 1) \Rightarrow 9 = 16(e^2 - 1) \Rightarrow e = \frac{5}{4}$ . Hence foci are  $(\pm ae, 0) = \left(\pm 4 \cdot \frac{5}{4}, 0\right)$  i.e.,  $(\pm 5, 0)$

**Example: 5** If the foci of the ellipse  $\frac{x^2}{16} + \frac{y^2}{b^2} = 1$  and the hyperbola  $\frac{x^2}{144} - \frac{y^2}{81} = \frac{1}{25}$  coincide, then the value of  $b^2$  is

- (a) 1 (b) 5 (c) 7 (d) 9

**Solution:** (c) For hyperbola,  $\frac{x^2}{144} - \frac{y^2}{81} = \frac{1}{25}$

$$A = \sqrt{\frac{144}{25}}, B = \sqrt{\frac{81}{25}}, e_1 = \sqrt{1 + \frac{B^2}{A^2}} = \sqrt{1 + \frac{81}{144}} = \sqrt{\frac{225}{144}} = \frac{5}{4}$$

Therefore foci =  $(\pm ae_1, 0) = \left(\pm \frac{12}{5} \cdot \frac{5}{4}, 0\right) = (\pm 3, 0)$ . Therefore foci of ellipse i.e.,  $(\pm 4e, 0) = (\pm 3, 0)$

(For ellipse  $a = 4$ )

$$\Rightarrow e = \frac{3}{4}, \text{ Hence } b^2 = 16\left(1 - \frac{9}{16}\right) = 7.$$

**Example: 6** If  $PQ$  is a double ordinate of hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  such that  $CPQ$  is an equilateral triangle,  $C$  being the centre of the hyperbola. Then the eccentricity  $e$  of the hyperbola satisfies

- (a)  $1 < e < 2/\sqrt{3}$  (b)  $e = 2/\sqrt{3}$  (c)  $e = \sqrt{3}/2$  (d)  $e > 2/\sqrt{3}$


**Solution:** (d) Let  $P(a \sec \theta, b \tan \theta)$ ;  $Q(a \sec \theta, -b \tan \theta)$  be end points of double ordinates and  $C(0, 0)$  is the centre of the hyperbola

$$\text{Now } PQ = 2b \tan \theta; \quad CQ = CP = \sqrt{a^2 \sec^2 \theta + b^2 \tan^2 \theta}$$

$$\text{Since } CQ = CP = PQ, \therefore 4b^2 \tan^2 \theta = a^2 \sec^2 \theta + b^2 \tan^2 \theta$$

$$\Rightarrow 3b^2 \tan^2 \theta = a^2 \sec^2 \theta \Rightarrow 3b^2 \sin^2 \theta = a^2$$

$$\Rightarrow 3a^2(e^2 - 1)\sin^2 \theta = a^2 \Rightarrow 3(e^2 - 1)\sin^2 \theta = 1$$




$$\Rightarrow \frac{1}{3(e^2 - 1)} = \sin^2 \theta < 1 \quad (\ominus \sin^2 \theta < 1)$$

$$\Rightarrow \frac{1}{e^2 - 1} < 3 \Rightarrow e^2 - 1 > \frac{1}{3} \Rightarrow e^2 > \frac{4}{3} \Rightarrow e > \frac{2}{\sqrt{3}}$$

### 5.3.3 CONJUGATE HYPERBOLA

The hyperbola whose transverse and conjugate axis are respectively the conjugate and transverse axis of a given hyperbola is called conjugate hyperbola of the given hyperbola.

Hyperbola Fundamentals	$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$	$-\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ or $\frac{x^2}{a^2} - \frac{y^2}{b^2} = -1$
Centre	(0, 0)	(0, 0)
Length of transverse axis	2a	2b
Length of conjugate axis	2b	2a
Foci	(± ae, 0)	(0, ± be)
Equation of directrices	$x = \pm a/e$	$y = \pm b/e$
Eccentricity	$e = \sqrt{\left(\frac{a^2 + b^2}{a^2}\right)}$	$e = \sqrt{\left(\frac{a^2 + b^2}{b^2}\right)}$
Length of latus rectum	$\frac{2b^2}{a}$	$\frac{2a^2}{b}$
Parametric co-ordinates	$(a \sec \phi, b \tan \phi), 0 \leq \phi < 2\pi$	$(b \sec \phi, a \tan \phi), 0 \leq \phi < 2\pi$
Focal radii	$SP = ex_1 - a$ & $SP = ex_1 + a$	$SP = ey_1 - b$ & $SP = ey_1 + b$
Difference of focal radii (SP - SP)	2a	2b
Tangents at the vertices	$x = -a, x = a$	$y = -b, y = b$
Equation of the transverse axis	$y = 0$	$x = 0$
Equation of the conjugate axis	$x = 0$	$y = 0$



Note : ☐ If  $e$  and  $e'$  are the eccentricities of a hyperbola and its conjugate, then  $\frac{1}{e^2} + \frac{1}{e'^2} = 1$ .

☐ The foci of a hyperbola and its conjugate are concyclic.

**Example: 7** The eccentricity of the conjugate hyperbola of the hyperbola  $x^2 - 3y^2 = 1$ , is

(a) 2

(b)  $\frac{2}{\sqrt{3}}$

(c) 4

(d)  $\frac{4}{3}$

**Solution:** (a) The given hyperbola is  $\frac{x^2}{1} - \frac{y^2}{1/3} = 1$ . Here  $a^2 = 1$  and  $b^2 = \frac{1}{3}$

$$\text{Since } b^2 = a^2(e^2 - 1) \Rightarrow \frac{1}{3} = 1(e^2 - 1) \Rightarrow e^2 = \frac{4}{3} \Rightarrow e = \frac{2}{\sqrt{3}}$$

If  $e'$  is the eccentricity of the conjugate hyperbola, then  $\frac{1}{e^2} + \frac{1}{e'^2} = 1$

$$\Rightarrow \frac{1}{e'^2} = 1 - \frac{1}{e^2} = 1 - \frac{3}{4} = \frac{1}{4} \Rightarrow e' = 2.$$

### 5.3.4 SPECIAL FORM OF HYPERBOLA

If the centre of hyperbola is  $(h, k)$  and axes are parallel to the co-ordinate axes, then its equation is  $\frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1$ . By shifting the origin at  $(h, k)$  without rotating the co-ordinate axes, the above equation reduces to  $\frac{X^2}{a^2} - \frac{Y^2}{b^2} = 1$ , where  $x = X + h, y = Y + k$ .

**Example: 8** The equation of the hyperbola whose foci are  $(6, 4)$  and  $(-4, 4)$  and eccentricity 2 is given by

(a)  $12x^2 - 4y^2 - 24x + 32y - 127 = 0$

(b)  $12x^2 + 4y^2 + 24x - 32y - 127 = 0$

(c)  $12x^2 - 4y^2 - 24x - 32y + 127 = 0$

(d)  $12x^2 - 4y^2 + 24x + 32y + 127 = 0$

**Solution:** (a) Foci are  $(6, 4)$  and  $(-4, 4)$  and  $e = 2$ .

$$\therefore \text{Centre is } \left( \frac{6-4}{2}, \frac{4+4}{2} \right) = (1, 4)$$

$$\text{So, } ae + 1 = 6 \Rightarrow ae = 5 \Rightarrow a = \frac{5}{2} \text{ and } b = \frac{5}{2}\sqrt{3}$$

$$\text{Hence, the required equation is } \frac{(x-1)^2}{25/4} - \frac{(y-4)^2}{(75/4)} = 1 \text{ or } 12x^2 - 4y^2 - 24x + 32y - 127 = 0$$

**Example: 9** The equations of the directrices of the conic  $x^2 + 2x - y^2 + 5 = 0$  are

(a)  $x = \pm 1$

(b)  $y = \pm 2$

(c)  $y = \pm \sqrt{2}$

(d)  $x = \pm \sqrt{3}$

$$\text{Solution: (c) } (x+1)^2 - y^2 - 1 + 5 = 0 \Rightarrow -\frac{(x+1)^2}{4} + \frac{y^2}{4} = 1$$

$$\text{Equation of directrices of } \frac{y^2}{b^2} - \frac{x^2}{a^2} = 1 \text{ are } y = \pm \frac{b}{e}$$


$$\text{Here } b = 2, e = \sqrt{1+1} = \sqrt{2}. \text{ Hence, } y = \pm \frac{2}{\sqrt{2}} \Rightarrow y = \pm \sqrt{2}.$$

### 5.3.5 AUXILIARY CIRCLE OF HYPERBOLA

Let  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  be the hyperbola with centre  $C$  and transverse axis  $AA'$ . Therefore circle drawn with centre  $C$  and segment  $AA'$  as a diameter is called auxiliary circle of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$

$$\therefore \text{Equation of the auxiliary circle is } x^2 + y^2 = a^2$$

Let  $\angle QCN = \phi$



Here  $P$  and  $Q$  are the corresponding points on the hyperbola and the auxiliary circle ( $0 \leq \phi < 2\pi$ )

(1) **Parametric equations of hyperbola** : The equations  $x = a \sec \phi$  and  $y = b \tan \phi$  are known as the parametric equations of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ . This  $(a \sec \phi, b \tan \phi)$  lies on the hyperbola for all values of  $\phi$ .

Position of points $Q$ on auxiliary circle and the corresponding point $P$ which describes the hyperbola and $0 \leq \phi < 2\pi$		
$\phi$ varies from	$Q(a \cos \phi, a \sin \phi)$	$P(a \sec \phi, b \tan \phi)$
0 to $\frac{\pi}{2}$	I	I
$\frac{\pi}{2}$ to $\pi$	II	III
$\pi$ to $\frac{3\pi}{2}$	III	II
$\frac{3\pi}{2}$ to $2\pi$	IV	IV

Note :  $\square$  The equations  $x = a \cosh \theta$  and  $y = b \sinh \theta$  are also known as the parametric equations of the hyperbola and the co-ordinates of any point on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  are expressible as  $(a \cosh \theta, b \sinh \theta)$ , where  $\cosh \theta = \frac{e^\theta + e^{-\theta}}{2}$  and  $\sinh \theta = \frac{e^\theta - e^{-\theta}}{2}$ .

**Example: 10** The distance between the directrices of the hyperbola  $x = 8 \sec \theta$ ,  $y = 8 \tan \theta$  is

(a)  $16\sqrt{2}$

(b)  $\sqrt{2}$

(c)  $8\sqrt{2}$

(d)  $4\sqrt{2}$

**Solution:** (c) Equation of hyperbola is  $x = 8 \sec \theta, y = 8 \tan \theta \Rightarrow \frac{x}{8} = \sec \theta, \frac{y}{8} = \tan \theta$

$$\therefore \sec^2 \theta - \tan^2 \theta = 1 \Rightarrow \frac{x^2}{8^2} - \frac{y^2}{8^2} = 1$$


$$\text{Here } a = 8, b = 8. \text{ Now } e = \sqrt{1 + \frac{b^2}{a^2}} = \sqrt{1 + \frac{8^2}{8^2}} = \sqrt{2}$$

$$\therefore \text{Distance between directrices} = \frac{2a}{e} = \frac{2 \times 8}{\sqrt{2}} = 8\sqrt{2}.$$

### 5.3.6 POSITION OF A POINT WITH RESPECT TO A HYPERBOLA

Let the hyperbola be  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ .

Then  $P(x_1, y_1)$  will lie inside, on or outside the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  according as  $\frac{x_1^2}{a^2} - \frac{y_1^2}{b^2} - 1$  is positive, zero or negative.



**Example: 11** The number of tangents to the hyperbola  $\frac{x^2}{4} - \frac{y^2}{3} = 1$

through (4, 1) is

[AMU 1998]

(a) 1

(b) 2

(c) 0

(d) 3

**Solution:** (c) Since the point (4, 1) lies inside the hyperbola  $\left[ \ominus \frac{16}{4} - \frac{1}{3} - 1 > 0 \right]$ ;  $\therefore$  Number of tangents through (4, 1) is 0.

### 5.3.7 INTERSECTION OF A LINE AND A HYPERBOLA

The straight line  $y = mx + c$  will cut the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  in two points may be real, coincident or imaginary according as  $c^2 >, =, < a^2 m^2 - b^2$ .

**Condition of tangency :** If straight line  $y = mx + c$  touches the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , then  $c^2 = a^2 m^2 - b^2$ .

### 5.3.8 EQUATIONS OF TANGENT IN DIFFERENT FORMS

(1) **Point form :** The equation of the tangent to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  at  $(x_1, y_1)$  is  $\frac{xx_1}{a^2} - \frac{yy_1}{b^2} = 1$ .

(2) **Parametric form :** The equation of tangent to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  at  $(a \sec \phi, b \tan \phi)$  is

$$\frac{x}{a} \sec \phi - \frac{y}{b} \tan \phi = 1$$

(3) **Slope form :** The equations of tangents of slope  $m$  to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  are  $y = mx \pm \sqrt{a^2 m^2 - b^2}$  and the co-ordinates of points of contacts are  $\left( \pm \frac{a^2 m}{\sqrt{a^2 m^2 - b^2}}, \pm \frac{b^2}{\sqrt{a^2 m^2 - b^2}} \right)$ .

**Note :**  $\square$  If the straight line  $lx + my + n = 0$  touches the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , then  $a^2 l^2 - b^2 m^2 = n^2$ .

$\square$  If the straight line  $x \cos \alpha + y \sin \alpha = p$  touches the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ ,

then  $a^2 \cos^2 \alpha - b^2 \sin^2 \alpha = p^2$

$\square$  Two tangents can be drawn from an outside point to a hyperbola.

### Important Tips

$\Rightarrow$  For hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  and  $\frac{y^2}{a^2} - \frac{x^2}{b^2} = 1$ , the equation of common tangent is  $y = \pm x \pm \sqrt{a^2 - b^2}$ , points of contacts are  $\left( \pm \frac{a^2}{\sqrt{a^2 - b^2}}, \pm \frac{b^2}{\sqrt{a^2 - b^2}} \right)$  and length of common tangent is  $\sqrt{2} \cdot \frac{(a^2 + b^2)}{\sqrt{a^2 - b^2}}$ .

$\Rightarrow$  If the line  $y = mx \pm \sqrt{a^2 m^2 - b^2}$  touches the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  at the point  $(a \sec \theta, b \tan \theta)$ , then  $\theta = \sin^{-1} \left( \frac{b}{am} \right)$ .

**Example: 12** The value of  $m$  for which  $y = mx + 6$  is a tangent to the hyperbola  $\frac{x^2}{100} - \frac{y^2}{49} = 1$ , is

- (a)  $\sqrt{\frac{17}{20}}$  (b)  $\sqrt{\frac{20}{17}}$  (c)  $\sqrt{\frac{3}{20}}$  (d)  $\sqrt{\frac{20}{3}}$

**Solution:** (a) For condition of tangency,  $c^2 = a^2 m^2 - b^2$ . Here  $c = 6$ ,  $a = 10$ ,  $b = 7$

Then,  $(6)^2 = (10)^2 \cdot m^2 - (7)^2$

$$36 = 100m^2 - 49 \Rightarrow 100m^2 = 85 \Rightarrow m^2 = \frac{17}{20} \Rightarrow m = \sqrt{\frac{17}{20}}$$

**Example: 13** If  $m_1$  and  $m_2$  are the slopes of the tangents to the hyperbola  $\frac{x^2}{25} - \frac{y^2}{16} = 1$  which pass through the point  $(6, 2)$ , then

- (a)  $m_1 + m_2 = \frac{24}{11}$  (b)  $m_1 m_2 = \frac{20}{11}$  (c)  $m_1 + m_2 = \frac{48}{11}$  (d)  $m_1 m_2 = \frac{11}{20}$

**Solution:** (a, b) The line through  $(6, 2)$  is  $y - 2 = m(x - 6) \Rightarrow y = mx + 2 - 6m$

Now, from condition of tangency  $(2 - 6m)^2 = 25m^2 - 16$

$$\Rightarrow 36m^2 + 4 - 24m - 25m^2 + 16 = 0 \Rightarrow 11m^2 - 24m + 20 = 0$$

Obviously, its roots are  $m_1$  and  $m_2$ , therefore  $m_1 + m_2 = \frac{24}{11}$  and  $m_1 m_2 = \frac{20}{11}$

**Example: 14** The points of contact of the line  $y = x - 1$  with  $3x^2 - 4y^2 = 12$  is

- (a)  $(4, 3)$  (b)  $(3, 4)$  (c)  $(4, -3)$  (d) None of these

**Solution:** (a) The equation of line and hyperbola are  $y = x - 1$  .....(i) and  $3x^2 - 4y^2 = 12$  .....(ii)

From (i) and (ii), we get  $3x^2 - 4(x - 1)^2 = 12$

$$\Rightarrow 3x^2 - 4(x^2 - 2x + 1) = 12 \text{ or } x^2 - 8x + 16 = 0 \Rightarrow x = 4$$

From (i),  $y = 3$  so points of contact is  $(4, 3)$

**Trick :** Points of contact are  $\left( \pm \frac{a^2 m}{\sqrt{a^2 m^2 - b^2}}, \pm \frac{b^2}{\sqrt{a^2 m^2 - b^2}} \right)$ .

Here  $a^2 = 4$ ,  $b^2 = 3$  and  $m = 1$ . So the required points of contact is  $(4, 3)$ .

**Example: 15**  $P$  is a point on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ ,  $N$  is the foot of the perpendicular from  $P$  on the transverse axis. The tangent to the hyperbola at  $P$  meets the transverse axis at  $T$ . If  $O$  is the centre of the hyperbola, then  $OT \cdot ON$  is equal to

- (a)  $e^2$  (b)  $a^2$  (c)  $b^2$  (d)  $\frac{b^2}{a^2}$

**Solution:** (b) Let  $P(x_1, y_1)$  be a point on the hyperbola. Then the co-ordinates of  $N$  are  $(x_1, 0)$ .

The equation of the tangent at  $(x_1, y_1)$  is  $\frac{xx_1}{a^2} - \frac{yy_1}{b^2} = 1$

This meets  $x$ -axis at  $T\left(\frac{a^2}{x_1}, 0\right)$ ;  $\therefore OT \cdot ON = \frac{a^2}{x_1} \times x_1 = a^2$


**Example: 16** If the tangent at the point  $(2\sec\phi, 3\tan\phi)$  on the hyperbola  $\frac{x^2}{4} - \frac{y^2}{9} = 1$  is parallel to  $3x - y + 4 = 0$ , then the value of  $\phi$  is

- (a)  $45^\circ$  (b)  $60^\circ$  (c)  $30^\circ$  (d)  $75^\circ$

**Solution:** (c) Here  $x = 2\sec\phi$  and  $y = 3\tan\phi$

Differentiating w.r.t.  $\phi$

$$\frac{dx}{d\phi} = 2\sec\phi \tan\phi \text{ and } \frac{dy}{d\phi} = 3\sec^2\phi$$





$$\therefore \text{Gradient of tangent } \frac{dy}{dx} = \frac{dy/d\phi}{dx/d\phi} = \frac{3\sec^2\phi}{2\sec\phi\tan\phi}; \therefore \frac{dy}{dx} = \frac{3}{2}\operatorname{cosec}\phi \quad \dots(i)$$

But tangent is parallel to  $3x - y + 4 = 0$ ;  $\therefore$  Gradient  $m = 3$

$$\text{From (i) and (ii), } \frac{3}{2}\operatorname{cosec}\phi = 3 \Rightarrow \operatorname{cosec}\phi = 2, \therefore \phi = 30^\circ$$

**Example: 17** The slopes of the common tangents to the hyperbola  $\frac{x^2}{9} - \frac{y^2}{16} = 1$  and  $\frac{y^2}{9} - \frac{x^2}{16} = 1$  are

(a)  $-2, 2$

(b)  $-1, 1$

(c)  $1, 2$

(d)  $2, 1$

**Solution:** (b) Given hyperbola are  $\frac{x^2}{9} - \frac{y^2}{16} = 1$  .....(i) and  $\frac{y^2}{9} - \frac{x^2}{16} = 1$  .....(ii)

Any tangent to (i) having slope  $m$  is  $y = mx \pm \sqrt{9m^2 - 16}$  .....(iii)

Putting in (ii), we get,  $16[mx \pm \sqrt{9m^2 - 16}]^2 - 9x^2 = 144$

$$\Rightarrow (16m^2 - 9)x^2 \pm 32m\sqrt{9m^2 - 16}x + 144m^2 - 256 - 144 = 0$$

$$\Rightarrow (16m^2 - 9)x^2 \pm 32m\sqrt{9m^2 - 16}x + (144m^2 - 400) = 0 \quad \dots(iv)$$

If (iii) is a tangent to (ii), then the roots of (iv) are real and equal.

$$\therefore \text{Discriminant} = 0; 32 \times 32m^2(9m^2 - 16) = 4(16m^2 - 9)(144m^2 - 400) = 64(16m^2 - 9)(9m^2 - 25)$$

$$\Rightarrow 16m^2(9m^2 - 16) = (16m^2 - 9)(9m^2 - 25) \Rightarrow 144m^4 - 256m^2 = 144m^4 - 481m^2 + 225$$


$$\Rightarrow 225m^2 = 225 \Rightarrow m^2 = 1 \Rightarrow m = \pm 1$$

### 5.3.9 EQUATION OF PAIR OF TANGENTS

If  $P(x_1, y_1)$  be any point outside the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  then a pair of tangents  $PQ, PR$  can be drawn to it from  $P$ .


The equation of pair of tangents  $PQ$  and  $PR$  is  $SS_1 = T^2$

$$\text{where, } S = \frac{x^2}{a^2} - \frac{y^2}{b^2} - 1, S_1 = \frac{x_1^2}{a^2} - \frac{y_1^2}{b^2} - 1, T = \frac{xx_1}{a^2} - \frac{yy_1}{b^2} - 1$$



**Director circle :** The director circle is the locus of points from which perpendicular tangents are drawn to the given hyperbola. The equation of the director circle of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  is

$$x^2 + y^2 = a^2 - b^2$$



**Example: 18** The locus of the point of intersection of tangents to the hyperbola  $4x^2 - 9y^2 = 36$  which meet at a constant angle  $\pi/4$ , is

(a)  $(x^2 + y^2 - 5)^2 = 4(9y^2 - 4x^2 + 36)$

(b)  $(x^2 + y^2 - 5) = 4(9y^2 - 4x^2 + 36)$

(c)  $4(x^2 + y^2 - 5)^2 = (9y^2 - 4x^2 + 36)$

(d) None of these

**Solution:** (a) Let the point of intersection of tangents be  $P(x_1, y_1)$ . Then the equation of pair of tangents from  $P(x_1, y_1)$  to the given hyperbola is  $(4x^2 - 9y^2 - 36)(4x_1^2 - 9y_1^2 - 36) = [4x_1x - 9y_1y - 36]^2$  .....(i)

From  $SS_1 = T^2$  or  $x^2(y_1^2 + 4) + 2x_1y_1xy + y^2(x_1^2 - 9) + \dots = 0$  .....(ii)

Since angle between the tangents is  $\pi/4$ .

$\therefore \tan(\pi/4) = \frac{2\sqrt{[x_1^2y_1^2 - (y_1^2 + 4)(x_1^2 - 9)]}}{y_1^2 + 4 + x_1^2 - 9}$ . Hence locus of  $P(x_1, y_1)$  is  $(x^2 + y^2 - 5)^2 = 4(9y^2 - 4x^2 + 36)$ .

### 5.3.10 EQUATIONS OF NORMAL IN DIFFERENT FORMS

(1) **Point form** : The equation of normal to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  at  $(x_1, y_1)$  is


$$\frac{a^2x}{x_1} + \frac{b^2y}{y_1} = a^2 + b^2.$$

(2) **Parametric form**: The equation of normal at  $(a \sec \theta, b \tan \theta)$  to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  is

$$a x \cos \theta + b y \cot \theta = a^2 + b^2$$

(3) **Slope form**: The equation of the normal to the hyperbola

$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  in terms of the slope  $m$  of the normal is  $y = mx \pm \frac{m(a^2 + b^2)}{\sqrt{a^2 - b^2m^2}}$



(4) **Condition for normality** : If  $y = mx + c$  is the normal of  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$

then  $c = \pm \frac{m(a^2 + b^2)}{\sqrt{a^2 - m^2b^2}}$  or  $c^2 = \frac{m^2(a^2 + b^2)^2}{(a^2 - m^2b^2)}$ , which is condition of normality.

(5) **Points of contact** : Co-ordinates of points of contact are  $\left( \pm \frac{a^2}{\sqrt{a^2 - b^2m^2}}, \pm \frac{mb^2}{\sqrt{a^2 - b^2m^2}} \right)$

Note :  $\square$  If the line  $lx + my + n = 0$  will be normal to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , then

$$\frac{a^2}{l^2} - \frac{b^2}{m^2} = \frac{(a^2 + b^2)^2}{n^2}.$$

### Important Tip

In general, four normals can be drawn to a hyperbola from any point and if  $\alpha, \beta, \gamma, \delta$  be the eccentric angles of these four co-normal points, then  $\alpha + \beta + \gamma + \delta$  is an odd multiple of  $\pi$ .

If  $\alpha, \beta, \gamma$  are the eccentric angles of three points on the hyperbola.  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , the normals at which are concurrent, then,  $\sin(\alpha + \beta) + \sin(\beta + \gamma) + \sin(\gamma + \alpha) = 0$

If the normal at P meets the transverse axis in G, then  $SG = e.SP$ . Also the tangent and normal bisect the angle between the focal distances of P.

The feet of the normals to  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  from  $(h, k)$  lie on  $a^2y(x - h) + b^2x(y - k) = 0$ .

**Example: 19** The equation of the normal to the hyperbola  $\frac{x^2}{16} - \frac{y^2}{9} = 1$  at the point  $(8, 3\sqrt{3})$  is

- (a)  $\sqrt{3}x + 2y = 25$  (b)  $x + y = 25$  (c)  $y + 2x = 25$  (d)  $2x + \sqrt{3}y = 25$

**Solution:** (d) From  $\frac{a^2x}{x_1} + \frac{b^2y}{y_1} = a^2 + b^2$

Here  $a^2 = 16$ ,  $b^2 = 9$  and  $(x_1, y_1) = (8, 3\sqrt{3})$

$$\Rightarrow \frac{16x}{8} + \frac{9y}{3\sqrt{3}} = 16 + 9 \text{ i.e., } 2x + \sqrt{3}y = 25.$$

**Example: 20** If the normal at ' $\phi$ ' on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  meets transverse axis at  $G$ , then  $AGAG =$

(Where  $A$  and  $A'$  are the vertices of the hyperbola)

- (a)  $a^2(e^4 \sec^2 \phi - 1)$  (b)  $(a^2 e^4 \sec^2 \phi - 1)$  (c)  $a^2(1 - e^4 \sec^2 \phi)$  (d) None of these

**Solution:** (a) The equation of normal at  $(a \sec \phi, b \tan \phi)$  to the given hyperbola is  $ax \cos \phi + by \cot \phi = (a^2 + b^2)$

This meets the transverse axis i.e.,  $x$ -axis at  $G$ . So the co-ordinates of  $G$  are

$\left( \left( \frac{a^2 + b^2}{a} \right) \sec \phi, 0 \right)$  and the co-ordinates of the vertices  $A$  and  $A'$  are  $A(a, 0)$  and  $A'(-a, 0)$  respectively.

$$\therefore AGAG = \left( -a + \left( \frac{a^2 + b^2}{a} \right) \sec \phi \right) \left( a + \left( \frac{a^2 + b^2}{a} \right) \sec \phi \right) = \left( \frac{a^2 + b^2}{a} \right)^2 \sec^2 \phi - a^2 = (ae^2)^2 \sec^2 \phi - a^2 = a^2(e^4 \sec^2 \phi - 1)$$

**Example: 21** The normal at  $P$  to a hyperbola of eccentricity  $e$ , intersects its transverse and conjugate axis at  $L$  and  $M$  respectively, then the locus of the middle point of  $LM$  is a hyperbola whose eccentricity is

- (a)  $\frac{e}{\sqrt{e^2 - 1}}$  (b)  $\frac{e}{\sqrt{e^4 - 1}}$  (c)  $\frac{e}{\sqrt{a^2 e^2 - 1}}$  (d) None of these

**Solution:** (a) The equation of the normal at  $P(a \sec \phi, b \tan \phi)$  to the hyperbola is  $ax \cos \phi + by \cot \phi = a^2 + b^2 = a^2 e^2$

It meets the transverse and conjugate axes at  $L$  and  $M$ , then  $L(ae^2 \sec \phi, 0)$ ;  $M\left(0, \frac{a^2 e^2 \tan \phi}{b}\right)$

Let the middle point of  $LM$  is  $(\alpha, \beta)$ ; then  $\alpha = \frac{ae^2 \sec \phi}{2} \Rightarrow \sec \phi = \frac{2\alpha}{ae^2}$  .....(i)

and  $\beta = \frac{a^2 e^2 \tan \phi}{2b} \Rightarrow \tan \phi = \frac{2b\beta}{a^2 e^2}$  .....(ii)

$$\ominus \quad 1 = \sec^2 \phi - \tan^2 \phi; \quad 1 = \frac{4\alpha^2}{a^2 e^4} - \frac{4b^2 \beta^2}{a^4 e^4}, \therefore \text{Locus of } (\alpha, \beta) \text{ is } \frac{x^2}{\left(\frac{a^2 e^4}{4}\right)} - \frac{y^2}{\left(\frac{a^4 e^4}{4b^2}\right)} = 1$$

It is a hyperbola, let its eccentricity  $e_1 = \frac{\sqrt{\left(\frac{a^2 e^4}{4} + \frac{a^4 e^4}{4b^2}\right)}}{\left(\frac{a^2 e^4}{4}\right)} = \sqrt{1 + \frac{a^2}{b^2}} = \sqrt{\frac{a^2 + b^2}{b^2}} = \sqrt{\frac{a^2 e^2}{a^2(e^2 - 1)}}; \therefore$

$$e_1 = \frac{e}{\sqrt{e^2 - 1}}.$$


### 5.3.11 EQUATION OF CHORD OF CONTACT OF TANGENTS DRAWN FROM A POINT TO A HYPERBOLA

Let  $PQ$  and  $PR$  be tangents to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  drawn from any external point  $P(x_1, y_1)$ .

Then equation of chord of contact  $QR$  is

or  $\frac{xx_1}{a^2} - \frac{yy_1}{b^2} = 1$

or  $T = 0$  (At  $x_1, y_1$ )




### 5.3.12 EQUATION OF THE CHORD OF THE HYPERBOLA WHOSE MID POINT $(x_1, y_1)$ IS GIVEN

Equation of the chord of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , bisected at the

given point  $(x_1, y_1)$  is  $\frac{xx_1}{a^2} - \frac{yy_1}{b^2} - 1 = \frac{x_1^2}{a^2} - \frac{y_1^2}{b^2} - 1$

i.e.,  $T = S_1$



Note :  $\square$  The length of chord cut off by hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  from the line  $y = mx + c$  is

$$\frac{2ab\sqrt{[c^2 - (a^2m^2 - b^2)](1 + m^2)}}{(b^2 - a^2m^2)}$$

### 5.3.13 EQUATION OF THE CHORD JOINING TWO POINTS ON THE HYPERBOLA

The equation of the chord joining the points  $P(a \sec \phi_1, b \tan \phi_1)$  and  $Q(a \sec \phi_2, b \tan \phi_2)$  is

$$y - b \tan \phi_1 = \frac{b \tan \phi_2 - b \tan \phi_1}{a \sec \phi_2 - a \sec \phi_1} (x - a \sec \phi_1)$$

$$\frac{x}{a} \cos \left( \frac{\phi_1 - \phi_2}{2} \right) - \frac{y}{b} \sin \left( \frac{\phi_1 + \phi_2}{2} \right) = \cos \left( \frac{\phi_1 + \phi_2}{2} \right)$$

Note :  $\square$  If the chord joining two points  $(a \sec \theta_1, b \tan \theta_1)$  and  $(a \sec \theta_2, b \tan \theta_2)$  passes through the

focus of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , then  $\tan \frac{\theta_1}{2} \tan \frac{\theta_2}{2} = \frac{1 - e}{1 + e}$ .

**Example: 22** The equation of the chord of contact of tangents drawn from a point  $(2, -1)$  to the hyperbola  $16x^2 - 9y^2 = 144$  is

(a)  $32x + 9y = 144$

(b)  $32x + 9y = 55$

(c)  $32x + 9y + 144 = 0$

(d)  $32x + 9y + 55 = 0$

**Solution:** (a) From  $T = 0$  i.e.,  $\frac{xx_1}{a^2} - \frac{yy_1}{b^2} = 1$ . Here,  $16x^2 - 9y^2 = 144$  i.e.,  $\frac{x^2}{9} - \frac{y^2}{16} = 1$

So, the equation of chord of contact of tangents drawn from a point  $(2, -1)$  to the hyperbola is  $\frac{2x}{9} - \frac{(-1)y}{16} = 1$

i.e.,  $32x + 9y = 144$

**Example: 23** The point of intersection of tangents drawn to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  at the points where it is intersected by the line  $lx + my + n = 0$  is

- (a)  $\left(-\frac{a^2 l}{n}, \frac{b^2 m}{n}\right)$  (b)  $\left(\frac{a^2 l}{n}, -\frac{b^2 m}{n}\right)$  (c)  $\left(-\frac{a^2 n}{l}, \frac{b^2 n}{m}\right)$  (d)  $\left(\frac{a^2 n}{l}, -\frac{b^2 n}{m}\right)$

**Solution:** (a) Let  $(x_1, y_1)$  be the required point. Then the equation of the chord of contact of tangents drawn from  $(x_1, y_1)$  to the given hyperbola is  $\frac{xx_1}{a^2} - \frac{yy_1}{b^2} = 1$  .....(i)

The given line is  $lx + my + n = 0$  .....(ii)

Equation (i) and (ii) represent the same line

$$\therefore \frac{x_1}{a^2 l} = -\frac{y_1}{b^2 m} = \frac{1}{-n} \Rightarrow x_1 = -\frac{a^2 l}{n}, y_1 = \frac{b^2 m}{n}; \text{ Hence the required point is } \left(-\frac{a^2 l}{n}, \frac{b^2 m}{n}\right).$$

**Example: 24** What will be equation of that chord of hyperbola  $25x^2 - 16y^2 = 400$ , whose mid point is (5, 3) [UPSEAT 1999]

- (a)  $115x - 117y = 17$  (b)  $125x - 48y = 481$  (c)  $127x + 33y = 341$  (d)  $15x + 12y = 105$

**Solution:** (b) According to question,  $S = 25x^2 - 16y^2 - 400 = 0$

Equation of required chord is  $S_1 = T$  .....(i)

$$\text{Here } S_1 = 25(5)^2 - 16(3)^2 - 400 = 625 - 144 - 400 = 81 \text{ and } T = 25xx_1 - 16yy_1 - 400, \text{ where } x_1 = 5, y_1 = 3$$

$$\Rightarrow 25x(5) - 16y(3) - 400 = 125x - 48y - 400$$

So, from (i) required chord is  $125x - 48y - 400 = 81 \Rightarrow 125x - 48y = 481$ .

**Example: 25** The locus of the mid-points of the chords of the circle  $x^2 + y^2 = 16$  which are tangent to the hyperbola  $9x^2 - 16y^2 = 144$  is

- (a)  $(x^2 + y^2)^2 = 16x^2 - 9y^2$  (b)  $(x^2 + y^2)^2 = 9x^2 - 16y^2$   
(c)  $(x^2 - y^2)^2 = 16x^2 - 9y^2$  (d) None of these

**Solution:** (a) The given hyperbola is  $\frac{x^2}{16} - \frac{y^2}{9} = 1$  .....(i)

Any tangent to (i) is  $y = mx + \sqrt{16m^2 - 9}$  .....(ii)

Let  $(x_1, y_1)$  be the mid point of the chord of the circle  $x^2 + y^2 = 16$

Then equation of the chord is  $T = S_1$  i.e.,  $xx_1 + yy_1 - (x_1^2 + y_1^2) = 0$  .....(iii)

Since (ii) and (iii) represent the same line.

$$\therefore \frac{m}{x_1} = \frac{-1}{y_1} = \frac{\sqrt{16m^2 - 9}}{-(x_1^2 + y_1^2)}$$

$$\Rightarrow m = -\frac{x_1}{y_1} \text{ and } (x_1^2 + y_1^2)^2 = y_1^2(16m^2 - 9) \Rightarrow (x_1^2 + y_1^2)^2 = 16 \cdot \frac{x_1^2}{y_1^2} y_1^2 - 9y_1^2 = 16x_1^2 - 9y_1^2$$

$\therefore$  Locus of  $(x_1, y_1)$  is  $(x^2 + y^2)^2 = 16x^2 - 9y^2$ .

### 5.3.14 POLE AND POLAR

Let  $P$  be any point inside or outside the hyperbola. If any straight line drawn through  $P$  intersects the hyperbola at  $A$  and  $B$ . Then the locus of the point of intersection of the tangents to the hyperbola at  $A$  and  $B$  is called the polar of the given point  $P$  with respect to the hyperbola and the point  $P$  is called the pole of the polar.

The equation of the required polar with  $(x_1, y_1)$  as its pole is



$$\frac{xx_1}{a^2} - \frac{yy_1}{b^2} = 1$$

Note :  $\square$  Polar of the focus is the directrix.

$\square$  Any tangent is the polar of its point of contact.

(1) **Pole of a given line :** The pole of a given line  $lx + my + n = 0$  with respect to the hyperbola

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1 \text{ is } (x_1, y_1) = \left( -\frac{a^2 l}{n}, \frac{b^2 m}{n} \right)$$



(2) **Properties of pole and polar**

(i) If the polar of  $P(x_1, y_1)$  passes through  $Q(x_2, y_2)$ , then the polar of  $Q(x_2, y_2)$  goes through  $P(x_1, y_1)$  and such points are said to be conjugate points.

(ii) If the pole of a line  $lx + my + n = 0$  lies on the another line  $l'x + m'y + n' = 0$  then the pole of the second line will lie on the first and such lines are said to be conjugate lines.

(iii) Pole of a given line is same as point of intersection of tangents as its extremities.

### Important Tips

$\Rightarrow$  If the polars of  $(x_1, y_1)$  and  $(x_2, y_2)$  with respect to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  are at right angles, then

$$\frac{x_1 x_2}{y_1 y_2} + \frac{a^4}{b^4} = 0$$

**Example: 26** If the polar of a point w.r.t.  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  touches the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , then the locus of the point is

(a) Given hyperbola

(b) Ellipse

(c) Circle

(d) None of these

**Solution:** (a) Let  $(x_1, y_1)$  be the given point.

$$\text{Its polar w.r.t. } \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \text{ is } \frac{xx_1}{a^2} + \frac{yy_1}{b^2} = 1 \text{ i.e., } y = \frac{b^2}{y_1} \left( 1 - \frac{xx_1}{a^2} \right) = -\frac{b^2 x_1}{a^2 y_1} x + \frac{b^2}{y_1}$$

$$\text{This touches } \frac{x^2}{a^2} - \frac{y^2}{b^2} = 1 \text{ if } \left( \frac{b^2}{y_1} \right)^2 = a^2 \cdot \left( \frac{b^2 x_1}{a^2 y_1} \right) - b^2 \Rightarrow \frac{b^4}{y_1^2} = \frac{a^2 b^4 x_1^2}{a^4 y_1^2} - b^2 \Rightarrow \frac{b^2}{y_1^2} = \frac{b^2 x_1^2}{a^2 y_1^2} - 1$$

$$\Rightarrow \frac{x_1^2}{a^2} - \frac{y_1^2}{b^2} = 1$$

$\therefore$  Locus of  $(x_1, y_1)$  is  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ . Which is the same hyperbola.

**Example: 27** The locus of the poles of the chords of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , which subtend a right angle at the centre is

$$(a) \frac{x^2}{a^4} + \frac{y^2}{b^4} = \frac{1}{a^2} - \frac{1}{b^2} \quad (b) \frac{x^2}{a^2} - \frac{y^2}{b^2} = \frac{1}{a^2} - \frac{1}{b^2} \quad (c) \frac{x^2}{a^4} - \frac{y^2}{b^4} = \frac{1}{a^2} + \frac{1}{b^2} \quad (d) \frac{x^2}{a^4} - \frac{y^2}{b^4} = \frac{1}{a^2} - \frac{1}{b^2}$$

**Solution:** (a) Let  $(x_1, y_1)$  be the pole w.r.t.  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  .....(i)

Then equation of polar is  $\frac{hx}{a^2} - \frac{ky}{b^2} = 1$  .....(ii)

The equation of lines joining the origin to the points of intersection of (i) and (ii) is obtained by making homogeneous (i) with the help of (ii), then  $\left(\frac{x^2}{a^2} - \frac{y^2}{b^2}\right) = \left(\frac{hx}{a^2} - \frac{ky}{b^2}\right)^2 \Rightarrow$

$$x^2\left(\frac{1}{a^2} - \frac{h^2}{a^4}\right) - y^2\left(\frac{1}{b^2} + \frac{k^2}{b^4}\right) + \frac{2hk}{a^2b^2}xy = 0$$


Since the lines are perpendicular, then coefficient of  $x^2$  + coefficient of  $y^2 = 0$

$$\frac{1}{a^2} - \frac{h^2}{a^4} - \frac{1}{b^2} - \frac{k^2}{b^4} = 0 \text{ or } \frac{h^2}{a^4} + \frac{k^2}{b^4} = \frac{1}{a^2} - \frac{1}{b^2}. \text{ Hence required locus is } \frac{x^2}{a^4} + \frac{y^2}{b^4} = \frac{1}{a^2} - \frac{1}{b^2}$$

### 5.3.15 DIAMETER OF THE HYPERBOLA

The locus of the middle points of a system of parallel chords of a hyperbola is called a diameter and the point where the diameter intersects the hyperbola is called the vertex of the diameter.

Let  $y = mx + c$  a system of parallel chords to  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  for different chords then the equation of diameter of the hyperbola is  $y = \frac{b^2x}{a^2m}$ , which is passing through (0, 0)



**Conjugate diameter :** Two diameters are said to be conjugate when each bisects all chords parallel to the others.

If  $y = m_1x$ ,  $y = m_2x$  be conjugate diameters, then  $m_1m_2 = \frac{b^2}{a^2}$ .

Note : ☐ If a pair of diameters be conjugate with respect to a hyperbola, they are conjugate with respect to its conjugate hyperbola also.

☐ In a pair of conjugate diameters of a hyperbola. Only one meets the curve in real points.

☐ The condition for the lines  $AX^2 + 2HXY + BY^2 = 0$  to be conjugate diameters of

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1 \text{ is } a^2A = b^2B.$$

### Important Tips

If  $CD$  is the conjugate diameter of a diameter  $CP$  of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , where  $P$  is  $(asec\phi, btan\phi)$  then coordinates of  $D$  is  $(atan\phi, bsec\phi)$ , where  $C$  is (0, 0).

**Example: 28** If a pair of conjugate diameters meet the hyperbola and its conjugate in  $P$  and  $D$  respectively, then  $CP^2 - CD^2 =$

- (a)  $a^2 + b^2$  (b)  $a^2 - b^2$  (c)  $\frac{a^2}{b^2}$  (d) None of these

**Solution:** (b) Coordinates of  $P$  and  $D$  are  $(asec\phi, btan\phi)$  and  $(atan\phi, bsec\phi)$  respectively.

$$\begin{aligned} \text{Then } (CP)^2 - (CD)^2 &= a^2 \sec^2 \phi + b^2 \tan^2 \phi - a^2 \tan^2 \phi - b^2 \sec^2 \phi \\ &= a^2 (\sec^2 \phi - \tan^2 \phi) - b^2 (\sec^2 \phi - \tan^2 \phi) = a^2 (1) - b^2 (1) = a^2 - b^2. \end{aligned}$$

**Example: 29** If the line  $lx + my + n = 0$  passes through the extremities of a pair of conjugate diameters of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  then

- (a)  $a^2 l^2 - b^2 m^2 = 0$  (b)  $a^2 l^2 + b^2 m^2 = 0$  (c)  $a^2 l^2 + b^2 m^2 = n^2$  (d) None of these

**Solution:** (a) The extremities of a pair of conjugate diameters of  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  are  $(asec\phi, btan\phi)$  and  $(atan\phi, bsec\phi)$  respectively.

According to the question, since extremities of a pair of conjugate diameters lie on  $lx + my + n = 0$

$$\therefore l(asec\phi) + m(btan\phi) + n = 0 \Rightarrow l(atan\phi) + m(bsec\phi) + n = 0 \quad \dots\dots(i)$$

$$\text{Then from (i), } a l sec\phi + b m tan\phi = -n \text{ or } a^2 l^2 sec^2 \phi + b^2 m^2 tan^2 \phi + 2ablm sec\phi tan\phi = n^2 \quad \dots\dots(ii)$$

$$\text{And from (ii), } a l tan\phi + b m sec\phi = -n \text{ or } a^2 l^2 tan^2 \phi + b^2 m^2 sec^2 \phi + 2ablm sec\phi tan\phi = n^2 \quad \dots\dots(iii)$$

Then subtracting (ii) from (iii)


$$\therefore a^2 l^2 (sec^2 \phi - tan^2 \phi) + b^2 m^2 (tan^2 \phi - sec^2 \phi) = 0 \text{ or } a^2 l^2 - b^2 m^2 = 0.$$

### 5.3.16 SUBTANGENT AND SUBNORMAL OF THE HYPERBOLA

Let the tangent and normal at  $P(x_1, y_1)$  meet the  $x$ -axis at  $A$  and  $B$  respectively.

$$\text{Length of subtangent } AN = CN - CA = x_1 - \frac{a^2}{x_1}$$


$$\text{Length of subnormal } BN = CB - CN = \frac{(a^2 + b^2)}{a^2} x_1 - x_1 = \frac{b^2}{a^2} x_1 = (e^2 - 1)x_1$$



### 5.3.17 Reflection property of the Hyperbola

If an incoming light ray passing through one focus ( $S$ ) strike convex side of the hyperbola then it will get reflected towards other focus ( $S'$ )

$$\angle TPS = \angle LPM = \alpha$$





**Example: 30** A ray emanating from the point (5, 0) is incident on the hyperbola  $9x^2 - 16y^2 = 144$  at the point  $P$  with abscissa 8; then the equation of reflected ray after first reflection is (Point  $P$  lies in first quadrant)

- (a)  $3\sqrt{3}x - 13y + 15\sqrt{3} = 0$  (b)  $3x - 13y + 15 = 0$  (c)  $3\sqrt{3}x + 13y - 15\sqrt{3} = 0$  (d) None of these

**Solution:** (a) Given hyperbola is  $9x^2 - 26y^2 = 144$ . This equation can be rewritten as  $\frac{x^2}{16} - \frac{y^2}{9} = 1$

Since  $x$  coordinate of  $P$  is 8. Let  $y$ -coordinate of  $P$  is  $\alpha$


$\therefore (8, \alpha)$  lies on (i)

$$\therefore \frac{64}{16} - \frac{\alpha^2}{9} = 1; \quad \therefore \alpha = 27$$

$$\alpha = 3\sqrt{3}$$

Hence coordinate of point  $P$  is  $(8, 3\sqrt{3})$

$\ominus P$  lies in first



$\ominus$  Equation of reflected ray passing through  $P(8, 3\sqrt{3})$  and  $S(-5, 0)$ ;  $\therefore$  Its equation is

$$y - 3\sqrt{3} = \frac{0 - 3\sqrt{3}}{-5 - 8}(x - 8)$$

$$\text{or } 13y - 39\sqrt{3} = 3\sqrt{3}x - 24\sqrt{3} \text{ or } 3\sqrt{3}x - 13y + 15\sqrt{3} = 0$$

### 5.3.18 ASYMPTOTES OF A HYPERBOLA

An asymptote to a curve is a straight line, at a finite distance from the origin, to which the tangent to a curve tends as the point of contact goes to infinity.

The equations of two asymptotes of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  are  $y = \pm \frac{b}{a}x$  or  $\frac{x}{a} \pm \frac{y}{b} = 0$ .

Note :  $\square$  The combined equation of the asymptotes of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  is  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 0$ .

$\square$  When  $b = a$  i.e. the asymptotes of rectangular hyperbola  $x^2 - y^2 = a^2$  are  $y = \pm x$ , which are at right angles.

$\square$  A hyperbola and its conjugate hyperbola have the same asymptotes.


$\square$  The equation of the pair of asymptotes differ the hyperbola and the conjugate hyperbola by the same constant only i.e.  
Hyperbola - Asymptotes = Asymptotes - Conjugated hyperbola

$$\left(\frac{x^2}{a^2} - \frac{y^2}{b^2} - 1\right) - \left(\frac{x^2}{a^2} - \frac{y^2}{b^2}\right) = \left(\frac{x^2}{a^2} - \frac{y^2}{b^2}\right) - \left(\frac{x^2}{a^2} - \frac{y^2}{b^2} + 1\right).$$

or,

$\square$  The asymptotes pass through the centre of the hyperbola.

$\square$  The bisectors of the angles between the asymptotes are the coordinate axes.




- The angle between the asymptotes of the hyperbola  $S=0$  i.e.,  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  is  $2\tan^{-1} \frac{b}{a}$  or  $2\sec^{-1} e$ .
- Asymptotes are equally inclined to the axes of the hyperbola.

### Important Tips

- ☞ The parallelogram formed by the tangents at the extremities of conjugate diameters of a hyperbola has its vertices lying on the asymptotes and is of constant area.

Area of parallelogram  $QRQR = 4(\text{Area of parallelogram QDCP}) = 4ab = \text{Constant}$



- ☞ The product of length of perpendiculars drawn from any point on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  to the asymptotes is  $\frac{a^2 b^2}{a^2 + b^2}$ .

**Example: 31** From any point on the hyperbola,  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  tangents are drawn to the hyperbola

$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 2$ . The area cut-off by the chord of contact on the asymptotes is equal to

- (a)  $\frac{ab}{2}$  (b)  $ab$  (c)  $2ab$  (d)  $4ab$

**Solution:** (d) Let  $P(x_1, y_1)$  be a point on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , then  $\frac{x_1^2}{a^2} - \frac{y_1^2}{b^2} = 1$

The chord of contact of tangent from  $P$  to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 2$  is  $\frac{xx_1}{a^2} - \frac{yy_1}{b^2} = 2$

The equation of asymptotes are  $\frac{x}{a} - \frac{y}{b} = 0$  .....(ii)

And  $\frac{x}{a} + \frac{y}{b} = 0$  .....(iii)

The point of intersection of the asymptotes and chord are  $\left(\frac{2a}{x_1/a - y_1/b}, \frac{2b}{x_1/a - y_1/b}\right); \left(\frac{2a}{x_1/a + y_1/b}, \frac{-2b}{x_1/a + y_1/b}\right), (0, 0)$

$$\therefore \text{Area of triangle} = \frac{1}{2} |(x_1 y_2 - x_2 y_1)| = \frac{1}{2} \left| \left( \frac{-8ab}{x_1^2/a^2 - y_1^2/b^2} \right) \right| = 4ab.$$

**Example: 32** The combined equation of the asymptotes of the hyperbola  $2x^2 + 5xy + 2y^2 + 4x + 5y = 0$

- (a)  $2x^2 + 5xy + 2y^2 = 0$  (b)  $2x^2 + 5xy + 2y^2 - 4x + 5y + 2 = 0$   
 (c)  $2x^2 + 5xy + 2y^2 + 4x + 5y - 2 = 0$  (d)  $2x^2 + 5xy + 2y^2 + 4x + 5y + 2 = 0$

**Solution:** (d) Given, equation of hyperbola  $2x^2 + 5xy + 2y^2 + 4x + 5y = 0$  and equation of asymptotes

$2x^2 + 5xy + 2y^2 + 4x + 5y + \lambda = 0$  .....(i) which is the equation of a pair of straight lines. We know that the standard equation of a pair of straight lines is  $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$

Comparing equation (i) with standard equation, we get  $a=2, b=2, h=\frac{5}{2}, g=2, f=\frac{5}{2}$  and  $c=\lambda$ .

We also know that the condition for a pair of straight lines is  $abc + 2fgh - af^2 - bg^2 - ch^2 = 0$ .

Therefore,  $4\lambda + 25 - \frac{25}{2} - 8 - \frac{25}{4}\lambda = 0$  or  $\frac{-9\lambda}{4} + \frac{9}{2} = 0$  or  $\lambda = 2$

Substituting value of  $\lambda$  in equation (i), we get  $2x^2 + 5xy + 2y^2 + 4x + 5y + 2 = 0$ .

### 5.3.19 RECTANGULAR OR EQUILATERAL HYPERBOLA

(1) **Definition** : A hyperbola whose asymptotes are at right angles to each other is called a rectangular hyperbola. The eccentricity of rectangular hyperbola is always  $\sqrt{2}$ .

The general equation of second degree represents a rectangular hyperbola if  $\Delta \neq 0$ ,  $h^2 > ab$  and coefficient of  $x^2$  + coefficient of  $y^2 = 0$

The equation of the asymptotes of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  are given by  $y = \pm \frac{b}{a}x$ .

The angle between these two asymptotes is given by  $\tan\theta = \frac{\frac{b}{a} - \left(-\frac{b}{a}\right)}{1 + \frac{b}{a}\left(-\frac{b}{a}\right)} = \frac{2b/a}{1 - b^2/a^2} = \frac{2ab}{a^2 - b^2}$ .

If the asymptotes are at right angles, then  $\theta = \pi/2 \Rightarrow \tan\theta = \tan\frac{\pi}{2} \Rightarrow \frac{2ab}{a^2 - b^2} = \tan\frac{\pi}{2} \Rightarrow a^2 - b^2 = 0 \Rightarrow a = b \Rightarrow 2a = 2b$ . Thus the transverse and conjugate axis of a rectangular hyperbola are equal and the equation is  $x^2 - y^2 = a^2$ . The equations of the asymptotes of the rectangular hyperbola are  $y = \pm x$  i.e.,  $y = x$  and  $y = -x$ . Clearly, each of these two asymptotes is inclined at  $45^\circ$  to the transverse axis.

(2) **Equation of the rectangular hyperbola referred to its asymptotes as the axes of coordinates** : Referred to the transverse and conjugate axis as the axes of coordinates, the equation of the rectangular hyperbola is


$$x^2 - y^2 = a^2 \quad \dots(i)$$

The asymptotes of (i) are  $y = x$  and  $y = -x$ . Each of these two asymptotes is inclined at an angle of  $45^\circ$  with the transverse axis, So, if we rotate the coordinate axes through an angle of  $-\pi/4$  keeping the origin fixed, then the axes coincide with the asymptotes of the hyperbola and  $x = X\cos(-\pi/4) - Y\sin(-\pi/4) = \frac{X+Y}{\sqrt{2}}$  and  $y = X\sin(-\pi/4) + Y\cos(-\pi/4) = \frac{Y-X}{\sqrt{2}}$ .

Substituting the values of  $x$  and  $y$  in (i),

$$\text{We obtain the } \left(\frac{X+Y}{\sqrt{2}}\right)^2 - \left(\frac{Y-X}{\sqrt{2}}\right)^2 = a^2 \Rightarrow XY = \frac{a^2}{2} \Rightarrow XY = c^2$$

$$\text{where } c^2 = \frac{a^2}{2}.$$



This is transformed equation of the rectangular hyperbola (i).

(3) **Parametric co-ordinates of a point on the hyperbola  $XY = c^2$**  : If  $t$  is non-zero variable, the coordinates of any point on the rectangular hyperbola  $xy = c^2$  can be written as  $(ct^2/c/t)$ . The point  $(ct^2/c/t)$  on the hyperbola  $xy = c^2$  is generally referred as the point 't'.

For rectangular hyperbola the coordinates of foci are  $(\pm a\sqrt{2}, 0)$  and directrices are  $x = \pm a\sqrt{2}$ .

For rectangular hyperbola  $xy = c^2$ , the coordinates of foci are  $(\pm c\sqrt{2}, \pm c\sqrt{2})$  and directrices are  $x + y = \pm c\sqrt{2}$ .

(4) **Equation of the chord joining points  $t_1$  and  $t_2$**  : The equation of the chord joining two points

$$\left(ct_1, \frac{c}{t_1}\right) \text{ and } \left(ct_2, \frac{c}{t_2}\right) \text{ on the hyperbola } xy = c^2 \text{ is } y - \frac{c}{t_1} = \frac{\frac{c}{t_2} - \frac{c}{t_1}}{ct_2 - ct_1}(x - ct_1) \Rightarrow x + yt_1t_2 = c(t_1 + t_2).$$

(5) **Equation of tangent in different forms**

(i) **Point form** : The equation of tangent at  $(x_1, y_1)$  to the hyperbola  $xy = c^2$  is  $xy_1 + yx_1 = 2c^2$  or  $\frac{x}{x_1} + \frac{y}{y_1} = 2$

(ii) **Parametric form** : The equation of the tangent at  $\left(ct, \frac{c}{t}\right)$  to the hyperbola  $xy = c^2$  is  $\frac{x}{t} + yt = 2c$ . On replacing  $x_1$  by  $ct$  and  $y_1$  by  $\frac{c}{t}$  on the equation of the tangent at  $(x_1, y_1)$  i.e.  $xy_1 + yx_1 = 2c^2$  we get  $\frac{x}{t} + yt = 2c$ .

Note :  $\square$  Point of intersection of tangents at ' $t_1$ ' and ' $t_2$ ' is  $\left(\frac{2ct_1t_2}{t_1 + t_2}, \frac{2c}{t_1 + t_2}\right)$

(6) **Equation of the normal in different forms** : (i) **Point form** : The equation of the normal at  $(x_1, y_1)$  to the hyperbola  $xy = c^2$  is  $xx_1 - yy_1 = x_1^2 - y_1^2$ . As discussed in the equation of the tangent, we have  $\left(\frac{dy}{dx}\right)_{(x_1, y_1)} = -\frac{y_1}{x_1}$

So, the equation of the normal at  $(x_1, y_1)$  is  $y - y_1 = \frac{-1}{\left(\frac{dy}{dx}\right)_{(x_1, y_1)}}(x - x_1) \Rightarrow y - y_1 = \frac{x_1}{y_1}(x - x_1)$

$$\Rightarrow yy_1 - y_1^2 = xx_1 - x_1^2 \Rightarrow xx_1 - yy_1 = x_1^2 - y_1^2$$

This is the required equation of the normal at  $(x_1, y_1)$ .

(ii) **Parametric form**: The equation of the normal at  $\left(ct, \frac{c}{t}\right)$  to the hyperbola  $xy = c^2$  is  $xt^3 - yt - ct^2 + c = 0$ . On replacing  $x_1$  by  $ct$  and  $y_1$  by  $c/t$  in the equation.

$$\text{We obtain } xx_1 - yy_1 = x_1^2 - y_1^2, \quad xct - \frac{yc}{t} = c^2t^2 - \frac{c^2}{t^2} \Rightarrow xt^3 - yt - ct^2 + c = 0$$

Note :  $\square$  The equation of the normal at  $\left(ct, \frac{c}{t}\right)$  is a fourth degree in  $t$ . So, in general, four normals can be drawn from a point to the hyperbola  $xy = c^2$

$\square$  If the normal at  $\left(ct, \frac{c}{t}\right)$  on the curve  $xy = c^2$  meets the curve again in ' $t'$ ' then;  $t = \frac{-1}{t'}$ .

$\square$  Point of intersection of normals at ' $t_1$ ' and ' $t_2$ ' is  $\left(\frac{c\{t_1t_2(t_1^2 + t_1t_2 + t_2^2) - 1\}}{t_1t_2(t_1 + t_2)}, \frac{c\{t_1^3t_2^3 + (t_1^2 + t_1t_2 + t_2^2)\}}{t_1t_2(t_1 + t_2)}\right)$

### Important Tips

- ☞ A triangle has its vertices on a rectangular hyperbola; then the orthocentre of the triangle also lies on the same hyperbola.
- ☞ All conics passing through the intersection of two rectangular hyperbolas are themselves rectangular hyperbolas.
- ☞ An infinite number of triangles can be inscribed in the rectangular hyperbola  $xy = c^2$  whose all sides touch the parabola  $y^2 = 4ax$ .

**Example: 33** If  $5x^2 + \lambda y^2 = 20$  represents a rectangular hyperbola, then  $\lambda$  equals

- (a) 5                      (b) 4                      (c) -5                      (d) None of these

**Solution:** (c) Since the general equation of second degree represents a rectangular hyperbola if  $\Delta \neq 0, h^2 > ab$  and coefficient of  $x^2 + \text{coefficient of } y^2 = 0$ . Therefore the given equation represents a rectangular hyperbola if  $\lambda + 5 = 0$  i.e.,  $\lambda = -5$

**Example: 34** If  $PN$  is the perpendicular from a point on a rectangular hyperbola to its asymptotes, the locus, the mid-point of  $PN$  is

- (a) Circle                      (b) Parabola                      (c) Ellipse                      (d) Hyperbola


**Solution:** (d) Let  $xy = c^2$  be the rectangular hyperbola, and let  $P(x_1, y_1)$  be a point on it. Let  $Q(h, k)$  be the mid-point of  $PN$ . Then the coordinates of  $Q$  are  $(x_1, \frac{y_1}{2})$ .

$$\therefore x_1 = h \text{ and } \frac{y_1}{2} = k \Rightarrow x_1 = h \text{ and } y_1 = 2k$$

But  $(x_1, y_1)$  lies on  $xy = c^2$ .

$$\therefore h \cdot (2k) = c^2 \Rightarrow hk = c^2/2$$

Hence, the locus of  $(h, k)$  is  $xy = c^2/2$ , which is a hyperbola.



**Example: 35** If the normal at  $(ct, \frac{c}{t})$  on the curve  $xy = c^2$  meets the curve again in  $t'$ , then

- (a)  $t = -\frac{1}{t^3}$                       (b)  $t = -\frac{1}{t}$                       (c)  $t = \frac{1}{t^2}$                       (d)  $t^2 = -\frac{1}{t^2}$

**Solution:** (a) The equation of the tangent at  $(ct, \frac{c}{t})$  is  $ty = t^3x - ct^4 + c$

If it passes through  $(ct, \frac{c}{t})$  then

$$\Rightarrow \frac{tc}{t} = t^3 ct - ct^4 + c \Rightarrow t = t^3 t^2 - t^4 t + t \Rightarrow t - t = t^3 t(t - t) \Rightarrow t = -\frac{1}{t^3}$$

**Example: 36** If the tangent and normal to a rectangular hyperbola cut off intercepts  $a_1$  and  $a_2$  on one axis and  $b_1$  and  $b_2$  on the other axis, then

- (a)  $a_1 b_1 + a_2 b_2 = 0$                       (b)  $a_1 b_2 + b_2 a_1 = 0$                       (c)  $a_1 a_2 + b_1 b_2 = 0$                       (d) None of these

**Solution:** (c) Let the hyperbola be  $xy = c^2$ . Tangent at any point  $t$  is  $x + yt^2 - 2ct = 0$

Putting  $y = 0$  and then  $x = 0$  intercepts on the axes are  $a_1 = 2ct$  and  $b_1 = \frac{2c}{t}$

Normal is  $xt^3 - yt - ct^4 + c = 0$ .

Intercepts as above are  $a_2 = \frac{c(t^4 - 1)}{t^3}$ ,  $b^2 = \frac{-c(t^4 - 1)}{t}$

$$\therefore a_1 a_2 + b_1 b_2 = 2ct \times \frac{c(t^4 - 1)}{t^3} + \frac{2c}{t} \times \frac{-c(t^4 - 1)}{t} = \frac{2c^2}{t^2} (t^4 - 1) - \frac{2c^2}{t^2} (t^4 - 1) = 0; \therefore a_1 a_2 + b_1 b_2 = 0.$$

**Example: 37** A variable straight line of slope 4 intersects the hyperbola  $xy=1$  at two points. The locus of the point which divides the line segment between these two points in the ratio 1 : 2 is

- (a)  $16x^2 + 10xy + y^2 = 2$  (b)  $16x^2 - 10xy + y^2 = 2$  (c)  $16x^2 + 10xy + y^2 = 4$  (d) None of these

**Solution:** (a) Let  $P(h, k)$  be any point on the locus. Equation of the line through  $P$  and having slope 4 is

$$y - k = 4(x - h) \quad \dots(i)$$

Suppose this meets  $xy=1$  .....(ii) in  $A(x_1, y_1)$  and  $B(x_2, y_2)$

Eliminating  $y$  between (i) and (ii), we get  $\frac{1}{x} - k = 4(x - h)$

$$\Rightarrow 1 - xk = 4x^2 - 4hx \Rightarrow 4x^2 - (4h - k)x - 1 = 0 \quad \dots(iii)$$

This has two roots say  $x_1, x_2$ ;  $x_1 + x_2 = \frac{4h - k}{4}$  .....(iv) and  $x_1 x_2 = -\frac{1}{4}$  .....(v)

$$\text{Also, } \frac{2x_1 + x_2}{3} = h \quad [\because P \text{ divides } AB \text{ in the ratio } 1 : 2]$$

$$\text{i.e., } 2x_1 + x_2 = 3h \quad \dots(vi)$$

$$(vi) - (iv) \text{ gives, } x_1 = 3h - \frac{4h - k}{4} = \frac{8h + k}{4} \text{ and } x_2 = 3h - 2 \cdot \frac{8h + k}{4} = -\frac{2h + k}{2}$$

$$\text{Putting in (v), we get } \frac{8h + k}{4} \left( -\frac{2h + k}{2} \right) = -\frac{1}{4}$$

$$\Rightarrow (8h + k)(2h + k) = 2 \Rightarrow 16h^2 + 10hk + k^2 = 2$$

$\therefore$  Required locus of  $P(h, k)$  is  $16x^2 + 10xy + y^2 = 2$ .

**Example: 38**  $PQ$  and  $RS$  are two perpendicular chords of the rectangular hyperbola  $xy = c^2$ . If  $C$  is the centre of the rectangular hyperbola, then the product of the slopes of  $CP$ ,  $CQ$ ,  $CR$  and  $CS$  is equal to

- (a)  $-1$  (b)  $1$  (c)  $0$  (d) None of these

**Solution:** (b) Let  $t_1, t_2, t_3, t_4$  be the parameters of the points  $P, Q, R$  and  $S$  respectively. Then, the

coordinates of  $P, Q, R$  and  $S$  are  $\left( ct_1, \frac{c}{t_1} \right)$ ,  $\left( ct_2, \frac{c}{t_2} \right)$ ,  $\left( ct_3, \frac{c}{t_3} \right)$  and  $\left( ct_4, \frac{c}{t_4} \right)$  respectively.

$$\text{Now, } PQ \perp RS \Rightarrow \frac{\frac{c}{t_2} - \frac{c}{t_1}}{ct_2 - ct_1} \times \frac{\frac{c}{t_4} - \frac{c}{t_3}}{ct_4 - ct_3} = -1 \Rightarrow -\frac{1}{t_1 t_2} \times -\frac{1}{t_3 t_4} = -1 \Rightarrow t_1 t_2 t_3 t_4 = -1 \dots(i)$$

$\therefore$  Product of the slopes of  $CP, CQ, CR$  and  $CS$

$$\frac{1}{t_1^2} \times \frac{1}{t_2^2} \times \frac{1}{t_3^2} \times \frac{1}{t_4^2} = \frac{1}{t_1^2 t_2^2 t_3^2 t_4^2} = 1 \quad [\text{Using (i)}]$$

### 5.3.20 INTERSECTION OF A CIRCLE AND A RECTANGULAR HYPERBOLA

If a circle  $x^2 + y^2 + 2gx + 2fy + k = 0$  cuts a rectangular hyperbola  $xy = c^2$  in A, B, C and D and the parameters of these four points be  $t_1, t_2, t_3$  and  $t_4$  respectively; then

$$(1) (i) \sum t_1 = -\frac{2g}{c}$$

$$(ii) \sum t_1 t_2 = \frac{k}{c^2}$$

$$(iii) \sum t_1 t_2 t_3 = -\frac{2f}{c}$$

$$(iv) t_1 t_2 t_3 t_4 = 1 \quad (v) \sum \frac{1}{t_1} = -\frac{2f}{c}$$

$$(2) \text{Orthocentre of } \triangle ABC \text{ is } H\left(-ct_4, -\frac{c}{t_4}\right) \text{ but D is } \left(ct_4, \frac{c}{t_4}\right)$$

Hence H and D are the extremities of a diagonal of rectangular hyperbola.

$$(3) \text{Centre of mean position of four points is } \left\{ \frac{c}{4} \sum t_1, \frac{c}{4} \sum \left( \frac{1}{t_1} \right) \right\} \text{ i.e., } \left( -\frac{g}{2}, -\frac{f}{2} \right)$$

∴ Centres of the circles and rectangular hyperbola are  $(-g, -f)$  and  $(0, 0)$ ; mid point of centres of circle and hyperbola is  $\left( -\frac{g}{2}, -\frac{f}{2} \right)$ . Hence the centre of the mean position of the four points bisects the distance between the centres of the two curves (circle and rectangular hyperbola)

(4) If the circle passing through ABC meet the hyperbola in fourth points D; then centre of circle is  $(-g, -f)$

$$\text{i.e., } \left\{ \frac{c}{2} \left( t_1 + t_2 + t_3 + \frac{1}{t_1 t_2 t_3} \right); \frac{c}{2} \left( \frac{1}{t_1} + \frac{1}{t_2} + \frac{1}{t_3} + t_1 t_2 t_3 \right) \right\}$$

**Example: 39** If a circle cuts a rectangular hyperbola  $xy = c^2$  in A, B, C, D and the parameters of these four points be  $t_1, t_2, t_3$  and  $t_4$  respectively. Then

$$(a) t_1 t_2 = t_3 t_4$$

$$(b) t_1 t_2 t_3 t_4 = 1$$

$$(c) t_1 = t_2$$

$$(d) t_3 = t_4$$

**Solution:** (b) Let the equation of circle be  $x^2 + y^2 = a^2$  .....(i)

Parametric equation of rectangular hyperbola is  $x = ct, y = \frac{c}{t}$

Put the values of x and y in equation (i) we get  $c^2 t^2 + \frac{c^2}{t^2} = a^2 \Rightarrow c^2 t^4 - a^2 t^2 + c^2 = 0$

Hence product of roots  $t_1 t_2 t_3 t_4 = \frac{c^2}{c^2} = 1$

**Example: 40** If the circle  $x^2 + y^2 = a^2$  intersects the hyperbola  $xy = c^2$  in four points  $P(x_1, y_1), Q(x_2, y_2), R(x_3, y_3), S(x_4, y_4)$  then

$$(a) x_1 + x_2 + x_3 + x_4 = 0$$

$$(b) y_1 + y_2 + y_3 + y_4 = 0$$

$$(c) x_1 x_2 x_3 x_4 = c^4$$

$$(d) y_1 y_2 y_3 y_4 = c^4$$

**Solution:** (a,b,c,d) Given, circle is  $x^2 + y^2 = a^2$  .....(i) and hyperbola be  $xy = c^2$  .....(ii)

from (ii)  $y = \frac{c^2}{x}$ . Putting in (i), we get  $x^2 + \frac{c^4}{x^2} = a^2 \Rightarrow x^4 - a^2 x^2 + c^4 = 0$

$$\therefore x_1 + x_2 + x_3 + x_4 = 0, x_1 x_2 x_3 x_4 = c^4$$

Since both the curves are symmetric in x and y,  $\therefore y_1 + y_2 + y_3 + y_4 = 0; y_1 y_2 y_3 y_4 = c^4$ .

## **ASSIGNMENT**

### **DEFINITION STANDARD FORM OF HYPERBOLA , CONJUGATE OF HYPERBOLA**

#### *Basic Level*

1. The locus of the centre of a circle, which touches externally the given two circle, is  
(a) Circle (b) Parabola (c) Hyperbola (d) Ellipse
2. The locus of a point which moves such that the difference of its distances from two fixed points is always a constant is  
(a) A straight line (b) A circle (c) An ellipse (d) A hyperbola
3. The one which does not represent a hyperbola is  
(a)  $xy=1$  (b)  $x^2 - y^2 = 5$  (c)  $(x-1)(y-3)=3$  (d)  $x^2 - y^2 = 0$
4. The equation of the hyperbola whose directrix is  $x+2y=1$ , focus  $(2, 1)$  and eccentricity 2 will be  
(a)  $x^2 - 16xy - 11y^2 - 12x + 6y + 21 = 0$  (b)  $3x^2 + 16xy + 15y^2 - 4x - 14y - 1 = 0$   
(c)  $x^2 + 16xy + 11y^2 - 12x - 6y + 21 = 0$  (d) None of these
5. The locus of the point of intersection of the lines  $\sqrt{3}x - y - 4\sqrt{3}k = 0$  and  $\sqrt{3}kx + ky - 4\sqrt{3} = 0$  for different value of  $k$  is  
(a) Circle (b) Parabola (c) Hyperbola (d) Ellipse
6. Locus of the point of intersection of straight line  $\frac{x}{a} - \frac{y}{b} = m$  and  $\frac{x}{a} + \frac{y}{b} = \frac{1}{m}$  is  
(a) An ellipse (b) A circle (c) A hyperbola (d) A parabola
7. The eccentricity of the hyperbola  $2x^2 - y^2 = 6$  is  
(a)  $\sqrt{2}$  (b) 2 (c) 3 (d)  $\sqrt{3}$
8. Centre of hyperbola  $9x^2 - 16y^2 + 18x + 32y - 151 = 0$  is  
(a)  $(1, -1)$  (b)  $(-1, 1)$  (c)  $(-1, -1)$  (d)  $(1, 1)$
9. The eccentricity of the conic  $x^2 - 4y^2 = 1$ , is  
(a)  $\frac{2}{\sqrt{3}}$  (b)  $\frac{\sqrt{3}}{2}$  (c)  $\frac{2}{\sqrt{5}}$  (d)  $\frac{\sqrt{5}}{2}$
10. The eccentricity of a hyperbola passing through the point  $(3, 0)$ ,  $(3\sqrt{2}, 2)$  will be  
(a)  $\sqrt{13}$  (b)  $\frac{\sqrt{13}}{3}$  (c)  $\frac{\sqrt{13}}{4}$  (d)  $\frac{\sqrt{13}}{2}$
11. If  $(4, 0)$  and  $(-4, 0)$  be the vertices and  $(6, 0)$  and  $(-6, 0)$  be the foci of a hyperbola, then its eccentricity is  
(a)  $5/2$  (b) 2 (c)  $3/2$  (d)  $\sqrt{2}$
12. If  $e$  and  $e'$  are eccentricities of hyperbola and its conjugate respectively, then  
(a)  $\left(\frac{1}{e}\right)^2 + \left(\frac{1}{e'}\right)^2 = 1$  (b)  $\frac{1}{e} + \frac{1}{e'} = 1$  (c)  $\left(\frac{1}{e}\right)^2 + \left(\frac{1}{e'}\right)^2 = 0$  (d)  $\frac{1}{e} + \frac{1}{e'} = 2$



13. If  $e$  and  $e'$  are the eccentricities of the ellipse  $5x^2 + 9y^2 = 45$  and the hyperbola  $5x^2 - 4y^2 = 45$  respectively, then  $ee' =$
- (a) 9 (b) 4 (c) 5 (d) 1
14. The directrix of the hyperbola is  $\frac{x^2}{9} - \frac{y^2}{4} = 1$
- (a)  $x = 9/\sqrt{13}$  (b)  $y = 9/\sqrt{13}$  (c)  $x = 6/\sqrt{13}$  (d)  $y = 6/\sqrt{13}$
15. The latus rectum of the hyperbola  $16x^2 - 9y^2 = 144$ , is
- (a)  $\frac{16}{3}$  (b)  $\frac{32}{3}$  (c)  $\frac{8}{3}$  (d)  $\frac{4}{3}$
16. The foci of the hyperbola  $2x^2 - 3y^2 = 5$ , is
- (a)  $\left(\pm \frac{5}{\sqrt{6}}, 0\right)$  (b)  $\left(\pm \frac{5}{6}, 0\right)$  (c)  $\left(\pm \frac{\sqrt{5}}{6}, 0\right)$  (d) None of these
17. The distance between the directrices of a rectangular hyperbola is 10 units, then distance between its focus
- (a)  $10\sqrt{2}$  (b) 5 (c)  $5\sqrt{2}$  (d) 20
18. The difference of the focal distances of any point on the hyperbola  $9x^2 - 16y^2 = 144$ , is
- (a) 8 (b) 7 (c) 6 (d) 4
19. If the length of the transverse and conjugate axes of a hyperbola be 8 and 6 respectively, then the difference of focal distances of any point of the hyperbola will be
- (a) 8 (b) 6 (c) 14 (d) 2
20. The length of transverse axis of the hyperbola  $3x^2 - 4y^2 = 32$  is
- (a)  $\frac{8\sqrt{2}}{\sqrt{3}}$  (b)  $\frac{16\sqrt{2}}{\sqrt{3}}$  (c)  $\frac{3}{32}$  (d)  $\frac{64}{3}$
21. A hyperbola passes through the points (3, 2) and (-17, 12) and has its centre at origin and transverse axis is along  $x$ -axis. The length of its transverse axis is
- (a) 2 (b) 4 (c) 6 (d) None of these
22. The equation of the hyperbola whose foci are the foci of the ellipse  $\frac{x^2}{25} + \frac{y^2}{9} = 1$  and the eccentricity is 2, is
- (a)  $\frac{x^2}{4} + \frac{y^2}{12} = 1$  (b)  $\frac{x^2}{4} - \frac{y^2}{12} = 1$  (c)  $\frac{x^2}{12} + \frac{y^2}{4} = 1$  (d)  $\frac{x^2}{12} - \frac{y^2}{4} = 1$
23. The distance between the foci of a hyperbola is double the distance between its vertices and the length of its conjugate axis is 6. The equation of the hyperbola referred to its axes as axes of coordinates is
- (a)  $3x^2 - y^2 = 3$  (b)  $x^2 - 3y^2 = 3$  (c)  $3x^2 - y^2 = 9$  (d)  $x^2 - 3y^2 = 9$
24. If  $(0, \pm 4)$  and  $(0, \pm 2)$  be the foci and vertices of a hyperbola then its equation is
- (a)  $\frac{x^2}{4} - \frac{y^2}{12} = 1$  (b)  $\frac{x^2}{12} - \frac{y^2}{4} = 1$  (c)  $\frac{y^2}{4} - \frac{x^2}{12} = 1$  (d)  $\frac{y^2}{12} - \frac{x^2}{4} = 1$

25. The length of the transverse axis of a hyperbola is 7 and it passes through the point  $(5, -2)$ , the equation of the hyperbola is  
 (a)  $\frac{4}{49}x^2 - \frac{196}{51}y^2 = 1$  (b)  $\frac{49}{4}x^2 - \frac{51}{196}y^2 = 1$  (c)  $\frac{4}{49}x^2 - \frac{51}{196}y^2 = 1$  (d) None of these
26. If the centre, vertex and focus of a hyperbola be  $(0, 0)$ ,  $(4, 0)$  and  $(6, 0)$  respectively, then the equation of the hyperbola is  
 (a)  $4x^2 - 5y^2 = 8$  (b)  $4x^2 - 5y^2 = 80$  (c)  $5x^2 - 4y^2 = 80$  (d)  $5x^2 - 4y^2 = 8$
27. The equation of a hyperbola, whose foci are  $(5, 0)$  and  $(-5, 0)$  and the length of whose conjugate axis is 8, is  
 (a)  $9x^2 - 16y^2 = 144$  (b)  $16x^2 - 9y^2 = 144$  (c)  $9x^2 - 16y^2 = 12$  (d)  $16x^2 - 9y^2 = 12$
28. If the latus rectum of an hyperbola be 8 and eccentricity be  $3/\sqrt{5}$ , then the equation of the hyperbola is  
 (a)  $4x^2 - 5y^2 = 100$  (b)  $5x^2 - 4y^2 = 100$  (c)  $4x^2 + 5y^2 = 100$  (d)  $5x^2 + 4y^2 = 100$
29. The equation of the hyperbola whose conjugate axis is 5 and the distance between the foci is 13, is  
 (a)  $25x^2 - 144y^2 = 900$  (b)  $144x^2 - 25y^2 = 900$  (c)  $144x^2 + 25y^2 = 900$  (d)  $25x^2 + 144y^2 = 900$
30. For hyperbola  $\frac{x^2}{\cos^2 \alpha} - \frac{y^2}{\sin^2 \alpha} = 1$  which of the following remains constant with change in ' $\alpha$ '  
 (a) Abscissae of vertices (b) Abscissae of foci (c) Eccentricity (d)
31. The hyperbola is the conic with eccentricity  
 (a)  $e > 1$  (b)  $e < 1$  (c)  $e = 1$  (d)  $e = 0$
32. The eccentricity of the conic  $9x^2 - 16y^2 = 144$  is  
 (a)  $\frac{4}{5}$  (b)  $\frac{5}{4}$  (c)  $\frac{4}{3}$  (d)  $\sqrt{7}$
33. If  $e, e'$  be the eccentricities of two conics  $S$  and  $S'$  and if  $e^2 + e'^2 = 3$ , then both  $S$  and  $S'$  can be  
 (a) Ellipses (b) Parabolas (c) Hyperbolas (d) None of these
34. If  $e_1, e_2$  be respectively the eccentricities of ellipse  $9x^2 + 4y^2 = 36$  and hyperbola  $9x^2 - 4y^2 = 36$ , then  
 (a)  $e_1^2 + e_2^2 > 3$  (b)  $e_1^2 + e_2^2 = 2$  (c)  $e_1^2 + e_2^2 > 4$  (d)  $e_1^2 + e_2^2 < 4$
35. The length of the latus rectum of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = -1$  is  
 (a)  $\frac{2a^2}{b}$  (b)  $\frac{2b^2}{a}$  (c)  $\frac{b^2}{a}$  (d)  $\frac{a^2}{b}$
36. The distance between the foci of a hyperbola is 16 and its eccentricity is  $\sqrt{2}$ , then the equation of hyperbola is  
 (a)  $x^2 + y^2 = 32$  (b)  $x^2 - y^2 = 16$  (c)  $x^2 + y^2 = 16$  (d)  $x^2 - y^2 = 32$

37. The equation of the hyperbola with vertices (3, 0) and (−3, 0) and semi-latus-rectum 4, is given by  
 (a)  $4x^2 - 3y^2 + 36 = 0$  (b)  $4x^2 - 3y^2 + 12 = 0$  (c)  $4x^2 - 3y^2 - 36 = 0$  (d) None of these
38. Equation of the hyperbola with eccentricity  $3/2$  and foci at  $(\pm 2, 0)$  is  
 (a)  $\frac{x^2}{4} - \frac{y^2}{5} = \frac{4}{9}$  (b)  $\frac{x^2}{9} - \frac{y^2}{9} = \frac{4}{9}$  (c)  $\frac{x^2}{4} - \frac{y^2}{9} = 1$  (d) None of these
39. The eccentricity of the hyperbola with latus rectum 12 and semi-conjugate axis  $2\sqrt{3}$ , is  
 (a) 2 (b) 3 (c)  $\frac{\sqrt{3}}{2}$  (d)  $2\sqrt{3}$
40. The eccentricity of the hyperbola  $3x^2 - 4y^2 = -12$  is  
 (a)  $\sqrt{\frac{7}{3}}$  (b)  $\frac{\sqrt{7}}{2}$  (c)  $-\sqrt{\frac{7}{3}}$  (d)  $-\frac{\sqrt{7}}{2}$
41. The equation  $\frac{x^2}{12-k} + \frac{y^2}{8-k} = 1$  represents  
 (a) A hyperbola if  $k < 8$  (b) An ellipse if  $k > 8$   
 (c) A hyperbola if  $8 < k < 12$  (d) None of these

## PARAMERIC EQUATIONS OF HYPERBOLA , SPECIAL FORMS OF HYPERBOLA

### Basic Level

42. The auxiliary equation of circle of hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , is  
 (a)  $x^2 + y^2 = a^2$  (b)  $x^2 + y^2 = b^2$  (c)  $x^2 + y^2 = a^2 + b^2$  (d)  $x^2 + y^2 = a^2 - b^2$
43. A point on the curve  $\frac{x^2}{A^2} - \frac{y^2}{B^2} = 1$  is  
 (a)  $(A\cos\theta, B\sin\theta)$  (b)  $(A\sec\theta, B\tan\theta)$  (c)  $(A\cos^2\theta, B\sin^2\theta)$  (d) None of these
44. The locus of the point of intersection of the lines  $ax\sec\theta + by\tan\theta = a$  and  $ax\tan\theta + by\sec\theta = b$ , where  $\theta$  is the parameter, is  
 (a) A straight line (b) A circle (c) An ellipse (d) A hyperbola
45. The eccentricity of the conic represented by  $x^2 - y^2 - 4x + 4y + 16 = 0$  is  
 (a) 1 (b)  $\sqrt{2}$  (c) 2 (d)  $1/2$
46. The latus rectum of the hyperbola  $9x^2 - 16y^2 - 18x - 32y - 151 = 0$  is  
 (a)  $\frac{9}{4}$  (b) 9 (c)  $\frac{3}{2}$  (d)  $\frac{9}{2}$
47. The vertices of a hyperbola are at  $(0, 0)$  and  $(10, 0)$  and one of its foci is at  $(18, 0)$ . The equation of the hyperbola is  
 (a)  $\frac{x^2}{25} - \frac{y^2}{144} = 1$  (b)  $\frac{(x-5)^2}{25} - \frac{y^2}{144} = 1$  (c)  $\frac{x^2}{25} - \frac{(y-5)^2}{144} = 1$  (d)  $\frac{(x-5)^2}{25} - \frac{(y-5)^2}{144} = 1$
48. The equations of the transverse and conjugate axis of the hyperbola  $16x^2 - y^2 + 64x + 4y + 44 = 0$  are  
 (a)  $x = 2, y + 2 = 0$  (b)  $x = 2, y = 2$  (c)  $y = 2, x + 2 = 0$  (d) None of these

49. Foci of the hyperbola  $\frac{x^2}{16} - \frac{(y-2)^2}{9} = 1$  are  
 (a) (5, 2), (-5, 2) (b) (5, 2), (5, -2) (c) (5, 2), (-5, -2) (d) None of these
50. The eccentricity of the conic  $x^2 - 2x - 4y^2 = 0$  is  
 (a)  $\frac{1}{4}$  (b)  $\frac{3}{2}$  (c)  $\frac{\sqrt{5}}{2}$  (d)  $\frac{\sqrt{5}}{4}$
51. The equation  $16x^2 - 3y^2 - 32x + 12y - 44 = 0$  represents a hyperbola  
 (a) The length of whose transverse axis is  $4\sqrt{3}$  (b) The length of whose conjugate axis is 4  
 (c) Whose centre is (-1, 2) (d) Whose eccentricity is  $\sqrt{\frac{19}{3}}$
52. The equation of the hyperbola whose foci are (6, 5), (-4, 5) and eccentricity  $\frac{5}{4}$  is  
 (a)  $\frac{(x-1)^2}{16} - \frac{(y-5)^2}{9} = 1$  (b)  $\frac{x^2}{16} - \frac{y^2}{9} = 1$  (c)  $\frac{(x-1)^2}{16} - \frac{(y-5)^2}{9} = -1$  (d) None of these
53. The equation  $x = \frac{e^t + e^{-t}}{2}; y = \frac{e^t - e^{-t}}{2}; t \in \mathbb{R}$  represents  
 (a) An ellipse (b) A parabola (c) A hyperbola (d) A circle
54. The vertices of the hyperbola  $9x^2 - 16y^2 - 36x + 96y - 252 = 0$  are  
 (a) (6, 3) and (-6, 3) (b) (6, 3) and (-2, 3) (c) (-6, 3) and (-6, -3) (d) None of these
55. The curve represented by  $x = a(\cosh\theta + \sinh\theta), y = b(\cosh\theta - \sinh\theta)$  is  
 (a) A hyperbola (b) An ellipse (c) A parabola (d) A circle
56. The foci of the hyperbola  $9x^2 - 16y^2 + 18x + 32y - 151 = 0$  are  
 (a) (2, 3), (5, 7) (b) (4, 1), (-6, 1) (c) (0, 0), (5, 3) (d) None of these

### Advance Level

57. The equations of the transverse and conjugate axes of a hyperbola respectively are  $x + 2y - 3 = 0$ ,  $2x - y + 4 = 0$  and their respective lengths are  $\sqrt{2}$  and  $\frac{2}{\sqrt{3}}$ . The equation of the hyperbola is  
 (a)  $\frac{2}{5}(x + 2y - 3)^2 - \frac{3}{5}(2x - y + 4)^2 = 1$  (b)  $\frac{2}{5}(2x - y + 4)^2 - \frac{3}{5}(x + 2y - 3)^2 = 1$   
 (c)  $2(2x - y + 4)^2 - 3(x + 2y - 3)^2 = 1$  (d)  $2(x + 2y - 3)^2 - 3(2x - y + 4)^2 = 1$
58. The points of intersection of the curves whose parametric equations are  $x = t^2 + 1, y = 2t$  and  $x = 2s, y = 2/s$  is given by  
 (a) (1, -3) (b) (2, 2) (c) (-2, 4) (d) (1, 2)
59. Equation  $\frac{1}{r} = \frac{1}{8} + \frac{3}{8}\cos\theta$  represents  
 (a) A rectangular hyperbola (b) A hyperbola (c) An ellipse (d) A parabola

## POSITION OF A POINT , TANGENTS , PAIR OF TANGENTS & DIRECTOR CIRCLE OF AN HYPERBOLA

### Basic Level

60. The line  $y = mx + c$  touches the curve  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , if
- (a)  $c^2 = a^2 m^2 + b^2$       (b)  $c^2 = a^2 m^2 - b^2$       (c)  $c^2 = b^2 m^2 - a^2$       (d)  $a^2 = b^2 m^2 + c^2$
61. The line  $lx + my + n = 0$  will be a tangent to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , if
- (a)  $a^2 l^2 + b^2 m^2 = n^2$       (b)  $a^2 l^2 - b^2 m^2 = n^2$       (c)  $a^2 m^2 - b^2 l^2 = a^2 l^2$       (d) None of these
62. If the straight line  $x \cos \alpha + y \sin \alpha = p$  be a tangent to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , then
- (a)  $a^2 \cos^2 \alpha + b^2 \sin^2 \alpha = p^2$       (b)  $a^2 \cos^2 \alpha - b^2 \sin^2 \alpha = p^2$   
 (c)  $a^2 \sin^2 \alpha + b^2 \cos^2 \alpha = p^2$       (d)  $a^2 \sin^2 \alpha - b^2 \cos^2 \alpha = p^2$
63. The equation of the tangent at the point  $(a \sec \theta, b \tan \theta)$  of the conic  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , is
- (a)  $x \sec \theta - y \tan \theta = 1$       (b)  $\frac{x}{a} \sec \theta - \frac{y}{b} \tan \theta = 1$   
 (c)  $\frac{x + a \sec \theta}{a^2} - \frac{y + b \tan \theta}{b^2} = 1$       (d) None of these
64. If the line  $y = 2x + \lambda$  be a tangent to the hyperbola  $36x^2 - 25y^2 = 3600$  then  $\lambda =$
- (a) 16      (b) -16      (c)  $\pm 16$       (d) None of these
65. The equation of the tangent to the hyperbola  $4y^2 = x^2 - 1$  at the point (1, 0) is
- (a)  $x = 1$       (b)  $y = 1$       (c)  $y = 4$       (d)  $x = 4$
66. The straight line  $x + y = \sqrt{2}p$  will touch the hyperbola  $4x^2 - 9y^2 = 36$ , is
- (a)  $p^2 = 2$       (b)  $p^2 = 5$       (c)  $5p^2 = 2$       (d)  $2p^2 = 5$
67. The equation of the tangent to the hyperbola  $2x^2 - 3y^2 = 6$  which is parallel to the line  $y = 3x + 4$ , is
- (a)  $y = 3x + 5$       (b)  $y = 3x - 5$       (c)  $y = 3x + 5$  and  $y = 3x - 5$       (d) None of these
68. The equation of tangents to the hyperbola  $3x^2 - 4y^2 = 12$  which cuts equal intercepts from the axes, are
- (a)  $y + x = \pm 1$       (b)  $y - x = \pm 1$       (c)  $3x + 4y = \pm 1$       (d)  $3x - 4y = \pm 1$
69. The line  $3x - 4y = 5$  is a tangent to the hyperbola  $x^2 - 4y^2 = 5$ . The point of contact is
- (a) (3, 1)      (b) (2, 1/4)      (c) (1, 3)      (d) None of these
70. The equation of a common tangent to the conics  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  and  $\frac{y^2}{a^2} - \frac{x^2}{b^2} = 1$ , is
- (a)  $x + y = a^2 - b^2$       (b)  $x + y = \sqrt{a^2 - b^2}$       (c)  $x - y = \sqrt{a^2 - b^2}$       (d)  $x + y = \sqrt{b^2 - a^2}$
71. The equation of common tangents to the parabola  $y^2 = 8x$  and hyperbola  $3x^2 - y^2 = 3$ , is
- (a)  $2x \pm y + 1 = 0$       (b)  $2x \pm y - 1 = 0$       (c)  $x \pm 2y + 1 = 0$       (d)  $x \pm 2y - 1 = 0$
72. The radius of the director circle of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , is
- (a)  $a - b$       (b)  $\sqrt{a - b}$       (c)  $\sqrt{a^2 - b^2}$       (d)  $\sqrt{a^2 + b^2}$

73. The tangents to the hyperbola  $x^2 - y^2 = 3$  are parallel to the straight line  $2x + y + 8 = 0$  at the following points.  
 (a) (2, 1) or (1, 2)      (b) (2, -1) or (-2, 1)      (c) (-1, -2)      (d) (-2, -1)
74. The line  $y = 4x + c$  touches the hyperbola  $x^2 - y^2 = 1$  iff  
 (a)  $c = 0$       (b)  $c = \pm\sqrt{2}$       (c)  $c = \pm\sqrt{15}$       (d)  $c = \pm\sqrt{17}$
75. The line  $5x + 12y = 9$  touches the hyperbola  $x^2 - 9y^2 = 9$  at the point  
 (a)  $\left(-5, \frac{4}{3}\right)$       (b)  $\left(5, -\frac{4}{3}\right)$       (c)  $\left(3, -\frac{1}{2}\right)$       (d) None of these
76. The number of tangents to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  from an external point is  
 (a) 2      (b) 4      (c) 6      (d) 5
77. The slope of the tangent to the hyperbola  $2x^2 - 3y^2 = 6$  at (3, 2) is  
 (a) -1      (b) 1      (c) 0      (d) 2
78. A common tangent to  $9x^2 - 16y^2 = 144$  and  $x^2 + y^2 = 9$  is  
 (a)  $y = \frac{3}{\sqrt{7}}x + \frac{\pi}{\sqrt{7}}$       (b)  $y = 3\sqrt{\frac{2}{7}}x + \frac{15}{\sqrt{7}}$       (c)  $y = 2\sqrt{\frac{3}{7}}x + 15\sqrt{7}$       (d) None of these
79. The product of the perpendiculars from two foci on any tangent to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$   
 (a)  $a^2$       (b)  $-a^2$       (c)  $b^2$       (d)  $-b^2$
80. If the two intersecting lines intersect the hyperbola and neither of them is a tangent to it, then number of intersecting points are  
 (a) 1      (b) 2      (c) 2, 3 or 4      (d) 2 or 3
81. The equation of a tangent parallel to  $y = x$  drawn to  $\frac{x^2}{3} - \frac{y^2}{2} = 1$  is  
 (a)  $x - y + 1 = 0$       (b)  $x + y + 2 = 0$       (c)  $x + y - 1 = 0$       (d)  $x - y + 2 = 0$
82. The equation of the tangent to the conic  $x^2 - y^2 - 8x + 2y + 11 = 0$  at (2, 1) is  
 (a)  $x + 2 = 0$       (b)  $2x + 1 = 0$       (c)  $x - 2 = 0$       (d)  $x + y + 1 = 0$
83. The equation of tangents to the hyperbola  $x^2 - 4y^2 = 36$  which are perpendicular to the line  $x - y + 4 = 0$   
 (a)  $y = -x + 3\sqrt{3}$       (b)  $y = -x - 3\sqrt{3}$       (c)  $y = -x \pm 2$       (d) None of these
84. The position of point (5, -4) relative to the hyperbola  $9x^2 - y^2 = 1$   
 (a) Outside the hyperbola (b) Inside the hyperbola (c) On the conjugate axis (d) On the hyperbola

### Advance Level

85. If the two tangents drawn on hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  in such a way that the product of their gradients is  $c^2$ , then they intersect on the curve
- (a)  $y^2 + b^2 = c^2(x^2 - a^2)$  (b)  $y^2 + b^2 = c^2(x^2 + a^2)$  (c)  $ax^2 + by^2 = c^2$  (d) None of these
86.  $C$  the centre of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ . The tangent at any point  $P$  on this hyperbola meets the straight lines  $bx - ay = 0$  and  $bx + ay = 0$  in the points  $Q$  and  $R$  respectively. Then  $CQ \cdot CR =$
- (a)  $a^2 + b^2$  (b)  $a^2 - b^2$  (c)  $\frac{1}{a^2} + \frac{1}{b^2}$  (d)  $\frac{1}{a^2} - \frac{1}{b^2}$
87. Let  $P(a \sec \theta, b \tan \theta)$  and  $Q(a \sec \phi, b \tan \phi)$ , where  $\theta + \phi = \frac{\pi}{2}$ , be two points on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ . If  $(h, k)$  is the point of intersection of the normals at  $P$  and  $Q$ , then  $k$  is equal to
- (a)  $\frac{a^2 + b^2}{a}$  (b)  $-\left(\frac{a^2 + b^2}{a}\right)$  (c)  $\frac{a^2 + b^2}{b}$  (d)  $-\left(\frac{a^2 + b^2}{b}\right)$
88. Let  $P$  be a point on the hyperbola  $x^2 - y^2 = a^2$  where  $a$  is a parameter such that  $P$  is nearest to the line  $y = 2x$ . The locus of  $P$  is
- (a)  $x - 2y = 0$  (b)  $2y - x = 0$  (c)  $x + 2y = 0$  (d)  $2y + x = 0$
89. An ellipse has eccentricity  $\frac{1}{2}$  and one focus at the point  $P\left(\frac{1}{2}, 1\right)$ . Its one directrix is the common tangent nearer to the point  $P$ , to the circle  $x^2 + y^2 = 1$  and the hyperbola  $x^2 - y^2 = 1$ . The equation of the ellipse in the standard form, is
- (a)  $\frac{(x-1/3)^2}{1/9} + \frac{(y-1)^2}{1/12} = 1$  (b)  $\frac{(x-1/3)^2}{1/9} + \frac{(y+1)^2}{1/12} = 1$
- (c)  $\frac{(x-1/3)^2}{1/9} - \frac{(y-1)^2}{1/12} = 1$  (d)  $\frac{(x-1/3)^2}{1/9} - \frac{(y+1)^2}{1/12} = 1$

### NORMALS , CO – NORMAL POINTS

#### Basic Level

90. The condition that the straight line  $lx + my = n$  may be a normal to the hyperbola  $b^2x^2 - a^2y^2 = a^2b^2$  is given by
- (a)  $\frac{a^2}{l^2} - \frac{b^2}{m^2} = \frac{(a^2 + b^2)^2}{n^2}$  (b)  $\frac{l^2}{a^2} - \frac{m^2}{b^2} = \frac{(a^2 + b^2)^2}{n^2}$  (c)  $\frac{a^2}{l^2} + \frac{b^2}{m^2} = \frac{(a^2 - b^2)^2}{n^2}$  (d)  $\frac{l^2}{a^2} + \frac{m^2}{b^2} = \frac{(a^2 - b^2)^2}{n^2}$
91. The equation of the normal to the hyperbola  $\frac{x^2}{16} - \frac{y^2}{9} = 1$  at  $(-4, 0)$  is
- (a)  $y = 0$  (b)  $y = x$  (c)  $x = 0$  (d)  $x = -y$
92. The equation of the normal at the point  $(a \sec \theta, b \tan \theta)$  of the curve  $b^2x^2 - a^2y^2 = a^2b^2$  is
- (a)  $\frac{ax}{\cos \theta} + \frac{by}{\sin \theta} = a^2 + b^2$  (b)  $\frac{ax}{\tan \theta} + \frac{by}{\sec \theta} = a^2 + b^2$  (c)  $\frac{ax}{\sec \theta} + \frac{by}{\tan \theta} = a^2 + b^2$  (d)  $\frac{ax}{\sec \theta} + \frac{by}{\tan \theta} = a^2 - b^2$
93. The number of normals to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  from an external point is
- (a) 2 (b) 4 (c) 6 (d) 5

## CHORD OF CONTACT , EQUATION OF THE CHORD WHOSE MID POINT IS GIVEN & EQUATION OF CHORD JOINING TWO POINTS

### Basic Level

94. The locus of the middle points of the chords of hyperbola  $3x^2 - 2y^2 + 4x - 6y = 0$  parallel to  $y = 2x$  is  
 (a)  $3x - 4y = 4$  (b)  $3y - 4x + 4 = 0$  (c)  $4x - 4y = 3$  (d)  $3x - 4y = 2$
95. The equation of the chord of the hyperbola  $x^2 - y^2 = 9$  which is bisected at  $(5, -3)$  is  
 (a)  $5x + 3y = 9$  (b)  $5x - 3y = 16$  (c)  $5x + 3y = 16$  (d)  $5x - 3y = 9$
96. If the chords of contact of tangents from two points  $(x_1, y_1)$  and  $(x_2, y_2)$  to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  are at right angles, then  $\frac{x_1 x_2}{y_1 y_2}$  is equal to  
 (a)  $-\frac{a^2}{b^2}$  (b)  $-\frac{b^2}{a^2}$  (c)  $-\frac{b^4}{a^4}$  (d)  $-\frac{a^4}{b^4}$
97. Equation of the chord of the hyperbola  $25x^2 - 16y^2 = 400$  which is bisected at the point  $(6, 2)$  is  
 (a)  $16x - 75y = 418$  (b)  $75x - 16y = 418$  (c)  $25x - 4y = 400$  (d) None of these

### Advance Level

98. If  $x = 9$  is the chord of contact of the hyperbola  $x^2 - y^2 = 9$ , then the equation of the corresponding pair of tangent is  
 (a)  $9x^2 - 8y^2 + 18x - 9 = 0$  (b)  $9x^2 - 8y^2 - 18x + 9 = 0$  (c)  $9x^2 - 8y^2 - 18x - 9 = 0$  (d)  $9x^2 - 8y^2 + 18x + 9 = 0$
99. If  $(a \sec \theta, b \tan \theta)$  and  $(a \sec \phi, b \tan \phi)$  are the ends of a focal chord of  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , then  $\tan \frac{\theta}{2} \tan \frac{\phi}{2}$  equals to  
 (a)  $\frac{e-1}{e+1}$  (b)  $\frac{1-e}{1+e}$  (c)  $\frac{1+e}{1-e}$  (d)  $\frac{e+1}{e-1}$
100. If  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  ( $a > b$ ) and  $x^2 - y^2 = c^2$  cut at right angles, then  
 (a)  $a^2 + b^2 = 2c^2$  (b)  $b^2 - a^2 = 2c^2$  (c)  $a^2 - b^2 = 2c^2$  (d)  $a^2 b^2 = 2c^2$
101. The locus of the middle points of the chords of contact of tangents to the hyperbola  $x^2 - y^2 = a^2$  from points on the auxiliary circle, is  
 (a)  $a^2(x^2 + y^2) = (x^2 - y^2)$  (b)  $a^2(x^2 + y^2) = (x^2 - y^2)^2$  (c)  $a^2(x^2 + y^2) = (x - y)^2$  (d) None of these
102. The locus of the mid points of the chords of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , which subtend a right angle at the origin  
 (a)  $\left(\frac{x^2}{a^2} - \frac{y^2}{b^2}\right)^2 \left(\frac{1}{a^2} - \frac{1}{b^2}\right) = \frac{x^2}{a^4} + \frac{y^2}{b^4}$  (b)  $\left(\frac{x^2}{a^2} - \frac{y^2}{b^2}\right)^2 \left(\frac{1}{a^2} - \frac{1}{b^2}\right) = \frac{x^2}{a^2} + \frac{y^2}{b^2}$   
 (c)  $\left(\frac{x^2}{a^2} - \frac{y^2}{b^2}\right) \left(\frac{1}{a^2} - \frac{1}{b^2}\right) = \frac{x^2}{a^2} + \frac{y^2}{b^2}$  (d) None of these



## POLE AND POLAR , DIAMETER OF POLAR POINTS

### Basic Level

103. The diameter of  $16x^2 - 9y^2 = 144$  which is conjugate to  $x = 2y$  is

- (a)  $y = \frac{16}{9}x$  (b)  $y = \frac{32}{9}x$  (c)  $x = \frac{16}{9}y$  (d)  $x = \frac{32}{9}y$

104. The lines  $2x + 3y + 4 = 0$  and  $3x - 2y + 5 = 0$  may be conjugate w.r.t the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ , if

- (a)  $a^2 + b^2 = \frac{10}{3}$  (b)  $a^2 - b^2 = \frac{10}{3}$  (c)  $b^2 - a^2 = \frac{10}{3}$  (d) None of these

105. The polars of  $(x_1, y_1)$  and  $(x_2, y_2)$  w.r.t  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  are perpendicular to each other if

- (a)  $\frac{x_1 x_2}{y_1 y_2} = -\frac{b^2}{a^4}$  (b)  $\frac{x_1 x_2}{y_1 y_2} = -\frac{a^4}{b^4}$  (c)  $x_1 x_2 + y_1 y_2 = \frac{a^2}{b^2}$  (d)  $x_1 x_2 - y_1 y_2 = \frac{a^2}{b^2}$

### Advance Level

106. The locus of the pole of normal chords of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  is

- (a)  $a^6 / x^2 - b^6 / y^2 = (a^2 + b^2)^2$  (b)  $x^2 / a^2 - y^2 / b^2 = (a^2 + b^2)^2$   
(c)  $a^2 / x^2 - b^2 / y^2 = (a^2 + b^2)^2$  (d) None of these

107. The locus of the pole with respect to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  of any tangent to the circle, whose diameter is the line joining the foci is the

- (a) Ellipse (b) Hyperbola (c) Parabola (d) None of these

## ASYMPTOTES OF HYPERBOLA

### Basic Level

108. The product of the lengths of perpendicular drawn from any point on the hyperbola  $x^2 - 2y^2 - 2 = 0$  to its asymptotes is

- (a)  $\frac{1}{2}$  (b)  $\frac{2}{3}$  (c)  $\frac{3}{2}$  (d) 2

109. The angle between the asymptotes of  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  is equal to

- (a)  $2\tan^{-1}\left(\frac{b}{a}\right)$  (b)  $2\tan^{-1}\frac{a}{b}$  (c)  $\tan^{-1}\frac{a}{b}$  (d)  $\tan^{-1}\frac{b}{a}$

### Advance Level

110. The product of perpendicular drawn from any point on a hyperbola to its asymptotes is

- (a)  $\frac{a^2 b^2}{a^2 + b^2}$  (b)  $\frac{a^2 + b^2}{a^2 b^2}$  (c)  $\frac{ab}{\sqrt{a} + \sqrt{b}}$  (d)  $\frac{ab}{a^2 + b^2}$

111. From any point on the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  tangents are drawn to the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 2$ .

The area cut-off by the chord of contact on the asymptotes is equal to

- (a)  $\frac{ab}{2}$  (b)  $ab$  (c)  $2ab$  (d)  $4ab$

112. The equation of the hyperbola whose asymptotes are the straight lines  $3x - 4y + 7 = 0$  and  $4x + 3y + 1 = 0$  and which passes through origin is

- (a)  $(3x - 4y + 7)(4x + 3y + 1) = 0$  (b)  $12x^2 - 7xy - 12y^2 + 31x + 17y = 0$   
(c)  $12x^2 - 7xy + 2y^2 = 0$  (d) None of these

113. The equation of the asymptotes of the hyperbola  $2x^2 + 5xy + 2y^2 - 11x - 7y - 4 = 0$  are

- (a)  $2x^2 + 5xy + 2y^2 - 11x - 7y - 5 = 0$  (b)  $2x^2 + 4xy + 2y^2 - 7x - 11y + 5 = 0$   
(c)  $2x^2 + 5xy + 2y^2 - 11x - 7y + 5 = 0$  (d) None of these

## RECTANGULAR HYPERBOLA

### Basic Level

114. Eccentricity of the curve  $x^2 - y^2 = a^2$  is

- (a) 2 (b)  $\sqrt{2}$  (c) 4 (d) None of these

115. The eccentricity of curve  $x^2 - y^2 = 1$  is

- (a)  $\frac{1}{2}$  (b)  $\frac{1}{\sqrt{2}}$  (c) 2 (d)  $\sqrt{2}$

116. The eccentricity of the hyperbola  $x^2 - y^2 = 25$  is

- (a)  $\sqrt{2}$  (b)  $\frac{1}{\sqrt{2}}$  (c) 2 (d)  $1 + \sqrt{2}$

117. If transverse and conjugate axes of a hyperbola are equal, then its eccentricity is

- (a)  $\sqrt{3}$  (b)  $\sqrt{2}$  (c)  $\frac{1}{\sqrt{2}}$  (d) 2

118. The eccentricity of the hyperbola  $\frac{\sqrt{1999}}{3}(x^2 - y^2) = 1$  is

- (a)  $\sqrt{3}$  (b)  $\sqrt{2}$  (c) 2 (d)  $2\sqrt{2}$

119. Eccentricity of the rectangular hyperbola  $\int_0^1 e^x \left( \frac{1}{x} - \frac{1}{x^3} \right) dx$  is

- (a) 2 (b)  $\sqrt{2}$  (c) 1 (d)  $\frac{1}{\sqrt{2}}$

120. The reciprocal of the eccentricity of rectangular hyperbola, is

- (a) 2 (b)  $\frac{1}{2}$  (c)  $\sqrt{2}$  (d)  $\frac{1}{\sqrt{2}}$

121. The locus of the point of intersection of the lines  $(x + y)t = a$  and  $x - y = at$ , where  $t$  is the parameter, is

- (a) A circle (b) An ellipse  
(c) A rectangular hyperbola (d) None of these

122. Curve  $xy = c^2$  is said to be

- (a) Parabola (b) Rectangular hyperbola (c) Hyperbola (d) Ellipse

123. What is the slope of the tangent line drawn to the hyperbola  $xy = a (a \neq 0)$  at the point  $(a, 1)$
- (a)  $\frac{1}{a}$  (b)  $-\frac{1}{a}$  (c)  $a$  (d)  $-a$
124. The coordinates of the foci of the rectangular hyperbola  $xy = c^2$  are
- (a)  $(\pm c, c)$  (b)  $(\pm c\sqrt{2}, \pm c\sqrt{2})$  (c)  $\left(\pm \frac{c}{\sqrt{2}}, \pm \frac{c}{\sqrt{2}}\right)$  (d) None of these
125. A tangent to a hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  intercepts a length of unity from each of the coordinate axes, then the point  $(a, b)$  lies on the rectangular hyperbola
- (a)  $x^2 - y^2 = 2$  (b)  $x^2 - y^2 = 1$  (c)  $x^2 - y^2 = -1$  (d) None of these
126. A rectangular hyperbola is one in which
- (a) The two axes are rectangular (b) The two axes are equal  
(c) The asymptotes are perpendicular (d) The two branches are perpendicular
127. If  $e$  and  $e_1$  are the eccentricities of the hyperbolas  $xy = c^2$  and  $x^2 - y^2 = c^2$ , then  $e^2 + e_1^2$  is equal to
- (a) 1 (b) 4 (c) 6 (d) 8
128. If the line  $ax + by + c = 0$  is a normal to the curve  $xy = 1$ , then
- (a)  $a > 0, b > 0$  (b)  $a > 0, b < 0$  or  $a < 0, b > 0$  (c)  $a < 0, b < 0$  (d) None of these
129. The number of normals that can be drawn from any point to the rectangular hyperbola  $xy = c^2$  is
- (a) 1 (b) 2 (c) 3 (d) 4
130. The equation of the chord joining two points  $(x_1, y_1)$  and  $(x_2, y_2)$  on the rectangular hyperbola  $xy = c^2$  is
- (a)  $\frac{x}{x_1 + x_2} + \frac{y}{y_1 + y_2} = 1$  (b)  $\frac{x}{x_1 - x_2} + \frac{y}{y_1 - y_2} = 1$  (c)  $\frac{x}{y_1 + y_2} + \frac{y}{x_1 + x_2} = 1$  (d)  $\frac{x}{y_1 - y_2} + \frac{y}{x_1 - x_2} = 1$
131. If a triangle is inscribed in a rectangular hyperbola, its orthocentre lies
- (a) Inside the curve (b) Outside the curve (c) On the curve (d) None of these
- Advance Level**
132. The equation of the common tangent to the curves  $y^2 = 8x$  and  $xy = -1$  is
- (a)  $3y = 9x + 2$  (b)  $y = 2x + 1$  (c)  $2y = x + 8$  (d)  $y = x + 2$
133. A rectangular hyperbola whose centre is  $C$  is cut by any circle of radius  $r$  in four points  $P, Q, R$  and  $S$ , then  $CP^2 + CQ^2 + CR^2 + CS^2 =$
- (a)  $r^2$  (b)  $2r^2$  (c)  $3r^2$  (d)  $4r^2$
134. If  $P(x_1, y_1), Q(x_2, y_2), R(x_3, y_3)$  and  $S(x_4, y_4)$  are four concyclic points on the rectangular hyperbola  $xy = c^2$ , the coordinates of orthocentre of the  $\Delta PQR$  are
- (a)  $(x_4, -y_4)$  (b)  $(x_4, y_4)$  (c)  $(-x_4, -y_4)$  (d)  $(-x_4, y_4)$
135. If a circle cuts the rectangular hyperbola  $xy = 1$  in the points  $(x_r, y_r)$  where  $r = 1, 2, 3, 4$  then
- (a)  $x_1 x_2 x_3 x_4 = 2$  (b)  $x_1 x_2 x_3 x_4 = 1$  (c)  $x_1 + x_2 + x_3 + x_4 = 0$  (d)  $y_1 + y_2 + y_3 + y_4 = 0$

**ANSWER**  
**ASSIGNMENT ( BASIC & ADVANCE LEVEL )**

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
c	d	d	a	c	c	d	b	d	b	c	a	d	a	b	a	d	a	a	a
21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
a	b	c	c	c	c	b	a	a	b	a	b	c	a,d	a	d	c	a	a	a
41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
c	a	b	d	b	d	b	c	a	c	d	a	c	b	a	b	b	b	b	b
61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
b	b	b	c	a	d	c	b	a	b	a	c	b	c	b	a	b	b	c	c
81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
a	c	a,b	a	a	a	d	a,b	a	a	a	c	b	a	c	d	b	b	b	c
101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120
b	a	b	a	b	a	a	b	a	a	d	b	c	b	d	a	b	b	b	d
121	122	123	124	125	126	127	128	129	130	131	132	133	134	135					
c	b	b	b	b	a,b, c	b	b	d	a	c	d	d	d	b					