



SESSION LEAD CARD AND EMERGENCY ACTION PLAN



Club Name:	United Cornwall FC
Club Address:	Evolve Fitness, Cambrone College,
Postcode:	Trevenson Ln, TR15, Pool, Redruth TR15 3RE
Telephone:	07460118199 (Jack Greaves)

FIRST AIDERS / HELPER INFORMATION

Name	Mobile Number / Email
For queries about this EAP:	
Jack Greaves	07460118199 / jack@growcornwall.com
All Session Leads should hold suitable First Aid Qualification as outlined in the Risk Assessments. There should be a minimum of one First Aider at each session, who is responsible for bringing suitable kit and knowing the location specific information.	

FIRST AID EQUIPMENT AND FACILITIES

Item	Location
Defibrillator	Side of the pitch (on the building)
Stretcher	N/A
First Aid Room	Rooms available next to the pitch if required

ACCESS ROUTES

For Ambulance	Evolve Fitness, across from Pool School entrance
From Pitch to Ambulance	You can park above or alongside the pitch - access gates at bottom

OTHER INFORMATION

Nearest A&E / Trauma Hospital:	Treliske Truro Cornwall TR1 3LQ. Phone. 01872 250000.
---	---



SESSION LEAD CARD AND EMERGENCY ACTION PLAN



CHARTER STANDARD
CLUB

Fastest Route to A&E / Trauma Hospital:	
Distance & Journey Time:	19 minutes (without traffic)
Nearest Walk-in Centre:	Barncoose - Camborne - not 24 hours
Alternative Trauma Hospital:	N/A


Additional Information	
Timings	Monday evenings - 5-7pm
Toilets	Toilets are found in the building next to the pitch.
Changing Rooms	Changing Rooms are also available in the building next to the pitch
Parking for participants	There is parking next to the pitch for participants
Location Lead	Trudy Watts



SESSION LEAD CARD AND EMERGENCY ACTION PLAN



CHARTER STANDARD
CLUB






SESSION LEAD CARD AND EMERGENCY ACTION PLAN



CHARTER STANDARD
CLUB

Adult basic life support in community settings





SESSION LEAD CARD AND EMERGENCY ACTION PLAN



CHARTER STANDARD
CLUB

CRT6™



Concussion Recognition Tool

To Help Identify Concussion in Children, Adolescents and Adults

What is the Concussion Recognition Tool?

A concussion is a brain injury. The Concussion Recognition Tool 6 (CRT6) is to be used by non-medically trained individuals for the identification and immediate management of suspected concussion. It is not designed to diagnose concussion.

Recognise and Remove

Red Flags: CALL AN AMBULANCE

If **ANY** of the following signs are observed or complaints are reported after an impact to the head or body the athlete should be immediately removed from play/game/activity and transported for urgent medical care by a healthcare professional (HCP):

- Neck pain or tenderness
- Seizure, 'fits', or convulsion
- Loss of vision or double vision
- Loss of consciousness
- Increased confusion or deteriorating conscious state (becoming less responsive, drowsy)
- Weakness or numbness/tingling in more than one arm or leg
- Repeated Vomiting
- Severe or increasing headache
- Increasingly restless, agitated or combative
- Visible deformity of the skull

Remember

- In all cases, the basic principles of first aid should be followed: assess danger at the scene, check airway, breathing, circulation; look for reduced awareness of surroundings or slowness or difficulty answering questions.
- Do not attempt to move the athlete (other than required for airway support) unless trained to do so.
- Do not remove helmet (if present) or other equipment.
- Assume a possible spinal cord injury in all cases of head injury.
- Athletes with known physical or developmental disabilities should have a lower threshold for removal from play.

If there are no Red Flags, identification of possible concussion should proceed as follows:

Concussion should be suspected after an impact to the head or body when the athlete seems different than usual. Such changes include the presence of **any one or more** of the following: visible clues of concussion, signs and symptoms (such as headache or unsteadiness), impaired brain function (e.g. confusion), or unusual behaviour.

This tool may be freely copied in its current form for distribution to individuals, teams, groups, and organizations. Any alteration (including translations and digital re-formatting), re-branding, or sale for commercial gain is not permissible without the expressed written consent of BMJ.



SESSION LEAD CARD AND EMERGENCY ACTION PLAN



CHARTER STANDARD
CLUB

Concussion Recognition Tool 6 - CRT6™



CRT6

Concussion Recognition Tool

To Help Identify Concussion in Children, Adolescents and Adults



1: Visible Clues of Suspected Concussion

Visible clues that suggest concussion include:

- Loss of consciousness or responsiveness
- Lying motionless on the playing surface
- Falling unprotected to the playing surface
- Disorientation or confusion, staring or limited responsiveness, or an inability to respond appropriately to questions
- Dazed, blank, or vacant look
- Seizure, fits, or convulsions
- Slow to get up after a direct or indirect hit to the head
- Unsteady on feet / balance problems or falling over / poor coordination / wobbly
- Facial injury

2: Symptoms of Suspected Concussion

Physical Symptoms	Changes in Emotions
Headache	More emotional
"Pressure in head"	More irritable
Balance problems	Sadness
Nausea or vomiting	Nervous or anxious
Drowsiness	
Dizziness	
Blurred vision	
More sensitive to light	
More sensitive to noise	
Fatigue or low energy	
"Don't feel right"	
Neck Pain	

Changes in Thinking
Difficulty concentrating
Difficulty remembering
Feeling slowed down
Feeling like "in a fog"

Remember, symptoms may develop over minutes or hours following a head injury.

3: Awareness

(Modify each question appropriately for each sport and age of athlete)

Failure to answer any of these questions correctly may suggest a concussion:

"Where are we today?"

"What event were you doing?"

"Who scored last in this game?"

"What team did you play last week/game?"

"Did your team win the last game?"

Any athlete with a suspected concussion should be - IMMEDIATELY REMOVED FROM PRACTICE OR PLAY and should NOT RETURN TO ANY ACTIVITY WITH RISK OF HEAD CONTACT, FALL OR COLLISION, including SPORT ACTIVITY until ASSESSED MEDICALLY, even if the symptoms resolve.

Athletes with suspected concussion should **NOT**:

- Be left alone initially (at least for the first 3 hours). Worsening of symptoms should lead to immediate medical attention.
- Be sent home by themselves. They need to be with a responsible adult.
- Drink alcohol, use recreational drugs or drugs not prescribed by their HCP
- Drive a motor vehicle until cleared to do so by a healthcare professional