

FEMALE HEALTH ASSESSMENT QUESTIONNAIRE

NAME: _____ EMAIL: _____

TODAY'S DATE: _____ PHONE: _____

Please mark the appropriate box for each symptom you may be experiencing.

SYMPTOMS	NONE	MILD	MODERATE	SEVERE	VERY SEVERE
Physical Exhaustion (fatigue, lack of energy, stamina or motivation)					
Sleep Problems (difficulty falling asleep or sleeping through the night)					
Irritability (mood swings, feeling aggressive, angers easily)					
Anxiety (feeling overwhelmed, feeling panicky, or feeling nervous)					
Decline in drive or interest (loss of "zest for life," feeling down or sad)					
Joint and muscular symptoms (joint pain, muscle weakness, poor recovery after exercise)					
Difficulties with memory (concentration, finding the right word, or retaining information)					
Vaginal dryness or difficulty with sexual intercourse					
Sexual Problems (change in desire, activity, orgasm and/or satisfaction)					
Sweating (night sweats or increased episodes of sweating)					
Hot Flashes (burst that starts in chest and lasts for short duration)					
Hair loss, thinning or change in texture of hair					
Feeling cold all the time, having cold hands or feet					
Headaches or migraines (increase in frequency or intensity)					
Weight (difficulty losing weight despite diet/exercise)					
Bladder problems (difficulty in urinating, increased need to urinate, incontinence)					

Other symptoms or unique health circumstances to take into consideration:

FEMALE HEALTH HISTORY & SYMPTOMS

For CDSS Continuous Round

PATIENT INFORMATION

Name: _____ Date: _____
 Date of Birth: _____ Age: _____ Weight: _____ Height: _____

PATIENT QUESTIONS

Have you been diagnosed with any cancer since initial pelleting (excluding basal cell carcinoma)?	Yes	No		
Blood clot, DVT, heart attack or stroke since being pelleted?	Yes	No		
Currently pregnant or trying to conceive?	Yes	No		
Had a recent mammogram (within last 12 months)?	Yes	No		
Have you had a hysterectomy since last insertion?	Yes	No		
If so, type of hysterectomy:	Complete (uterus and ovaries removed)	Partial (uterus only removed)		
Had menstrual cycle (within last 12 months)?	Yes	No		
Had endometrial ablation?	Yes	No		
Have you had any spotting or bleeding since last pellet?	Yes	No		
Are you on birth control?	Yes	No		
Name of birth control: _____				
Select types of hormones you are currently on:	Testosterone	Estrogen	Progesterone	Thyroid
Are you currently on statins?	Yes	No		
Are you a smoker?	Yes	No		
Are you currently on oral nitrates?	Yes	No		

PATIENT'S CURRENT AND PAST MEDICAL HISTORY

Select all that apply:

Cardiovascular Conditions:

- Tachycardia
- Hypertension
- Hyperlipidemia
- Obstructive Sleep Apnea
- Atrial Fibrillation

Neurological Conditions:

- Epilepsy or Seizure Disorder
- Depression/Anxiety
- Psychiatric Conditions
- Migraine with Aura
- Meningioma (since last pellet)

Gynecological Conditions:

- Pre-Menstrual Syndrome
- Endometriosis
- Fibrocystic Breast Disease
- Fibroids (since last pellet)
- Polyps (since last pellet)

FEMALE HEALTH HISTORY & SYMPTOMS

For CDSS Continuous Round

MEDICAL HISTORY

Endocrine and Metabolic:

PCOS
Diabetes Type 2 or Insulin Resistance
Hyperthyroid
Hypothyroid
Multiple Endocrine Neoplasia Type-2

Autoimmune Conditions:

Diabetes Type 1
Hashimoto's Thyroiditis
Graves' Disease
Rheumatoid Arthritis
Multiple Sclerosis
Systemic Lupus (Erythematosus)
Psoriasis
IBS (Irritable Bowel Syndrome)
Crohn's Disease
Ulcerative Colitis

Organ Specific Conditions:

Liver Disease (since last pellet)
Kidney Disease (since last pellet)
LAM (Lymphangioliomyomatosis)
Osteoporosis or Osteopenia
HIV
Hepatitis
Hemochromatosis
Pancreatitis (since last pellet)
History of or Gall Bladder Disease
Polycythemia Vera (PV)

SYMPTOMS AND CONCERNS

Select all that apply:

Hot Flashes	Cold Hands or Feet
Night Sweats	Brittle Nails
Vaginal Dryness	Dry or Flaking Skin
Decreased Interest in Sex	Lack of Energy (Fatigue)
Inability To or Delayed Orgasm	Decreased Muscle Mass
Painful Intercourse	Acne
Urinary Incontinence	Facial Hair
Frequent Urinary Tract Infection	Dry Eyes
Breast Tenderness	Joint Pain
Weight Gain	Difficulty Sleeping
Hair Loss	Mind Racing at Bedtime
Hair Thinning	Eating When Stressed
Thinning Eyebrows	

Female Hormone Replacement Insertion Consent

Name: _____ Date of Birth: _____

My physician/practitioner has recommended bioidentical hormone therapy delivered by a pellet inserted under my skin for treatment of symptoms I am experiencing related to low hormone levels. The following information has been explained to me prior to receiving the recommended therapy.

OVERVIEW

Bioidentical hormones are hormones that are biologically identical to that made in my own body. The levels of active estradiol and/or testosterone made by my body have decreased, and therapy using these hormones may have the same or similar effect(s) on my body as my own naturally produced hormones. The pellets are a delivery mechanism for estradiol and/or testosterone, and bioidentical hormone replacement therapy using pellets has been used since the 1930's. There are other formulations of estradiol and testosterone replacement available, and different methods can be used to deliver the therapy. There are no commercially available forms of testosterone that are formulated specifically for use in females. The risks associated with pellet therapy are generally similar to other forms of replacement therapy using bioidentical hormones.

PELLET ACTIVE INGREDIENTS

I understand that (please initial by the appropriate statement):

- I am receiving pellets today that contain testosterone only.
 I am receiving pellets today that contain estradiol and testosterone.
 I am receiving pellets today that contain testosterone and anastrozole.

RISKS/COMPLICATIONS

Risks associated with pellet insertion may include: bleeding from incision site, bruising, fever, infection, pain, swelling, scarring, keloid formation at incision site, vaginal engorgement, pellet extrusion which may occur several weeks or months after insertion, reaction to local anesthetic and/or preservatives, allergy to adhesives from bandage(s), steri strips or other adhesive agents.

Some individuals may experience one or more of the following side effects with testosterone: acne, abnormal bleeding or a change in menstrual cycle (if patient has a uterus), anxiety, breast or nipple tenderness or swelling, insomnia, depression, mood swings, fluid retention, headaches, excess facial hair, hair thinning and/or loss, voice deepening, clitoromegaly, rash, redness, itching, lack of effect hypersexuality (overactive libido) or decreased libido, overproduction of estrogen (called aromatization) or an increase in red blood cell formation or blood count (erythrocytosis). The latter may be diagnosed with a blood test called a complete blood count (CBC). This test should be done at least annually. Erythrocytosis can be reversed simply by donating blood periodically, but further workup or referral may be required if a more worrisome condition is suspected. If you are planning to start or expand your family soon, please talk to your provider about other non-testosterone options. In addition, testosterone may cause some breast cancers to grow more rapidly.

RISKS/COMPLICATIONS OF ESTRADIOL (Only applicable if receiving estradiol in the pellets)

The side-effects of estradiol are similar to those listed above for testosterone. Additionally, there is some risk, even when using bioidentical hormones, that estrogens may cause existing cases of some breast cancers to grow more rapidly. This risk may also apply to some undiagnosed forms of breast cancer.

I ACKNOWLEDGE THAT I HAVE RECEIVED A COPY AND UNDERSTAND THE INFORMATION ON THIS FORM

PATIENT:

Print Name: _____

Signature: _____ Date: _____

WITNESS:

Print Name: _____

Signature: _____ Date: _____

Using estrogen-alone (without progesterone) may increase the chance of getting cancer of the uterus. Endometrial sampling (biopsy) or surgery may be required if abnormal bleeding occurs.

Please initial if you are postmenopausal, have a uterus, and are getting estradiol.

I understand that I have a uterus and am receiving postmenopausal dosing of estradiol. I agree to take progesterone as directed by my health care provider while receiving estradiol.

RISKS/COMPLICATIONS OF ANASTROZOLE (Only applicable if receiving anastrozole in the pellets)

Anastrozole is a type of medication called an aromatase inhibitor. Aromatase inhibitors limit or prevent the conversion of testosterone into estrogen. Aromatase inhibitors can be used for a variety of conditions but are most commonly used in patients with a history of estrogen receptor positive breast cancer.

Anastrozole should not be used in pregnant women and should be used with caution in females with pre-existing ischemic heart disease. Anastrozole in pellets should not be given to premenopausal women nor to women taking oral aromatase inhibitors (anastrozole or letrozole) or selective estrogen receptor modulators (tamoxifen or raloxifene).

The amount of anastrozole used in pellets is very low. The most common side-effects for women taking anastrozole are hot flashes, joint pain, and muscle pain. Because of the low dose in the pellet, these effects are not usually seen with this type of therapy, however.

CONSENT FOR TREATMENT

I agree to immediately report any adverse reactions or problems that may be related to my therapy to my physician or health care provider's office, so that it may be reported to the manufacturer. Potential complications have been explained to me, and I acknowledge that I have received and understand this information, including the possible risks and potential complications and the potential benefits.

I also acknowledge that the nature of bioidentical therapy and other treatments have been explained to me, and I have had all my questions answered. Blood tests may be necessary on several occasions during the 1st year to help w/ dosing, and then annually or biannually at the discretion of the prescribing practitioner.

I understand that my blood tests may reveal that my levels are not optimal which, would mean I may need a higher or lower dose in the future. Furthermore, I have not been promised or guaranteed any specific benefits from the insertion of testosterone pellets.

I accept these risks and benefits, and I consent to the insertion of testosterone pellets under my skin performed by my provider. This consent is ongoing for this and all future insertions in this facility until I am no longer a patient here, but I do understand that I can revoke my consent at any time. I have been informed that I may experience any of the complications to this procedure as described above. I have read or have had this form read to me.

Female Post-Insertion Instructions

Name: _____ Date of Birth: _____

- Your insertion site has been covered with two layers of bandages. Keep both bandages on for 3 days.
- You may experience bruising, swelling, and/or redness of the insertion site which may last from a few days up to 2 to 3 weeks. If the redness worsens after the first 2-3 days, please contact the office.
- Do not take tub baths or get into a hot tub or swimming pool for 3-4 days. You may shower, but do not remove the bandage or steri-strips for 4 days.
- No heavy lifting or major exercises for the incision area for the next 3-4 days, which includes running, elliptical, squats, lunges, etc.
- The sodium bicarbonate in the anesthetic may cause the site to swell for 1-3 days.
- The insertion site may be uncomfortable for up to 2 to 3 weeks. If there is itching or redness contact the office.
- You may notice some pinkish or bloody discoloration of the outer bandage. This is normal.
- If you experience bleeding from the incision, apply firm pressure for 5 minutes.
- Please call if you have any bleeding not relieved with pressure (not oozing), as this is NOT normal.
- Please call if you have any pus coming out of the insertion site, as this is NOT normal.
- Please call if the area becomes red and warm to the touch.
- We recommend putting an ice pack on the area where the pellets are located a couple of times for about 20 minutes each time over the next 4 to 5 hours. You can continue this for swelling, if needed. Be sure to place something between the ice pack and your bandages/skin. Do not place ice packs directly on bare skin.

REMINDERS

- Remember to have your post-insertion blood work done 1-2 weeks prior to your next insertion. If you are not feeling any better by 4 weeks, however, please call the office to have your labs drawn early.
- It may take 2-3 rounds for some to see symptom improvement.
- Most women will need re-insertion of their pellets 3-4 months after their initial insertion. If you experience symptoms prior to this, please call the office.
- Please call as soon as symptoms that were relieved from the pellets start to return to make an appointment for your next insertion.

ADDITIONAL INSTRUCTIONS

I ACKNOWLEDGE THAT I HAVE RECEIVED A COPY AND UNDERSTAND THE INFORMATION ON THIS FORM

PATIENT:

Print Name: _____

Signature: _____ Date: _____

WITNESS:

Print Name: _____

Signature: _____ Date: _____