



Particle in a Two-Dimensional Triangular Box: A Complete Theoretical Analysis

- 1) **Himanshu Jain**, Assistant Professor, Head, Department of Chemistry, Desh Bhagat College, Bardwal-Dhuri

Abstract

Quantum confinement in non-rectangular geometries has attracted considerable interest due to its significant applications in nanotechnology, condensed matter physics, and quantum computing. This paper presents a complete theoretical analysis of a non-relativistic particle confined within a two-dimensional equilateral triangular infinite potential well. We systematically derive the solutions to the time-independent Schrödinger equation by employing an appropriate coordinate transformation and imposing strict boundary conditions along the triangle's edges. The eigenfunctions are constructed using sinusoidal components adapted to triangular boundaries, leading to a quantized energy spectrum that markedly differs from those of rectangular or circular wells. The resulting energy levels depend on two quantum numbers with a specific constraint reflecting the triangular geometry. We analyze the properties of eigenstates, their nodal patterns, and the influence of geometric symmetry on probability distributions. Furthermore, we discuss the physical relevance of such systems in modern applications, including graphene quantum dots, triangular nanomechanical resonators, and molecular clusters exhibiting triangular confinement. Our study not only enriches the fundamental understanding of quantum particles in reduced-symmetry domains but also provides a foundation for investigating perturbations, external field effects, and potential many-body interactions within triangular geometries. This work bridges theoretical concepts with experimental possibilities, offering new avenues for future research.

Keywords: *Quantum Confinement, Oblique Coordinates, Equilateral Triangle Potential well, Schrodinger Equation in Non-orthogonal coordinates.*

Quantum mechanics has profoundly transformed our understanding of the microscopic world, especially through the study of systems where particles are confined in restricted geometries. Traditional models such as the particle in a one-dimensional box, two-dimensional square well, and circular well have been extensively studied both for their theoretical significance and practical applications. However, less attention has been given to confinement in non-standard geometries, such as triangular domains, despite their growing relevance in emerging quantum technologies. Triangular quantum wells, especially equilateral triangular ones, represent a fascinating playground where symmetry and boundary conditions interplay uniquely to affect the energy eigenvalues and eigenfunctions of confined particles.

In recent years, advancements in nanofabrication techniques have allowed the realization of quantum dots and nanostructures with triangular shapes, particularly in materials like graphene and semiconductor heterostructures. These systems exhibit novel optical, electronic, and mechanical properties, many of which originate from the peculiar nature of quantum confinement in a triangle. The triangular confinement not only introduces complexity in the

solution of the Schrödinger equation but also results in characteristic energy level spacings and nodal structures that differ from those observed in rectangular or circular geometries.


The present work focuses on a detailed theoretical treatment of a non-relativistic particle confined in a two-dimensional equilateral triangular infinite potential well. We derive the exact form of eigenfunctions and eigenvalues, discuss their symmetry properties, and highlight distinctive physical consequences of triangular confinement. Our approach lays the groundwork for understanding more complex phenomena such as electron interactions, external field perturbations, and spin-related effects in triangular geometries. By providing an analytical framework, this study aims to contribute to the fundamental body of knowledge in quantum chemistry and quantum physics while paving the way for practical applications in nanoelectronics, quantum information processing, and materials science.

Mathematical Formulation

Problem Statement

Consider a particle of mass (m) confined within an equilateral triangular domain of side length (a). The potential energy $V(x, y)$ is defined as:

$V(x, y) = 0$ inside the triangle; $V(x, y) = \infty$ outside the triangle.



The time-independent Schrödinger equation governing the system is:

$$-(\hbar^2/2m)\nabla^2\Psi(x, y) = E\Psi(x, y)$$

with boundary condition $\Psi = 0$ along the edges of the triangle.

Choice of Coordinate System

Define oblique coordinates (u, v):

$$u = x,$$

$$v = (-x/2) + (\sqrt{3}/2)y.$$

The triangle is bounded by $u = 0$, $v = 0$, and $u + v = a$.

Ansatz for Solution

We propose:

$$\Psi_{(u, v)} = \sin(n\pi u/a) \sin(m\pi v/a)$$

where n, m are positive integers.

Application of Boundary Conditions

Boundary condition at $u + v = a$ implies:
 $n + m = p$ (p is an integer).

Derivation of Energy Eigenvalues

Operating the Laplacian yields:

$$E(n, m) = (\hbar^2 \pi^2 / 2ma^2)(n^2 + m^2 - nm).$$

Analysis of Eigenfunctions

The eigenfunctions $\Psi_{(n, m)(u, v)} = \sin(n\pi u/a) \sin(m\pi v/a)$ are subject to $n + m = p$.

The nodal lines form slanted patterns, with the probability density exhibiting triangular symmetry.

Properties of Energy Spectrum

- Degeneracy
Accidental degeneracies occur when distinct (n, m) yield the same $E(n, m)$, but they are rare due to the constraint $n+m=p$.
- Energy Ordering
For a given p , the lowest energy corresponds to $n \approx m$.

The theoretical framework of a particle confined in a two-dimensional equilateral triangular box finds multiple applications across modern physical and chemical systems. In nanotechnology, triangular quantum dots fabricated from materials like graphene exhibit unique electronic properties such as valley polarization, quantum interference patterns, and edge-dependent conductivity. These triangular confinements allow precise control over the electronic states, making them suitable candidates for quantum computing and information storage.

In optoelectronics, triangular nanostructures enhance light-matter interactions due to their sharp corners and confined modes, leading to the development of highly sensitive photodetectors and nanoscale lasers. Additionally, the study of vibrational modes in triangular membranes serves as a prototype for understanding mechanical resonances in nanomechanical resonators and sensors.

Molecular chemistry also benefits from triangular confinement models. Certain molecular clusters, such as metal trimers, naturally exhibit triangular symmetries, and understanding their quantum mechanical behavior is crucial for interpreting spectroscopic data and designing new catalytic materials. Furthermore, the principles derived from triangular quantum systems can extend to designing artificial molecules with tailored electronic properties.

Thus, beyond pure theoretical interest, the study of particles in triangular domains is fundamentally linked to advancements in nanoelectronics, photonics, molecular engineering, and quantum device fabrication, opening avenues for future experimental and technological innovations.

Conclusion

In this study, we have presented a comprehensive theoretical analysis of a non-relativistic quantum particle confined within a two-dimensional equilateral triangular infinite potential well. By solving the time-independent Schrödinger equation under the imposed boundary conditions, we derived explicit expressions for the eigenfunctions and corresponding energy

eigenvalues. The triangular geometry introduces unique quantization conditions that are distinct from those found in rectangular or circular systems, resulting in a rich and non-trivial energy spectrum.

The symmetry inherent in the equilateral triangle leads to characteristic features in the spatial distribution of the wavefunctions, including specific nodal patterns and degeneracies governed by the triangular point group symmetries. These results highlight the strong influence of geometric confinement on quantum behavior, emphasizing the critical role played by boundary shape in determining physical observables.

Our theoretical findings have important implications in contemporary research areas such as graphene-based quantum dots, nanoscale mechanical resonators, and triangular molecular clusters. The detailed understanding of quantum states in triangular confinements provides a foundation for exploring more complex systems, including the addition of external fields, spin-orbit coupling, and many-body interactions.

Furthermore, the methodology employed here can be extended to study deformed triangles, perturbative effects, and finite potential wells, thereby broadening the relevance of this work. As nanotechnology continues to evolve, the ability to predict and engineer quantum behavior within specific geometries will be vital. Thus, our study not only enriches the fundamental landscape of quantum confinement theories but also serves as a stepping stone toward practical applications in nanoelectronics, quantum optics, and material science. Future research can build upon this framework to model realistic experimental systems, offering deeper insights into quantum phenomena in low-dimensional, symmetry-restricted environments.

References

- 1) Dehesa, J. S., et al., 'Quantum Confinement in Triangular Systems,' *Annals of Physics*, 201, 223-261 (1990).
- 2) Ezawa, M., 'Peculiar Width Dependence of the Electronic Properties of Carbon Nanoribbons,' *Phys. Rev. B*, 73, 045432 (2006).
- 3) Griffiths, D. J., *Introduction to Quantum Mechanics*, 2nd ed., Pearson Prentice Hall (2005).
- 4) Heller, E. J., 'Bound-State Eigenfunctions of Classically Chaotic Hamiltonian Systems: Scars of Periodic Orbits,' *Phys. Rev. Lett.*, 53, 1515 (1984).
- 5) Novoselov, K. S., et al., 'Two-dimensional gas of massless Dirac fermions in graphene,' *Nature*, 438, 197-200 (2005).