


EFFECTS OF MICROORGANISMS IN CROP GROWTH SCIENCES PRODUCTS

A growing collection of research has been conducted on the interactions between humic substances and microorganisms. The importance of a healthy soil microbe population is well-known, but there are many factors that can impact the soil and plant bacterial community. Humic substances can play a positive role in this biological community in a variety of ways. It has been shown that many species of soil bacteria are still capable of growing when their only source of carbon is humic acid¹. In that same experiment, the researchers discovered that humic acids stimulated the growth of bacteria when bacteria were only provided glucose as a growth medium¹. A similar response has been shown regarding toxin-degrading organisms. When humic acid was applied to microbes degrading a toxic environmental contaminant, it increased the enzyme activity of the microbes, which resulted in accelerated microbial remediation².

Since we know that humic acid has a positive effect on microbial communities, it is not surprising that it also contains a large community of beneficial microorganisms on its own. If these microorganisms are provided the right conditions to survive and grow in a humic acid product, the biological effects could be multiplied. Our liquid humic acid and fulvic acid products contain large numbers of plant growth-promoting bacteria.

The species in our liquid humic products are mostly made up of the bacterial groups ***Bacillus***, ***Pseudomonas***, ***Paenibacillus***, and ***Lysinibacillus***. ***Bacillus*** species have been proven to play roles in improving the biomass of plants under heat stress³ and increase plant growth and seed nutrient contents under water stress, especially when paired with humic acid⁴. ***Bacillus*** has also been reported to alleviate the negative effects of salinity stress in plants⁵ and contribute towards hydrocarbon degradation⁶. The benefits of ***Pseudomonas*** include phosphorus solubilization⁷, production of siderophores, which help to limit plant disease⁸, and hydrocarbon degradation in soils⁶. Species in the bacterial group ***Paenibacillus*** can degrade humics in cold environments⁹, dissolve phosphorus, particularly in saline and alkaline soils¹⁰, and when exposed to root-knot nematode eggs and juveniles, it has caused mortality to the plant pathogen¹¹. ***Lysinibacillus*** species have exhibited insecticidal and larvicidal properties, with some species showing antimicrobial characteristics¹². This can contribute towards biological disease control and bioremediation. Many of these studies found that two or more groups of these species often work together to produce these positive effects, so the combination of these organisms in our product is significant.



Hume8 Liquid 12 key species:


Genus	Bacterial count (cfu/ml)
<i>Bacillus</i>	3.50×10^5
<i>Pseudomonas</i>	3.37×10^5
<i>Paenibacillus</i>	2.55×10^5
<i>Lysinibacillus</i>	2.70×10^5



Penetr8 - Low pH Fulvic Acid (pH 5)

Our Penetr8 fulvic product contains a different set of species. The dominant bacterial groups in our liquid fulvic are *Azoarcus*, *Delftia*, and the species *Derxia gummosa*. *Azoarcus* and *Derxia gummosa* are both important nitrogen-fixing bacteria for plants. This gives the organism – and the plants around it – an advantage for survival because they have the capability of transforming nitrogen gas in the atmosphere into nitrogen usable by plants^{13, 14}. *Azoarcus* is also effective at solubilizing insoluble phosphate and producing indoleacetic acid, which is an important plant hormone responsible for regulating plant growth¹⁵. This group has an endophytic lifestyle¹⁵, meaning that they live between living plant cells. They were found adhering internally to rice roots and did not cause plant pathogenic effects when applied to leaves¹⁵. In one study, looking at wheat, a mixture of nitrogen-fixing bacteria including *Azoarcus* was applied to the plants by foliar spraying while nitrogen applications were decreased over two years. The bacteria were discovered to be very effective at adhering to the surface of leaves and roots, and the researchers reported a reduction in nitrogen loss and an increase in the ability of plants to deal with environmental stress¹⁶. The group *Delftia* has shown unique hydrocarbon-degrading capabilities, specifically when humic acid is involved. One study found that the sorption of an environmental contaminant to humic acids caused a limitation to bioavailability that only a few groups of bacteria were able to break, including *Delftia*¹⁷.

Other bacterial groups present in our liquid fulvic play roles in the nitrogen cycle. This includes *Rhodobacter*, which contains species capable of photosynthesis and nitrogen-fixation¹⁸. There are organisms affiliated with the rhizosphere as well, including *Rhizobium* and *Mesorhizobium*. Species in *Rhizobium* are known to be siderophore-producing bacteria, they stimulate nitrogen fixation on root surfaces, and have provided resistance to fungal infection of plants¹⁹. *Mesorhizobium* has been linked to phosphate solubilization in acidic soils²⁰. There are also ammonia-oxidizing bacteria present – *Nitrosomonas* and *Nitrosospira* – and their role in the nitrogen cycle involves turning ammonia oxide into nitrite²¹. *Bacillus* and *Pseudomonas*, described previously as beneficial organisms for soil and plants, are also present in our liquid fulvic.



Penetr8 - Low pH FA key species:

Genus	Bacterial count (cfu/ml)
<i>Azoarcus</i>	5.18×10^6
<i>Delftia</i>	1.70×10^6
<i>Derxia gummosa</i>	1.56×10^6
<i>Rhodobacter</i>	9.56×10^5
<i>Bacillus</i>	5.45×10^5
<i>Pseudomonas</i>	2.83×10^5
<i>Rhizobium</i>	2.10×10^5
<i>Nitrosomonas</i>	1.79×10^5
<i>Nitrospira</i>	1.35×10^5
<i>Mesorhizobium</i>	1.14×10^5



More than just Humic Products

This assortment of microorganisms in our products can provide services including nitrogen fixation, ammonia oxidization, phosphorus solubilization, stress mitigation in plants, biological disease control, and hydrocarbon degradation. Our liquid humic contains nearly one million cfu/ml of these beneficial organisms, and our liquid fulvic contains over ten million cfu/ml of beneficial organisms. The microbial community in our products have occurred because the species present in the raw material have been given an environment that allows them to multiply through our liquid manufacturing process. There are no external species added, so we can be confident that the species present can live and function in equilibrium. These numbers are in line with many liquid microbial products currently on the market, but the addition of humic acid in our product also supports bacterial life in the soils and plants where it is being applied.



References:

1. Tikhonov VV, Yakushev AV, Zavgorodnyaya YA, Byzov BA, Demin VV. 2010. Effect of humic acids on the growth of bacteria. *Eurasian Soil Sc.* 43(3):305-313.
2. Wan J, Chen K, Chen J, Qin Z, Adrian L, Shen C. 2022. Enhanced perchloroethene dichlorination by humic acids via increasing the dehalogenase activity of Dehalococcoides strains. *FEMS Microbiol Ecol.* 98(4):1-9.
3. Khan MA, Asaf S, Khan AL, Jan R, Kang S, Kim K, Lee I. 2020. Extending thermotolerance to tomato seedlings by inoculation with SA1 isolate of *Bacillus cereus* and comparison with exogenous humic acid application. *PLOS ONE.* 15(4):e0232228.
4. Ekin Z. 2020. Field evaluation of rhizobacterial inoculants in combination with humic substances to improve seed and oil yields of safflower (*Carthamus tinctorius* L.) under irrigated and rainfed conditions. *Appl Ecol Environ Res.* 18(5):6377-6403.
5. Esringu A, Kaynar D, Turan M, Ercisli S. 2015. Ameliorative effect of humic acid and plant growth-promoting rhizobacteria (PGPR) on Hungarian Vetch plants under salinity stress. *Commun Soil Sci Plant Anal.* 47(5):602-618.
6. Kubota K, Daisuke K, Matsumiya Y, Chung S, Kubo M. 2008. Phylogenetic analysis of long-chain hydrocarbon-degrading bacteria and evaluation of their hydrocarbon-degradation by the 2,6-DCPIP assay. *Biodegradation.* 19:749-757.
7. Govindan S, Joshi P, Nazim S, Mishra PK, Bisht JK, Gupta HS. 2009. Phosphate solubilization and growth promotion by *Pseudomonas fragi* CS11RH1 (MTCC 8984), a psychrotolerant bacterium isolated from a high altitude Himalayan rhizosphere. *Biologia.* 64(2):239-245.
8. Gupta A, Gopal M. 2008. Siderophore production by plant growth promoting rhizobacteria. *Indian J Agric Res.* 42(2):153-156.
9. Park HJ, Kim D. 2013. Draft genome sequence of a humic substance-degrading *Paenibacillus* sp. isolated from the subarctic grasslands at low temperature. *Genome Announc.* 1(1).
10. Zhang Y, Lang L, Sun Z, Li M. 2021. Potential application of *Paenibacillus* sp. C1 to the amelioration of soda saline-alkaline soil. *Geomicrobiol.* 40(2):172-182.
11. Khan Z, Kim SG, Jeon YH, Khan HU, Son SH, Kim YH. 2008. A plant growth promoting rhizobacterium, *Paenibacillus polymyxa* strain GBR-1, suppresses root-knot nematode. *Bioresour Technol.* 99(8):3016-3023.
12. Jamal QMS, Ahmad V. 2022. Lysinibacilli: A biological factories intended for bio-insecticidal, bio-control, and bioremediation activities. *J Fungi.* 8(12):1288.
13. Martin-Moldes Z, Zamarro MT, del Cerro C, Valencia A, Gomez MJ, Arcas A, Udaondo Z, Garcia JL, Nogales J, Carmona M, Diaz E. 2015. Whole-genome analysis of *Azoarcus* sp. strain CIB provides genetic insights to its different lifestyles and predicts novel metabolic features. *Syst Appl Microbiol.* 38(7):462-471.
14. Jensen HL, Petersen EJ, De PK, Bhattacharya R. 1960. A new nitrogen-fixing bacterium: *Derxia gummosa* nov. gen. nov. spec. *Arch Microbiol.* 36(2): 182-195.
15. Fernandez H, Prandoni N, Fernandez-Pascual M, Fajardo S, Morcillo C, Diaz E, Carmona M. 2014. *Azoarcus* sp. CIB, an anaerobic biodegrader of aromatic compounds shows an endophytic lifestyle. *PLOS ONE.* 9(10):e110771.
16. Cortivo CD, Barion G, Visioli G, Mattarozzi M, Mosca G, Vamerali T. 2017. Increased root growth and nitrogen accumulation in common wheat following PGPR inoculation: Assessment of plant-microbe interactions by ESEM. *Agric Ecosyst Environ.* 247:396-408.
17. Vacca DJ, Bleam WF, Hickey WJ. 2005. Isolation of soil bacteria adapted to degrade humic acid-sorbed phenanthrene. *Appl Environ Microbiol.* 71(7):3797-3805.
18. Kern M, Kamp P, Paschen A, Masepohl B, Klipp W. 1998. Evidence for a regulatory link of nitrogen fixation and photosynthesis in *Rhodobacter capsulatus* via HvrA. *J Bacteriol.* 180(7):1965-1969.
19. Jaiswal SK, Mohammed M, Ibny FYI, Dakora FD. 2021. Rhizobia as a source of plant growth-promoting molecules: potential applications and possible operational mechanisms. *Front Sustain Food Syst.* 4:619676.
20. Muleta A, Tesfaye K, Selassie THH, Cook DR, Assefa F. 2021. Phosphate solubilization and multiple plant growth promoting properties of *Mesorhizobium* species nodulating chickpea from acidic soils of Ethiopia. *Arch Microbiol.* 203:2129-2137.
21. Zorz JK, Kozłowski JA, Stein LY, Strous M, Kleiner M. 2018. Comparative proteomics of three species of ammonia-oxidizing bacteria. *Front Microbiol.* 9.