

CATARACT SURGERY EYE DROP INSTRUCTIONS



1 – Tan

BESIVANCE (or Moxifloxacin / Ciprofloxacin)

Start 3 days before surgery. 4 times per day. Stop 1 week after Surgery.

2 – Gray

BROMSITE (or Ketorolac)

Start 3 days before surgery. 2 times per day. Stop 4 weeks after Surgery.

3 – Pink or White

DUREZOL (or Prednisolone)

Start AFTER Surgery.

4 times a day for 1 week, then decrease to
3 times a day for 1 week, then decrease to
2 times a day for 1 week, then decrease to
1 time a day for 1 week, then stop

WAIT 5 MINUTES IN BETWEEN EACH DROP
DO NOT RUB YOUR EYE FOR 1 WEEK AFTER SURGERY

PLEASE BRING YOUR EYE DROPS TO ALL APPOINTMENTS AFTER SURGERY

3 Days Prior	2 Days Prior	1 Day Prior
Besivance 	Besivance 	Besivance
Bromsite 	Bromsite 	Bromsite

[illegible]

WEEK 2	Day 8	Day 9	Day 10	Day 11	Day 12	Day 13	Day 14
	Bromsite 	Bromsite 	Bromsite 	Bromsite 	Bromsite 	Bromsite 	Bromsite
	Durezol 	Durezol 	Durezol 	Durezol 	Durezol 	Durezol 	Durezol

WEEK 3	Day 15	Day 16	Day 17	Day 18	Day 19	Day 20	Day 21
	Bromsite 	Bromsite 	Bromsite 	Bromsite 	Bromsite 	Bromsite 	Bromsite 
	Durezol 	Durezol 	Durezol 	Durezol 	Durezol 	Durezol 	Durezol 

WEEK 4	Day 22	Day 23	Day 24	Day 25	Day 26	Day 27	Day 28
	Bromsite 	Bromsite 	Bromsite 	Bromsite 	Bromsite 	Bromsite 	Bromsite 
	Durezol 	Durezol 	Durezol 	Durezol 	Durezol 	Durezol 	Durezol 