



BEATRIX
EICHERT
PSYCHOLOGISCHE
PSYCHOTHERAPEUTIN

ASD diagnosis for enhanced clarity and improved self-awareness.

You may also relate to this: the sensation of being distinct, finding social interactions draining, experiencing sensory overload, or possessing a profound need for structure and solitude. Numerous individuals affected by this condition often feel misinterpreted and perpetually fatigued.

Particularly among women, autism frequently remains unrecognized for extended periods due to their ability to adapt and conform to societal expectations. A comprehensive diagnosis can facilitate a deeper understanding of their experiences, promote self-compassion, and foster the development of new perspectives.



Private practice for psychological therapy in Weimar



ASD diagnosis offered as a service for self-paying patients, ensuring minimal waiting times.

In my practice, I provide comprehensive counseling and psychotherapy grounded in the latest scientific research, ensuring you receive optimal support. My team of seasoned professionals is available to address all your inquiries and facilitate a diagnostic process that is straightforward and stress-free.

It is of utmost importance to me to offer you access to high-quality diagnostics and evidence-based treatment.

Your mental health is in the most capable hands.




to deliver psychotherapeutic care without extended waiting periods.

I firmly believe that every individual is entitled to a prompt and comprehensive diagnosis. Consequently, I have streamlined the process to ensure you receive the necessary support without delay.

You are welcome to schedule an initial consultation to address your inquiries regarding ND diagnostics.

praxis@psychotherapie-eichert.de
www.psychotherapie-eichert.de

Welcome to my private practice.



You may also be acquainted with this: challenges in concentration, a sense of inner restlessness, forgetfulness, or the impression of perpetually underachieving. Numerous individuals experience self-doubt and fatigue.

ADHD is frequently underestimated, particularly in women, as they often mask their symptoms for extended periods, presenting themselves as highly competent. An accurate diagnosis can facilitate a deeper understanding of their experiences and open new avenues for progress.

Schedule your upcoming appointments effortlessly online.



ASD Diagnostics - Protocol

Initial consultation (anamnesis): A comprehensive interview regarding your development, current challenges, social environment, and potential comorbidities.



Documents and Questionnaires:

You will receive several questionnaires, often including those for close relatives who are familiar with your childhood. Please return these completed for evaluation, along with any available old school reports.



Diagnostic Interview:

Implementing standardized testing protocols while examining ASD symptoms in the context of your personal history.



Assessment meeting:

A comprehensive analysis of the test results along with a recommendation regarding the advisability of psychotherapeutic treatment.



Ultimately, you will obtain a written summary of the test results.

Important: To prevent the oversight of any additional mental illnesses that may have arisen, their symptoms are also documented.

Which symptoms are most prevalent in an ASD diagnosis?

Social perception challenges, social Recognizing signals, insecurity in communication, sense of being "different."

2. Misunderstandings in Communication Dialogue, literal comprehension, challenges with casual conversation or subtle cues

3. Perception and Stimulus Processing: Elevated Sensitivity in relation to auditory, visual, tactile, or other sensory stimuli

4. Framework & Procedures A strong desire for order and predictability, accompanied by stress induced by changes or uncertainty in daily life.

5. Childhood abnormalities: Were there any early signs? For instance, withdrawal, pronounced special interests, specific sensitivities, or social anxieties?

praxis@psychotherapie-eichert.de
www.psychotherapie-eichert.de